



US00PP18693P2

(12) **United States Plant Patent**  
**Bradford**

(10) **Patent No.:** **US PP18,693 P2**  
(45) **Date of Patent:** **Apr. 1, 2008**

(54) **NECTARINE TREE NAMED ‘SUGARRED I’**

(56) **References Cited**

(50) Latin Name: *Prunus persica*  
Varietal Denomination: **SUGARRED I**

U.S. PATENT DOCUMENTS

PP7,507 P \* 4/1991 Bradford et al.

(76) Inventor: **Lowell Glen Bradford**, 10237 E.  
Mariposa Way, Le Grand, CA (US)  
95333

\* cited by examiner

*Primary Examiner*—Kent Bell

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(57) **ABSTRACT**

The present invention relates to a new and distinct variety of  
nectarine tree, *Prunus persica*, broadly characterized by a  
medium size, moderately vigorous, hardy, self-fertile, pro-  
ductive and regular bearing tree. The fruit matures under the  
ecological conditions described in mid June, with first  
picking on Jun. 19, 2006. The fruit is uniformly large in size,  
mildly acidic and sweet in flavor, globose in shape, cling-  
stone in type, very firm in texture, yellow with strong red  
bleeding in flesh color, and full red in skin color.

(21) Appl. No.: **11/637,894**

(22) Filed: **Dec. 13, 2006**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(52) **U.S. Cl.** ..... **Plt./187**

(58) **Field of Classification Search** ..... Plt./187  
See application file for complete search history.

**1 Drawing Sheet**

**1**

Botanical classification: *Prunus persica*.  
Variety denomination: ‘Sugarred I’.

**BACKGROUND OF THE VARIETY**

In a continuing effort to improve the quality of shipping  
fruits, I, the inventor, typically hybridize a large number of  
peach, nectarine, plum, apricot, and cherry seedlings each  
year. The present invention relates to a new and distinct  
variety of nectarine tree, which has been denominated  
varietally as ‘SUGARRED I’. The present variety was  
hybridized by me in 2000, grown as a seedling on its own  
root in my greenhouse, and transplanted to a cultivated area  
of my experimental orchard located near Le Grand, Calif., in  
Merced County (San Joaquin Valley).

The variety was developed as a first generation cross  
using ‘Spring Bright’ (U.S. Plant Pat. No. 7,507), nectarine  
as the selected seed parent and an unnamed nectarine  
seedling as the selected pollen parent. A single tree from the  
stated cross was selected as the claimed variety. Subsequent  
to origination of the present variety of nectarine tree, I  
asexually reproduced it by budding and grafting in the  
experimental orchard described above, and such reproduc-  
tion of plant and fruit characteristics were true to the original  
plant in all respects. The reproduction of the variety included  
the use of ‘Nemaguard’ (unpatented) rootstock upon which  
the present variety was compatible and true to type.

The present variety is most similar to its selected seed  
parent, ‘Spring Bright’ nectarine, by producing nectarines  
that are nearly globose in shape, acidic and sweet in flavor,  
and full red in skin color, but is distinguished therefrom by  
producing nectarines that are yellow with much more red  
bleeding in flesh color, that are larger in size, and that mature  
about five days earlier.

**SUMMARY OF VARIETY**

In summary, the present variety is characterized by a  
medium size, moderately vigorous, hardy, self-fertile, pro-

**2**

ductive and regular bearing tree. The fruit matures under the  
ecological conditions described in mid June, with first  
picking on Jun. 19, 2006. The fruit is uniformly large in size,  
mildly acidic and sweet in flavor, globose in shape, cling-  
stone in type, very firm in texture, yellow with strong red  
bleeding in flesh color, and full red in skin color.

**DRAWING**

The accompanying photograph consists of four whole  
fruits positioned to display the characteristics of the skin  
color and form, one fruit divided transversely to the suture  
plane to reveal the flesh and stone, two insets to reveal buds  
and blossoms in various stages, various leaves, a clean dry  
stone, and a typical tip shoot.

**POMOLOGICAL CHARACTERISTICS**

Referring now more specifically to the pomological char-  
acteristics of this new and distinct variety of nectarine tree,  
the following has been observed under the ecological con-  
ditions prevailing near Le Grand, Merced County (San  
Joaquin Valley), Calif., and was developed at the state of  
firm ripe on Jun. 23, 2006, on the original tree during its  
sixth growing season. All major color code designations are  
by reference to the Inter-Society Color Council, National  
Bureau of Standards. Common color names are also used  
occasionally.

**Tree**

Size: Medium, reaching and maintaining a height of 8' [2.44  
m.] and a spread of 7' [2.13 m.] after six growing seasons  
utilizing typical dormant pruning.

Vigor: Moderate, responding typically to irrigation and  
fertilization. The variety grows about 3' [0.91 m.] of  
surplus top-growth during the spring and summer. The  
plant should be grown on a standard commercial root-  
stock for production purposes.



Growth: Spreading and dense.

Form: Pruned to a perpendicular "V".

Hardiness: Hardy with respect to central California winters.

Heat tolerance: Observed to perform adequately in typical central California climatic conditions, which typically include extended periods of heat.

Drought tolerance: Variety is developed for commercial orchards and requires regular irrigation.

Production: Productive, thinning necessary.

Fertility: Self-fertile.

Bearing: Regular bearer with no alternate bearing yet observed.

Approximate chilling requirement: 525 hours.

Trunk:

*Size*.—Medium, reaching a maximum diameter of 3½" [89 mm.] after the sixth growing season.

*Texture*.—Shaggy.

*Bark color*.—A Grayish olive [110. gy.Ol] and Dark brown [59. d.Br] variegation.

*Lenticels*.—Approximate Number Per Square Inch: 8.

*Color*: Moderate orange yellow [71. m.OY]. Typical

*Size*: ¾" [4.8 mm.] to 7/16" [11.1 mm.]. *Shape*:

Eye-shaped to elongated.

Branches:

*Size*.—Diameter of limb is 2⅛" [54 mm.] measured 12" above the crotch, 1⅝" [33 mm.] measured 12" above the first fork.

*Texture*.—Smooth on first and second year wood, increasing roughness with age.

*Color*.—1st Year Wood Topside: Grayish red [19. gy.R]. 1st Year Wood Underside: Brilliant yellow green [116. brill.YG]. Older Wood: Deep yellowish brown [75. deep yBr].

*Lenticels*.—Number Per Square Inch: More than 60 on second year wood. *Color*: Light yellowish brown [76. l.yBr]. Typical *size*: ¼" [0.4 mm.] to ⅜" [2.4 mm.] on second year wood. *Shape*: Elongated.

Leaves:

*Size*.—Medium. Average Length: 5⅜" [137 mm.]. Average Width: 1½" [38 mm.].

*Arrangement*.—Alternate.

*Thickness*.—Medium.

*Form*.—Elliptical.

*Apex*.—Acuminate.

*Base*.—Rounded to acute.

*Surface*.—Smooth.

*Color*.—Dorsal Surface: Moderate olive green [125. m.OlG]. Ventral Surface: Moderate yellow green [120. m.YG].

*Margin*.—Finely serrate.

*Venation*.—Pinnately net veined.

*Vein color*.—Light greenish yellow [101. l.gY].

*Petiole*.—Average Length: ¾" [9.5 mm.]. Average Thickness: 1/16" [1.6 mm.]. *Color*: Moderate yellow green [120. m.YG].

*Stipules*.—Number: 2 per leaf, up to 6 per growing tip. Average Length: ¾" [9.5 mm.]. *Color*: Brilliant yellow green [116. brill.YG] becoming Moderate reddish brown [43. m.rBr] with maturity.

*Glands*.—Number: Usually 2 per leaf. Position: Slightly alternate, positioned at the intersection of the petiole and the base of the leaf. *Size*: Small. *Form*: Globose. *Color*: Strong yellow green [117. s.YG] becoming Moderate yellow brown [77. m.yBr] with age.

*Leaf buds*.—Pointed, medium in size.

Flower buds:

*Hardiness*.—Hardy, with respect to central California winters.

*Diameter*.—Typically ⅜" [9.5 mm.] 1 week before bloom.

*Length*.—Typically 1¼" [17.5 mm.] 1 week before bloom.

*Form*.—Not appressed.

*Surface*.—Pubescent.

*Color*.—Light purplish pink [249. l.pPk].

Flowers: Perfect, complete, perigynous, usually a single pistil, typically thirty or more stamens, five and sepal and petal locations alternately positioned.

*Type*.—Showy, very large.

*Average flower diameter*.—2" [50.8 mm.].

*Number of petals*.—Usually five, some with extra petal fragments and some double blossoms observed.

*Petal shape*.—Circular to oval.

*Petal margin*.—Somewhat wavy.

*Average petal diameter*.—13/16" [20.6 mm.].

*Average petal length*.—13/16" [20.6 mm.].

*Petal apex*.—Rounded.

*Petal base*.—Rounded to somewhat truncate.

*Petal color*.—Pale pink [7. p.Pk] toward the apex, Light pink [4. l.Pk] toward the base.

*Anther color*.—Dark red [16. d.R] over a Light yellow [86. l.Y] center at bloom onset.

*Stigma color*.—Light greenish yellow [101. l.gY].

*Sepal color*.—Dark purplish red [259. d.pR].

*Sepal length*.—5/16" [7.9 mm.].

*Sepal width*.—¼" [6.4 mm.].

*Average pistil length*.—13/16" [20.6 mm.].

*Average stamen length*.—11/16" [17.5 mm.].

*Fragrance*.—Moderate.

*Blooming period*.—Medium, close to 'Spring Bright' (U.S. Plant Pat. No. 7,507) nectarine.

*Onset of bloom*.—One percent on Feb. 23, 2006.

*Date of full bloom*.—Mar. 8, 2006.

*Duration of bloom*.—One to two weeks, dependent on ambient temperature.

*Number per cluster*.—1 to 3 with single flowers most common.

## FRUIT

Maturity when described: Firm ripe, Jun. 23, 2006.

Date of first picking: Jun. 19, 2006.

Date of last picking: Jun. 29, 2006.

*Size*: Uniform, large.

*Average diameter axially*.—3¼" [78 mm.].

*Average diameter across check plane*.—2⅝" [75 mm.].

*Average diameter across cheek plane*.—3¼" [78 m.].

*Typical weight*.—8.2 ounces [232 grams].

*Form*: Uniform, globose, slightly asymmetrical.

*Longitudinal section form*.—Round to elliptical.

*Axial view*.—Round with a slight hump at the suture.

*Suture*: A very sharp groove near the base, a shallow rough along the side, and a sharp groove toward the apex, ending just beyond the pistil point with a moderate depression.

*Ventral surface*: Rounded, lipped stronger toward the apex.

*Lips*: Slightly unequal.

*Cavity*: Flaring, circular, suture usually showing on both sides, Strong orange yellow [68. s.OY] stem markings typical.

*Depth*.—½" [12.7 mm.].

*Breadth*.—1¼" [27.0 mm.].

Base: Truncate.  
 Apex: Rounded.  
 Pistil point: Both apical and oblique, negligible in length, depressed within the suture.  
 Stem: Medium.  
*Average length.*— $\frac{3}{8}$ " [9.5 mm.].  
*Average width.*— $\frac{3}{16}$ " [4.8 mm.].  
 Skin:  
*Thickness.*—Medium.  
*Surface.*—Smooth.  
*Tenacity.*—Tenacious to flesh.  
*Astringency.*—Moderately astringent.  
*Tendency to crack.*—Slight.  
*Color.*—Dark red [16. d.R] smoothly blending into a Strong reddish orange [35. s.rO] background with the slightest amount of Pale orange yellow [73. p.OY] freckling.  
 Flesh:  
*Color.*—Deep red [13. deep R] toward the skin finally blending into Brilliant orange yellow [67. brill.OY] close to the stone.  
*Surface of pit cavity.*—Deep red [13. deep R] broken fibers when twisted from stone.  
*Amygdalin.*—Abundant.  
*Juice.*—Abundant, rich.  
*Texture.*—Very firm, crisp.  
*Fibers.*—Abundant, fine.  
*Ripens.*—Slightly earlier at the apex.  
*Flavor.*—Acidic and sweet, typically 15 brix.  
*Aroma.*—Moderate.  
*Eating quality.*—Excellent.

## STONE

Type: Clingstone.  
 Form: Oval.  
 Hilum: Oblong.  
 Base: Straight oblique.  
 Apex: Rounded to obtuse with an average angle of 110 degrees.  
 Sides: Equal.  
 Surface: Regularly furrowed toward the apex, pitted toward the base.  
 Ridges: Jagged.  
 External color: Moderate yellowish brown [77. m.yBr] when dry.  
 Pit wall color when cracked: Light yellowish brown [76. l.yBr].

Cavity surface color: Deep brown [56. deep Br].  
 Average pit wall thickness:  $\frac{1}{4}$ " [6.4 mm.].  
 Average width:  $1\frac{7}{16}$ " [36.5 mm.].  
 Average length:  $1\frac{1}{8}$ " [28.6 mm.].  
 Average breadth:  $\frac{3}{4}$ " [19.1 mm.].  
 Tendency to split: Slight.  
 Kernel:  
*Form.*—Oval.  
*Skin color.*—Strong yellowish brown [74. s.yBr] when dry.  
*Pellicle color.*—Grayish brown [61. gy.Br].  
*Vein color.*—Dark brown [59. d.Br].  
*Taste.*—Sweet.  
*Viable.*—Yes.  
*Average width.*— $\frac{9}{16}$ " [14.3 mm.].  
*Average length.*— $\frac{3}{4}$ " [19.1 mm.].  
*Amygdalin.*—Moderate.

## USE

Market: Fresh market and long distance shipping.  
 Keeping quality: Good, fruit quality observed to remain in good condition after 21 days in standard cold room at 36° Fahrenheit [2° Celsius].  
 Shipping quality: Good.  
 Resistance to insects: No unusual susceptibilities noted.  
 Resistance to diseases: No unusual susceptibilities noted.

## Other Notes

Although the new variety of nectarine tree possesses the described characteristics under the ecological conditions at Le Grand, Calif., in the central part of the San Joaquin Valley, It is to be expected that variations in these characteristics may occur when farmed in areas with different climatic conditions, different soil types, and/or varying cultural practices.

I claim:

1. A new and distinct variety of nectarine tree, substantially as illustrated and described, that is most similar to its selected seed parent, 'Spring Bright' (U.S. Plant Pat. No. 7,507) nectarine, by producing nectarines that are nearly globose in shape, acidic and sweet in flavor, and full red in skin color, but is distinguished therefrom by producing nectarines that are yellow with much more red bleeding in flesh color, that are larger in size, and that mature about five days earlier.

\* \* \* \* \*



