



(12) **United States Plant Patent**
Ren

(10) **Patent No.:** **US PP18,777 P2**
(45) **Date of Patent:** **Apr. 29, 2008**

(54) **CALIBRACHOA PLANT NAMED
'BALCABWITIM'**

(50) Latin Name: ***Calibrachoa*×*hybrida***
Varietal Denomination: **Balcabwitim**

(75) Inventor: **Jianping Ren**, Geneva, IL (US)

(73) Assignee: **Ball Horticultural Company**, West
Chicago, IL (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/602,698**

(22) Filed: **Nov. 20, 2006**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./263**

(58) **Field of Classification Search** Plt./263
See application file for complete search history.

Primary Examiner—Kent Bell

Assistant Examiner—June Hwu

(74) *Attorney, Agent, or Firm*—Audrey Charles

(57) **ABSTRACT**

A new and distinct cultivar of *Calibrachoa* plant named
'Balcabwitim', characterized by its white-colored flowers,
medium green-colored foliage, and moderately vigorous,
mounded-trailing growth habit.

1 Drawing Sheet

1

Latin name of genus and species of plant claimed: *Calibrachoa*×*hybrida*.

Variety denomination: 'Balcabwitim'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Calibrachoa* plant botanically known as *Calibrachoa*×*hybrida* and hereinafter referred to by the cultivar name 'Balcabwitim'.

The new cultivar originated in a controlled breeding program in Elburn, Ill. during November 2002. The objective of the breeding program was the development of *Calibrachoa* cultivars with attractive flower coloration, medium green-colored foliage, and a moderately vigorous, mounded-trailing growth habit.

The new *Calibrachoa* cultivar is the result of cross-pollination. The female (seed) parent of the new cultivar is the proprietary *Calibrachoa*×*hybrida* breeding selection designated 1023-2, not patented, characterized by its rose-colored flowers, medium green-colored foliage, and trailing growth habit. The male (pollen) parent of the new cultivar is a *Calibrachoa*×*hybrida* mixed pollen source from plants having various colored flowers, medium to dark green-colored foliage, and prostrate to mounded growth habits. The new cultivar was discovered and selected as a single flowering plant within the progeny of the above stated cross-pollination during June 2004 in a controlled environment at Elburn, Ill.

Asexual reproduction of the new cultivar by terminal stem cuttings since June 2004 at Arroyo Grande, Calif. and West Chicago, Ill. has demonstrated that the new cultivar reproduces true to type with all of the characteristics, as herein described, firmly fixed and retained through successive generations of such asexual propagation.

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have been repeatedly observed and can be used to distinguish 'Balcabwitim' as a new and distinct cultivar of *Calibrachoa* plant:

1. White-colored flowers;
2. Medium green-colored foliage; and

2

3. Moderately vigorous, mounded-trailing growth habit.
Plants of the new cultivar differ from plants of the female parent primarily in flower color and growth habit.

Of the many commercially available *Calibrachoa* cultivars known to the inventor, the most similar in comparison of the new cultivar is Callie White 'Cal White', U.S. Plant Pat. No. 14,497. However, in side by side comparisons, plants of the new cultivar differ from plants of 'Cal White' in the following characteristics:

1. Plants of the new cultivar are narrower than plants of 'Cal White'; and
2. Plants of the new cultivar are better branched than plants of 'Cal White'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it is reasonably possible to make the same in color illustrations of this type, typical flower and foliage characteristics of the new cultivar. Colors in the photographs differ slightly from the color values cited in the detailed description, which accurately describes the colors of 'Balcabwitim'. The plants were grown in 4.5 inch pots for 12 weeks in a greenhouse at West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and flowering habit of 'Balcabwitim'.

FIG. 2 illustrates a close-up view of an individual flower of 'Balcabwitim'.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible environmental conditions to date. Accordingly, it is possible that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2001 edition, except where general color terms of ordinary significance are used. The color values were determined on Feb. 23, 2006 between 9:00

a.m. and 11:00 a.m. under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe plants produced from cuttings taken from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown at West Chicago, Ill. in 4.5 inch pots for 12 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 70° F. 77° F. (21° C. to 25° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Greenhouse light levels of 2,500 footcandles to 6,000 footcandles were maintained during the day.

Botanical classification: *Calibrachoa*×*hybrida* cultivar Balcabwitim.

Parentage:

Female parent.—Proprietary *Calibrachoa*×*hybrida* breeding selection designated 1023-2, not patented.

Male parent.—*Calibrachoa*×*hybrida* mixed pollen source.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 6 to 8 days.

Time to produce a rooted cutting.—Approximately 24 to 28 days.

Root description.—Fibrous.

Rooting habit.—Freely branching.

Plant description:

Commercial crop time.—Approximately 6 to 7 weeks from a rooted cutting.

Growth habit and general appearance.—Moderately vigorous, trailing, becoming mounded with age.

Size.—Height from soil level to top of plant plane: Approximately 7.9 cm. Width: Approximately 40.6 cm.

Branching habit.—Freely branching. Quantity of main branches per plant: Approximately 6.

Branch.—Strength: Strong. Length: Approximately 22.7 cm. Diameter: Approximately 2.0 mm. Length of central internode: Approximately 1.5 cm. Texture: Densely glandular pubescent. Color of young stem: 144C. Color of mature stem: 146D.

Foliage description:

General description.—Quantity of leaves per main branch: Approximately 26. Fragrance: Slight. Form: Simple. Arrangement on flowering stem: Opposite.

Leaves.—Aspect: At acute angle to stem with tip turning downward. Shape: Elliptic. Margin: Entire. Apex: Acute. Base: Attenuate. Venation pattern: Pinnate. Length of mature leaf: Approximately 3.2 cm. Width of mature leaf: Approximately 1.0 cm. Texture of upper and lower surfaces: Glandular pubescent. Color of upper surface of young foliage: 137C with venation of 144D. Color of lower surface of young foliage: 138C with venation of 144D. Color of upper surface of mature foliage: 137B with venation of 144D. Color of lower surface of mature foliage: 138B with venation of 144D.

Petiole.—Length: Approximately 2.8 cm. Diameter: Approximately 1.0 mm. Texture: Glandular pubescent. Color: 144D.

Flowering description:

Flowering habit.—‘Balcabwitim’ is freely flowering under outdoor growing conditions with substantially

continuous blooming from spring through autumn and year-round in greenhouse environment.

Lastingness of individual flower on the plant.—Approximately 5 to 6 days.

Flower description:

General description.—Type: Single, salverform. Quantity per plant: Approximately 44. Fragrance: None.

Bud.—Rate of opening: Generally takes 3 to 4 days for bud to progress from first color to fully open flower. Quantity per plant: Approximately 9.

Bud just before opening.—Shape: Oblong. Length: Approximately 1.7 cm. Diameter: Approximately 4.5 mm. Texture: Densely glandular pubescent. Color: Petals 4D and corolla tube 160C both with venation of 151D.

Corolla.—Diameter: Approximately 2.8 cm.

Petals.—Quantity: 5, fused to form a tube. Shape: Obovate. Margin: Entire. Apex: Obtuse, some acuminate. Length from tube: Approximately 1.1 cm. Length of free portion: Approximately 4.7 mm. Width: Approximately 1.2 cm. Texture of upper surface: Glabrous. Texture of lower surface: Glandular pubescent, dense along the veins. Color of all surfaces when first open and when fully open: 155D with midveins of N144A.

Corolla tube.—Length: Approximately 1.5 cm. Diameter at distal end: Approximately 6.0 mm. Diameter at proximal end: Approximately 1.0 mm. Texture of inner surface: Glabrous. Texture of outer surface: Glandular pubescent. Color of inner surface: 9A with venation of N144A. Color of outer surface: 160D with venation of 151D.

Peduncle.—Strength: Strong. Aspect: Acute angle to stem. Length: Approximately 1.2 cm. Diameter: Approximately 1.0 mm. Texture: Densely glandular pubescent. Color: 144C.

Sepals.—Quantity per flower: 5, fused along lower half. Shape: Lanceolate. Apex: Acute. Length: Approximately 1.1 cm. Width: Approximately 2.0 mm. Texture of upper and lower surfaces: Densely glandular pubescent. Color of upper surface: 138A. Color of lower surface: 138B with 144C at base.

Reproductive organs.—Androecium: Stamen quantity: 5 per flower, fused to inside of corolla tube. Stamen length: Approximately 1.1 cm. Filament length of free portion: Approximately 6.4 mm. Filament color: 145D. Anther shape: 2 parted, oval. Anther length: Approximately 0.5 mm. Anther color: 10A. Pollen amount: Moderate. Pollen color: 13B. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 9.6 mm. Stigma shape: Funnel. Stigma length: Approximately 0.5 mm. Stigma color: 144A. Style length: Approximately 7.6 mm. Style color: 145D. Ovary diameter: Approximately 1.5 mm. Ovary color: N144B.

Seed and fruit production: Neither seed nor fruit production has been observed.

Disease and petal resistance: Resistance to pathogens and pests common to *Calibrachoa* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Calibrachoa* plant named ‘Balcabwitim’, substantially as herein shown and described.

* * * * *



FIG. 1




FIG. 2