

(12) **United States Plant Patent**
Meyer

(10) **Patent No.:** **US PP18,852 P2**
(45) **Date of Patent:** **May 27, 2008**

(54) **GUINEA *IMPATIENS* PLANT NAMED
'BALCELIMPUR'**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(50) Latin Name: ***Impatiens hawkeri***
Varietal Denomination: **Balcelimpur**

(52) **U.S. Cl.** **Plt./318**

(58) **Field of Classification Search** Plt./318
See application file for complete search history.

(75) Inventor: **Kerry M. Meyer**, St. Thomas, MO
(US)

Primary Examiner—Annette H Para
(74) *Attorney, Agent, or Firm*—Audrey Charles

(73) Assignee: **Ball Horticultural Company**, West
Chicago, IL (US)

(57) **ABSTRACT**

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 67 days.

A new and distinct cultivar of New Guinea *Impatiens* plant
named 'Balcelimpur', characterized by its single type
medium purple-colored flowers, medium green-colored
foliage, and vigorous, upright-mounded growth habit.

(21) Appl. No.: **11/602,815**

1 Drawing Sheet

(22) Filed: **Nov. 21, 2006**

1

Latin name of genus and species of plant claimed: *Impa-
tiens hawkeri*.
Variety denomination: 'Balcelimpur'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of New Guinea *Impatiens* plant botanically known as *Impa-
tiens hawkeri* and hereinafter referred to by the cultivar
name 'Balcelimpur'.

The new cultivar originated in a controlled breeding
program in Arroyo Grande, Calif. during May 2002. The
objective of the breeding program was the development of
New Guinea *Impatiens* cultivars with numerous large
flowers, attractive flower coloration, and a freely basal
branching growth habit.

The new New Guinea *Impatiens* cultivar is the result of
cross-pollination. The female (seed) parent of the new
cultivar is Harmony Lavender Purple 'Danharpl', U.S. Plant
Pat. No. 12,094, characterized by its single type medium
red-purple-colored flowers, dark green-colored foliage, and
compact growth habit. The male (pollen) parent of the new
cultivar is Celebration 'Raspberry Rose', U.S. Plant Pat. No.
9,212, characterized by its single type dark rose-colored
flowers, medium green-colored foliage, and upright growth
habit. The new cultivar was discovered and selected as a
single flowering plant within the progeny of the above stated
cross-pollination during October 2002 in a controlled envi-
ronment at Arroyo Grande, Calif.

Asexual reproduction of the new cultivar by terminal stem
cuttings since October 2002 at Arroyo Grande, Calif. and
West Chicago, Ill. has demonstrated that the new cultivar
reproduces true to type with all of the characteristics, as
herein described, firmly fixed and retained through succes-
sive generations of such asexual propagation.

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have
been repeatedly observed and can be used to distinguish
'Balcelimpur' as a new and distinct cultivar of New Guinea
Impatiens plant:

2

1. Single type medium purple-colored flowers;
2. Medium green-colored foliage; and
3. Vigorous, upright-mounded growth habit.

Plants of the new cultivar differ from plants of the female
parent primarily in flower color and foliage color, and from
plants of the male parent primarily in flower color and
foliage color.

Of the many commercially available New Guinea *Impa-
tiens* cultivars known to the inventor, the most similar in
comparison to the new cultivar is Celebration Purple 'BFP-
661 Purple', U.S. Plant Pat. No. 10,092. However, in side by
side comparisons, plants of the new cultivar differ from
plants of 'BFP-661 Purple' in the following characteristics:

1. Plants of the new cultivar have fewer flowers than
plants of 'BFP-661 Purple'; and
2. Plants of the new cultivar have larger flowers than
plants of 'BFP-661 Purple'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it
is reasonably possible to make the same in color illustrations
of this type, typical flower and foliage characteristics of the
new cultivar. Colors in the photographs differ slightly from
the color values cited in the detailed description, which
accurately describes the colors of 'Balcelimpur'. The plants
were grown in 4.5 inch pots for 12 weeks in a greenhouse
at West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and
flowering habit of 'Balcelimpur'.

FIG. 2 illustrates a close-up view of an individual flower
of 'Balcelimpur'.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible
environmental conditions to date. Accordingly, it is possible
that the phenotype may vary somewhat with variations in the
environment, such as temperature, light intensity, and day
length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2001 edition, except where general color terms of ordinary significance are used. The color values were determined on Jul. 26, 2006 between 9:00 a.m. and 11:00 a.m. under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe plants produced from cuttings taken from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown at West Chicago, Ill. in 4.5 inch pots for 12 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 70° F. to 77° F. (21° C. to 25° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Greenhouse light levels of 2,500 footcandles to 6,000 footcandles were maintained during the day.

Botanical classification: *Impatiens hawkeri* cultivar Balcelimpur.

Parentage:

Female parent.—Harmony Lavender Purple ‘Danharpl’, U.S. Plant Pat. No. 12,094.

Male parent.—Celebration Raspberry Rose ‘Raspberry Rose’, U.S. Plant Pat. No. 9,212.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 7 to 8 days.

Time to produce a rooted cutting.—Approximately 21 to 24 days.

Root description.—Fibrous.

Rooting habit.—Freely branching.

Plant description:

Commercial crop time.—Approximately 7 to 9 weeks from a rooted cutting.

Growth habit and general appearance.—Vigorous, upright-mounded.

Size.—Height from soil level to top of plant plane: Approximately 14.7 cm. Width: Approximately 24.4 cm.

Branching habit.—Freely basal branching. Quantity of main branches per plant: Approximately 6.

Branch.—Strength: Strong. Length: Approximately 9.0 cm. Diameter: Approximately 5.6 mm. Length of central internode: Approximately 1.7 cm. Texture: Glabrous. Color of young stem: 144C. Color of mature stem: 144C with a faint overlay of 185B.

Foliage description:

General description.—Quantity of leaves per main branch: Approximately 7. Fragrance: None. Form: Simple. Arrangement: In whorls of 3 to 5 leaves.

Leaves.—Aspect: Petiole is at an acute angle to stem and leaf blade is perpendicular to stem. Shape: Elliptic. Margin: Serrate, ciliate. Apex: Acuminate. Base: Attenuate. Venation pattern: Pinnate. Length of mature leaf at center of stem: Approximately 9.0 cm. Width of mature leaf at center of stem: Approximately 3.8 cm. Texture of upper and lower surfaces: Glabrous. Color of upper surface of young foliage: 137A with venation of 145C. Color of lower surface of young and mature foliage: Closest to 138B with venation of 145B. Color of upper surface of mature foliage: 137A with venation of 145C and a faint overlay of 185B on lower half of midvein.

Petiole.—Length: Approximately 2.9 cm. Diameter: Approximately 3.7 mm. Texture: Glabrous. Color: Between 145B and 145C.

Flowering description:

Flowering habit.—‘Balcelimpur’ is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

Lastingness of individual flower on the plant.—Approximately 9 to 11 days.

Flower description:

General description.—Type: Single. Quantity per plant: Approximately 10. Fragrance: None.

Bud.—Rate of opening: Generally takes 3 to 4 days for bud to progress from first color to fully open flower.

Quantity showing color per plant: Approximately 2.

Bud just before opening.—Shape: Ovoid. Length: Approximately 2.0 cm. Diameter: Approximately 1.3 cm. Color: Darker than N74A.

Corolla.—Shape: Round, cupped when first open with petals turning downward with age. Width: Approximately 7.2 cm. Depth: Approximately 1.9 cm. Borne: Above and below foliage.

Petals.—Quantity: 5. Shape: Obovate. Aspect: Flat. Appearance: Indescent. Margin: Entire. Apex: Emarginate. Base of upper petal: Truncate. Base of lateral and lower petals: Attenuate. Length of upper petal: Approximately 3.1 cm. Width of upper petal: Approximately 5.1 cm. Length of lateral petals: Approximately 3.4 cm. Width of lateral petals: Approximately 2.9 cm. Length of lower petals: Approximately 3.8 cm. Width of lower petals: Approximately 3.6 cm. Texture of upper and lower surfaces: Glabrous. Color of upper surface when first and fully open: Darker than N74A fading to 75C near base; lateral and lower petals have an eye of 53C. Color of lower surface when first and fully open: N74B.

Spur.—Quantity: 1 per flower. Length: Approximately 5.8 cm. Diameter at tip: Approximately 1.0 mm. Diameter at base: Approximately 3.0 mm. Texture: Glabrous. Color: 187C and tip of 144C.

Peduncle.—Strength: Strong. Aspect: Acute angle to stem. Length: Approximately 4.4 cm. Diameter: Approximately 2.0 mm. Texture: Glabrous. Color: 144C with a faint overlay of 185B.

Sepals.—Quantity per flower: 4, with one fused to base of upper petal. Shape of lateral sepals: Ovate. Shape of lower sepal: Ovate. Apex of all sepals: Acuminate. Length of lateral sepals: Approximately 1.3 cm. Width of lateral sepals: Approximately 6.6 mm. Length of lower sepal: Approximately 1.6 cm. Width of lower sepal: Approximately 1.4 cm. Texture of upper and lower surfaces of all sepals: Glabrous. Color of lateral sepals: 145D with a faint overlay of 185C and tip of 144B. Color of lower sepal: N155B with an overlay of 186C and tip of 144B.

Reproductive organs.—Androecium: Stamen quantity: 5 per flower, fused around pistil at apex. Anther shape: Hooded. Anther length: Approximately 3.0 mm. Anther color: Lighter than 160D at apex transitioning to N66A toward filament. Filament color: N66A. Pollen amount: Abundant. Pollen color: 155D. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 4.0 mm. Stigma shape: 5-pointed star. Stigma color: Colorless, transparent. Style color: 144B. Ovary texture: Glabrous. Ovary color: 144B.

Seed and fruit production: Neither seed nor fruit production has been observed.
Disease and pest resistance: Resistance to pathogens and pests common to New Guinea New Guinea *Impatiens* has not been observed.

What is claimed is:
1. A new and distinct cultivar of New Guinea *Impatiens* plant named ‘Balcelimpur’, substantially as herein shown and described.

* * * * *




FIG. 1



FIG. 2