



US00PP18871P2

(12) **United States Plant Patent**  
**Sanders**

(10) **Patent No.:** **US PP18,871 P2**  
(45) **Date of Patent:** **Jun. 3, 2008**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED  
'INGFILU'**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(50) Latin Name: ***Impatiens hawkeri***  
Varietal Denomination: **Ingfilu**

(52) **U.S. Cl.** ..... **Plt./318**  
(58) **Field of Classification Search** ..... **Plt./318**  
See application file for complete search history.

(75) Inventor: **Monica Maria Adelheid Sanders,**  
Grootebroek (NL)

*Primary Examiner*—Annette H Para  
(74) *Attorney, Agent, or Firm*—Bruce Vrana

(73) Assignee: **Syngenta Seeds B.V.,** Enkhuizen (NL)

(57) **ABSTRACT**

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

A new and distinct cultivar of New Guinea *Impatiens* plant  
named 'Ingfilu' characterized by its large white with pink  
blotch colored flowers with small, dark green foliage, early  
flowering, good heat tolerance, nice rounded shaped growth  
habit with excellent basal branching and good vigor.

(21) Appl. No.: **11/438,650**

(22) Filed: **May 22, 2006**

**1 Drawing Sheet**

**1**

Latin name of the genus and species of the plant claimed:  
*Impatiens hawkeri*.  
Varietal denomination: 'Ingfilu'.

**BACKGROUND OF THE NEW PLANT**

The present invention comprises a new and distinct New  
Guinea *Impatiens* plant botanically known as *Impatiens*  
*hawkeri* and referred to by the cultivar name 'Ingfilu.'

The new cultivar was developed in a controlled breeding  
program conducted by the inventor in Enkhuizen, The  
Netherlands. The new New Guinea *Impatiens* cultivar is  
very floriferous with large flowers, early flowering on a  
strong growing, basal branching plant with good heat tol-  
erance.

The new cultivar is propagated from cuttings resulting  
from the cross in September 2001 of the proprietary New  
Guinea *Impatiens* selection identified as 'H193-2' as female  
parent and the proprietary New Guinea *Impatiens* selection  
identified as 'H197-3' as male parent. 'H193-2' is not  
commercially available and has not been patented. 'H197-3'  
is not commercially available and has not been patented.

As a result of this cross the present cultivar was selected  
in September 2002 in Enkhuizen, Netherlands and has been  
repeatedly asexually reproduced by cuttings in Enkhuizen,  
Netherlands, in Gilroy, Calif., and in Angers, France over a  
period of several years. The distinctive characteristics of this  
new *Impatiens* are stable and reproduced true to type in  
successive generations of asexual reproduction. It takes 9 to  
11 weeks to produce a finished plant, starting from a rooted  
plug and planted in a 12 cm pot, depending on the tempera-  
ture.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of 'Ingfilu.'  
These characteristics in combination distinguish 'Ingfilu' as  
a new and distinct New Guinea *Impatiens* cultivar:

1. Large pinkish white flowers with pink blotch from the  
center
2. Early flowering and floriferous habit
3. Rounded shaped habit

**2**

4. Strong growing and good heat tolerance

Plants of the New Guinea *Impatiens* differ primarily from  
the plants of the female parent selection in the following  
characteristic:

- 5 Plants of the New Guinea *Impatiens* have a pinkish white  
with pink blotch flower color, whereas the plants of the  
female parent selection have a light pink flower color.

Plants of the new New Guinea *Impatiens* differ primarily  
from the plants of the male parent selection in the following  
characteristic:

- 10 Plants of the New Guinea *Impatiens* have a pinkish white  
with pink blotch flower color, whereas the plants of the male  
parent selection have a pink with rose eye flower color.

Plants of the new New Guinea *Impatiens* can be compared  
to plants of 'Fisupnic Mapink,' U.S. Plant Pat. No. 16,222.  
15 In side-by-side comparisons conducted in Enkhuizen,  
Netherlands, plants of the new New Guinea *Impatiens*  
differed from plants of the cultivar 'Fisupnic Mapink' in the  
following characteristics:

- 20 1. Plants of the new New Guinea *Impatiens* have large  
pinkish white flowers with pink blotch from the center,  
whereas plants of 'Fisupnic Mapink' have medium  
large, light pink flowers with white blotch.
- 25 2. Plants of the new New Guinea *Impatiens* have small  
dark green foliage, whereas plants of 'Fisupnic  
Mapink' have large green foliage.

**DESCRIPTION OF THE DRAWING**

30 This new New Guinea *Impatiens* plant is illustrated by the  
accompanying photographic drawing which shows blooms,  
buds and foliage of the plant in full color, the color showing  
being as true as can be reasonably obtained by conventional  
photographic procedures.

**DESCRIPTION OF THE NEW CULTIVAR**

35 The following detailed descriptions set forth the distinc-  
tive characteristics of this new New Guinea *Impatiens*. The  
data which defines these characteristics were collected from  
asexual reproductions carried out in Enkhuizen, Nether-  
lands. The plant history was taken on 30 week old plants,  
40 blossomed under natural light in the field.



Color readings were taken in the laboratory under ambient light. Color references are primarily to The R.H.S. Colour Chart of The Royal Horticultural Society of London.

The plant:

*Classification*.—Botanical: *Impatiens hawkeri*.

Parentage:

*Female parent*.—Proprietary *Impatiens hawkeri* selection identified as code number 'H193-2,' not patented.

*Male parent*.—Proprietary *Impatiens hawkeri* selection identified as code number 'H197-3,' not patented.

Propagation:

*Type cutting*.—Terminal cuttings.

*Time to initiate roots*.—Approximately 7–14 days.

*Time to produce a rooted cutting*.—Approximately 21 days.

*Root description*.—Fine, fibrous, and white in color.

Plant description:

*General appearance*.—Outwardly spreading, low rounded and uniform mounded plant growth.

Habit:

*Growth and branching habit*.—Freely branching habit, freely flowering, moderate vigorous.

Crop time:

*Plant height*.—26–28 cm.

*Plant spread*.—45–50 cm.

Lateral branch description:

*Length*.—17–22 cm.

*Diameter*.—5–7 mm.

*Texture*.—Smooth, glabrous.

*Internode length*.—2–10 cm.

*Color*.—RHS 187A.

Foliage description:

*Arrangement*.—Primarily in whorls.

*Length, mature leaves*.—8.0 cm.

*Width, mature leaves*.—2.9 cm.

*Shape*.—Lanceolate.

*Apex*.—Acuminate.

*Base*.—Acute.

*Margin*.—Serrulate with fine ciliation.

*Texture*.—Smooth, glabrous.

*Venation pattern*.—Pinnate.

*Color young foliage, upper surface*.—RHS 147A.

*Color young foliage, lower surface*.—RHS 187A.

*Fully expanded foliage, upper surface*.—RHS 147A.

*Fully expanded foliage, lower surface*.—RHS 187A.

*Venation, upper surface*.—RHS 187A+146A.

*Venation, lower surface*.—RHS 187A.

*Anthocyanin may be present*.—RHS 187A.

*Petiole length*.—About 0.4 cm.

*Petiole diameter*.—About 2.0 mm.

*Texture*.—Smooth, glabrous.

*Color*.—RHS 187A.

Flower description:

*Flower type and flowering habit*.—Single.

*Number of flowers per leaf axil*.—1 or 2.

*Number of flowers and flower buds per lateral branch*.—6.

*Flower position*.—Above and beyond the foliage and typically facing upwards and outward.

*Flower shape*.—Rounded, mostly flat.

*Flowers lasting*.—About 10 days on the plant depending on environmental conditions.

*Petals*.—Self cleaning.

*Gynoecium*.—Persistent.

*Flowers*.—Not fragrant.

*Flowering*.—Indeterminate and continuous.

*Flowering season*.—Year round under greenhouse conditions. In the garden, flowering from spring until fall.

*Flower length*.—About 7.2 cm.

*Flower width*.—About 6.8 cm.

*Flower depth*.—About 0.6 cm.

Flower buds:

*Bud length*.—About 2.5 cm.

*Bud diameter*.—About 1.7 cm.

*Bud shape*.—Ellipsoidal.

*Texture*.—Smooth, glabrous.

*Color, just before opening*.—RHS 53C to 55B to 69D, base to apex.

Petals:

*Quantity*.—Single, five per flower, imbricate.

*Length of banner petal*.—About 3.1 cm.

*Width of banner petal*.—About 4.7 cm.

*Length lateral and basal petals*.—About 3.9 cm.

*Width lateral and basal petals*.—About 3.1 cm.

*Lateral and basal petal shape*.—Cordate.

*Banner petal shape*.—Roughly cordate.

*Petal apex*.—Retuse to emarginate.

*Petal base*.—Attenuate.

*Petal margin*.—Entire.

*Petal texture*.—Smooth, satiny.

*Petal color*.—Upper surface: When opening and fully opened: RHS 69D. Middle eye N57A. Lower surface: When opening and fully opened: RHS 69D. Middle eye 58A.

Spur:

*Length*.—About 5.5 cm.

*Texture*.—Smooth, glabrous.

*Aspect*.—Curved.

*Color*.—RHS 182C to 152D.

Peduncles:

*Length*.—About 5.1 cm.

*Strength*.—Strong, flexible.

*Diameter*.—0.15 cm.

*Aspect*.—About 45° from the stem.

*Color*.—RHS 183B.

Reproductive organs:

*Androecium*.—Stamen number: Five fused at anthers, filaments free, hooded.

*Anther shape*.—Obovate.

*Anther length*.—About 7 mm.

*Anther color*.—RHS 155C.

*Amount of pollen*.—Abundant.

*Pollen color*.—RHS 4D.

Gynoecium:

*Pistil number*.—One.

*Pistil length*.—About 5 mm.

*Stigma shape*.—Columnar, five segmented.

*Stigma color*.—Colorless.

*Style color*.—RHS 187A.

*Ovary arrangement*.—Five celled.

*Ovary color*.—RHS 144A.

Seed development: Seed development has not been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of New Guinea *Impatiens* plant named 'Ingfilu', as substantially illustrated and described herein.

\* \* \* \* \*



