



US00PP18943P2

(12) **United States Plant Patent**
Smith

(10) **Patent No.:** **US PP18,943 P2**
(45) **Date of Patent:** **Jun. 17, 2008**

(54) **CHRYSANTHEMUM PLANT NAMED ‘COOL YOCHERYL’**

(50) Latin Name: *Chrysanthemum*×*morifolium*
Varietal Denomination: **Cool Yocheryl**

(75) Inventor: **Mark A. Smith**, Fort Myers, FL (US)

(73) Assignee: **Yoder Brothers Inc.**, Barberton, OH (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/641,534**

(22) Filed: **Dec. 18, 2006**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./287**

(58) **Field of Classification Search** Plt./287,
Plt./294
See application file for complete search history.

(56) **References Cited**
U.S. PATENT DOCUMENTS

PP10,218 P * 1/1998 Pieters Plt./294
PP11,982 P2 * 7/2001 Glicenstein Plt./287
PP16,296 P2 * 2/2006 Smith Plt./291

* cited by examiner

Primary Examiner—Wendy C. Haas
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named ‘Cool Yocheryl’, characterized by its compact, upright and outwardly spreading plant habit; freely branching habit; dense and full plant habit; uniform and freely flowering habit; decorative-type inflorescences with elongated oblong to ligulate-shaped ray florets; white-colored ray florets with purple-colored apices; and natural season flowering about October 6th in the Northern Hemisphere.

1 Drawing Sheet

1

Botanical designation: *Chrysanthemum*×*morifolium*.
Cultivar denomination: ‘Cool Yocheryl’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Chrysanthemum* plant, botanically known as *Chrysanthemum*×*morifolium*, commercially grown as a garden *Chrysanthemum* and hereinafter referred to by the name ‘Cool Yocheryl’.

The objective of the breeding program is to create new garden-type *Chrysanthemum* cultivars having inflorescences with desirable inflorescence forms, attractive floret colors and good garden performance.

The new *Chrysanthemum* is a naturally-occurring whole plant mutation of the *Chrysanthemum*×*morifolium* cultivar Soft Yocheryl, disclosed in U.S. Plant Pat. No. 16,296. The new *Chrysanthemum* was discovered and selected by the Inventor as a single flowering plant within a population of plants of the parent cultivar in December, 2003, in Alva, Fla. The selection of this plant was based on its desirable inflorescence color and good form and substance.

Asexual reproduction of the new *Chrysanthemum* by vegetative cuttings was first conducted in Alva, Fla. in February, 2004. Asexual reproduction by cuttings has shown that the unique features of this new *Chrysanthemum* are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the cultivar Cool Yocheryl have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature, daylength and light intensity, without, however, any variance in genotype.

2

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Cool Yocheryl’. These characteristics in combination distinguish ‘Cool Yocheryl’ as a new and distinct garden *Chrysanthemum* cultivar:

1. Compact, upright and outwardly spreading plant habit.
2. Freely branching habit; dense and full plant habit.
3. Uniform and freely flowering habit.
4. Decorative-type inflorescences with elongated oblong to ligulate-shaped ray florets.
5. White-colored ray florets with purple-colored apices.
6. Natural season flowering about October 6th in the Northern Hemisphere.

In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the parent, the cultivar Soft Yocheryl, in the following characteristics:

1. Plants of the new *Chrysanthemum* were larger than plants of the cultivar Soft Yocheryl.
2. Plants of the new *Chrysanthemum* flowered later than plants of the cultivar Soft Yocheryl when grown under natural season conditions.
3. Ray florets of plants of the new *Chrysanthemum* were white and purple in color whereas ray florets of plants of the cultivar Soft Yocheryl were light lavender in color.

Plants of the new *Chrysanthemum* can be compared to plants of the *Chrysanthemum* cultivar Yoroxanne, disclosed in U.S. Plant Pat. No. 11,982. In side-by-side comparisons conducted in Alva, Fla., plants of the new *Chrysanthemum* differed from plants of the cultivar Yoroxanne in the following characteristics:

1. Plants of the new *Chrysanthemum* were larger and more mounding than plants of the cultivar Yoroxanne.

2. Plants of the new *Chrysanthemum* flowered more uniformly than plants of the cultivar Yoroxanne.
3. Plants of the new *Chrysanthemum* had slightly smaller inflorescences than plants of the cultivar Yoroxanne.
4. Plants of the new *Chrysanthemum* and the cultivar Yoroxanne differed slightly in ray floret coloration.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Chrysanthemum*. These photographs show the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Chrysanthemum*.

The photograph at the bottom of the sheet comprises a side perspective view of a typical flowering plant of 'Cool Yocheryl'.

The photograph at the top of the sheet is a close-up view of typical inflorescences of 'Cool Yocheryl'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations and measurements describe plants grown in Fletcher, N.C. during the summer in an outdoor nursery and under conditions and practices which approximate those generally used in commercial garden *Chrysanthemum* production. During the production of the plants, day temperatures averaged 29° C. and night temperatures averaged 16° C. Plants were grown in 15-containers, exposed to long day/short night conditions and pinched about two weeks later. About two weeks after the pinch, the photoinductive short day/long night treatments were started. Plants used in the photographs and for the description were about three months old. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Chrysanthemum* × *morifolium* cultivar Cool Yocheryl.

Parentage: Naturally-occurring whole plant mutation of the *Chrysanthemum* × *morifolium* cultivar Soft Yocheryl, disclosed in U.S. Plant Pat. No. 16,296.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate roots.—About four days at temperatures of about 21° C.

Time to produce a rooted young plant.—About ten to twelve days at temperatures of about 21° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching.

Plant description:

Appearance.—Herbaceous decorative-type garden *Chrysanthemum*. Stems upright and outwardly spreading giving a uniformly mounded appearance to the plant. Freely branching habit, about six lateral branches develop after removal of terminal apex (pinching) each with numerous secondary laterals; dense and full plant habit. Strong and vigorous growth habit.

Plant height.—About 18 cm.

Plant width.—About 24 cm.

Lateral branches.—Length: About 17 cm. Diameter: About 7 mm. Internode length: About 1 cm. Strength: Strong. Texture: Pubescent. Color: 146C.

Leaves.—Arrangement: Alternate, simple. Length: About 5.6 cm. Width: About 4.4 cm. Apex: Broadly acute. Base: Truncate to attenuate. Margin: Palmately lobed, sinuses between lateral lobes parallel to divergent. Texture, upper and lower surfaces: Fine pubescence; veins prominent on lower surface. Color: Developing foliage, upper and lower surfaces: 147A. Fully expanded foliage, upper surface: 147A; venation, 147C. Fully expanded foliage, lower surface: 147B; venation, 147C. Petiole: Length: About 1.3 cm. Diameter: About 2 mm. Texture, upper and lower surfaces: Pubescent. Color, upper and lower surfaces: 147B.

Inflorescence description:

Appearance.—Decorative-type inflorescence form with elongated oblong to ligulate-shaped ray florets. Inflorescences borne on terminals above foliage. Disk and ray florets arranged acropetally on a capitulum. Inflorescences not fragrant.

Flowering response.—Under natural season conditions, plants flower about October 6th in the Northern Hemisphere. At other times of the year, inflorescence initiation and development can be induced under short day/long night conditions (at least 13.5 hours of darkness). Early flowering habit; plants exposed to photoinductive short day/long night conditions flower about 53 days later.

Postproduction longevity.—Inflorescences maintain good color and substance for about four weeks in an outdoor nursery.

Quantity of inflorescences.—About 27 to 28 inflorescences develop per lateral branch.

Inflorescence bud.—Height: About 1.5 cm. Diameter: About 1.2 cm. Shape: Ovoid. Color: Towards the apex, 186C to 186D; towards the base, 155D.

Inflorescence size.—Diameter: About 4.7 cm. Depth (height): About 2 cm. Disc diameter: About 5 mm; inconspicuous. Receptacle diameter: About 1.6 cm. Receptacle height: About 7 mm.

Ray florets.—Shape: Elongated-oblong to ligulate. Orientation: Initially upright, then about 60° from vertical. Aspect: Initially incurved, then mostly concave. Length: About 2.2 cm. Width: About 8 mm. Apex: Emarginate. Base: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous, satiny. Number of ray florets per inflorescence: About 170 arranged in about 16 to 17 whorls. Color: When opening, upper surface: 155D. When opening, lower surface: 155A; towards the apex, 186C to 186D. Fully opened, upper and lower surfaces: 155D.

Disc florets.—Shape: Tubular, elongated. Length: About 3 mm. Diameter: Less than 1 mm. Number of disc florets per inflorescence: About 60. Color, immature and mature: Apex: Close to 183A. Mid-section: Close to 182D. Base: Close to 157A.

Phyllaries.—Number of phyllaries per inflorescence: About 30 arranged in about three whorls. Length: About 5 mm. Width: About 2 mm. Shape: Elliptical. Apex: Acute. Base: Truncate. Texture, upper surface: Smooth, waxy. Texture, lower surface: Pubescent. Color, upper surface: Close to 146A. Color, lower surface: Close to 147B.

5

Peduncles.—Length: About 4 cm. Diameter: About 2 mm. Angle: About 45° from vertical. Strength: Strong. Texture: Pubescent; longitudinally ridged. Color: Close to 148A.

Reproductive organs.—Androecium: None observed. Gynoecium: Pistil length: About 4 mm. Stigma shape: Bi-parted. Stigma color: Close to 2A. Style length: About 2 mm. Style color: Close to 2D. Ovary color: Close to 157A.

Seed/fruit.—Seed and fruit production has not been observed.

6

Disease/pest resistance: Resistance to pathogens and pests common to *Chrysanthemums* has not been observed on plants grown under commercial conditions.

Garden performance: Plants of the new *Chrysanthemum* have demonstrated excellent garden performance and to tolerate temperatures from about 0° C. to about 38° C.

It is claimed:

1. A new and distinct *Chrysanthemum* plant named 'Cool Yocheryl' as illustrated and described.

* * * * *

