



US00PP18958P2

(12) **United States Plant Patent**
Kristensen

(10) **Patent No.:** **US PP18,958 P2**
(45) **Date of Patent:** **Jun. 17, 2008**

(54) **OSTEOSPERMUM PLANT NAMED**
'SAKCADNUCOP'

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(50) Latin Name: *Osteospermum hybrida*
Varietal Denomination: **Sakcadnucop**

(52) **U.S. Cl.** **Plt./360**

(58) **Field of Classification Search** **Plt./360**
See application file for complete search history.

(75) Inventor: **Niels G. Kristensen**, Marslev (DK)

Primary Examiner—Kent Bell

Assistant Examiner—S. B. McCormick-Ewoldt

(73) Assignee: **Sakata Ornamentals Europe A/S**,
Marslev (DK)

(74) *Attorney, Agent, or Firm*—Jondle & Associates, P.C.

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(57) **ABSTRACT**

An *Osteospermum* cultivar particularly distinguished by its
purple-terracotta-copper flower color, large flower size with
flowers that remain open under low light conditions and its
compact plant growth habit.

(21) Appl. No.: **11/717,339**

1 Drawing Sheet

(22) Filed: **Mar. 13, 2007**

1

2

Genus and species: *Osteospermum hybrida*.
Variety denomination: 'Sakcadnucop'.

DESCRIPTION OF PHOTOGRAPHS

BACKGROUND OF THE NEW PLANT

This new *Osteospermum* plant is illustrated by the accom-
panying photographs which show the plant's form, foliage
and flowers. The colors shown are as true as can be reason-
ably obtained by conventional photographic procedures.

'Sakcadnucop' originated from a hybridization of propri-
etary hybrid *Osteospermum* breeding line '203005'
(unpatented) and commercial hybrid *Osteospermum* line
'Sunny Amanda' (U.S. Plant Pat. No. 16,522) in Aabyhoej,
Denmark. The male parent, 'Sunny Amanda', has a pale
yellow flower color with terracotta-brown at the flower petal
apices, medium flower size and brown disc florets. The
female parent, '203005' has a bright yellow flower color,
medium flower size and a compact and less branching plant
growth habit.

FIG. 1 shows overall plant habit.

FIG. 2 shows the mature inflorescence.

In spring 2003, the two *Osteospermum* lines were crossed
and 552 seeds were obtained. The seeds were sown and 442
plants were grown in pots for evaluation. Out of 42 F₁ lines,
plant number 196 was selected for its unique copper-purple
flower color, large flowers which remain open even under
low light conditions and dark green plant foliage.

DESCRIPTION OF THE NEW CULTIVAR

In February 2004, plant number 196 was vegetatively
propagated with cuttings and re-evaluated in an open field
and a greenhouse. Plant number 196 was given the code
number '204056'.

The following detailed descriptions set forth the distinc-
tive characteristics of 'Sakcadnucop'. The data which define
these characteristics were collected from asexual reproduc-
tions carried out in Salinas, Calif. Data was collected on
plants grown approximately five months from transplanting
rooted cuttings into 6-inch pots under greenhouse conditions
in Salinas, Calif. Color references are to the R.H.S. Colour
Chart of The Royal Horticultural Society of London
(R.H.S.).

DESCRIPTION OF THE NEW PLANT

In spring 2004, plants were evaluated again in pots and in
an open field. The selection was named 'Sakcadnucop' and
found to retain its distinctive characteristics through succes-
sive asexual propagations.

Classification:

Family.—Compositae.

Botanical.—*Osteospermum hybrida*.

Parentage:

Female parent.—Hybrid proprietary *Osteospermum*
line '203005' (unpatented).

Male parent.—*Osteospermum* line 'Sunny Amanda'
(U.S. Plant Pat. No. 16,522).

Growth:

Time to produce a rooted cutting.—Cuttings will colo-
nize a 2.5 cm diameter by 2.5 cm tall greenhouse tray
cell with peat-based plant media in approximately
four weeks. Cuttings are dipped in a normal dilution
(1:9) of Dip 'N Grow root-inducing solution in
water. The trays are misted hourly during rooting.

SUMMARY OF THE INVENTION

The following are the most outstanding and distinguish-
ing characteristics of this new cultivar when grown under
normal horticultural practices in Kakagawa, Japan.

Environmental conditions for plant growth.—Rooted
cuttings are transplanted to pots with a 16 cm
diameter, one plant per pot. Peat-based growing
media is used. The pots are watered using a 150-200
ppm fertilizer solution containing 18% nitrogen, 8%
phosphorus and 18% potassium. The soil is allowed
to dry between waterings. During the first few weeks
after transplanting, the plants should have evening
temperatures around 15-18° C. for good root growth.

1. Purple-terracotta-copper flower color
2. Large flower size;
3. Flowers that remain open under low light conditions; and
4. Compact plant growth habit.

When plants reach 7.5-10 cm in height they are pinched back to 5-6 leaves to promote branching. Spring and summer daytime high temperatures in Salinas, Calif., where the data was collected, range from 16-25° C.

Time to bloom from propagation.—Approximately four weeks when rooted vegetative cuttings are transferred to a 16 cm. diameter pot. Flowering season is all year in the United States. Vernalization is not required to induce flowering.

Plant description:

Habit.—Compact, mounded and branching.

Life cycle.—Perennial.

Form.—Branching, dense.

Height.—43 cm to 48 cm.

Width.—40 cm to 50 cm.

Stems:

Internode length.—1.5 cm to 3.0 cm.

Stem description.—Strong, erect.

Stem diameter.—0.4 cm to 0.5 cm.

Stem color.—RHS 145A (Yellow-green).

Pubescence.—Sparse.

Pubescence shape.—Linear.

Pubescence color.—RHS N155A (White).

Anthocyanin color.—RHS 64A (Red-purple).

Leaves:

Arrangement.—Alternate.

Shape.—Oblanceolate.

Apex.—Acute.

Base.—Decurrent.

Margin.—Serrate.

Venation.—Pinnate.

Surface.—Dull, smooth.

Surface pubescence.—Present.

Length.—7.5 cm to 8.5 cm.

Width.—2.5 cm to 3.0 cm.

Thickness.—0.1 cm.

Color.—Upper surface: RHS 137B (Green). Lower surface: RHS 138B (Green).

Venation color.—Upper surface: RHS 144B (Yellow-green). Lower surface: RHS 139C (Green).

Inflorescence:

Type.—Capitulum (head); Disc florets are perfect and ray florets are carpellate.

Flowering habit.—Determinate.

Lastingness of flowers.—3 to 4 days.

Fragrance.—None.

Seed production.—None observed.

Diameter.—7.0 cm to 8.0 cm.

Depth.—1.2 cm to 1.5 cm.

Disc diameter.—1.5 cm.

Disc floret number.—80 to 100 per inflorescence.

Peduncle length.—7.5 cm to 11 cm.

Peduncle diameter.—0.2 cm to 0.25 cm.

Peduncle color.—RHS 144A (Yellow-green).

Peduncle texture.—Dull, pubescent.

Phyllaries.—Arrangement: 20 to 24 per inflorescence, free, arranged symmetrically. Length: 1.0 cm to 1.2 cm. Width: 0.15 cm to 0.20 cm. Apex: Acute. Margin: Entire. Shape: Linear, acute. Color: Upper surface: RHS 144A (Yellow-green). Lower surface: RHS 145A (Yellow-green).

Ray florets (ligules):

Corolla.—One petal per ray floret; only the outer row of florets are the ray florets.

Number of ray florets per inflorescence.—21 to 30.

Petal length.—3.4 cm to 3.8 cm.

Petal width.—0.7 cm to 0.8 cm.

Petal shape.—Spatulate.

Petal apex.—Obtuse.

Petal margin.—Entire.

Petal color.—Upper surface: RHS N77B (Purple).

Lower surface: RHS 167C (Gray-orange) with streaks of RHS 183A (Gray-purple).

Petal pubescence.—Glabrous.

Ovary.—Superior.

Pistil form.—One style with two stigma branches.

Pistil length.—0.3 cm.

Stigma color.—RHS 83A (Purple).

Style color.—RHS 83A (Purple).

Disc florets:

Color.—RHS N89A (Violet-blue).

Shape.—Tubular.

Apex.—Rounded.

Surface.—Smooth, shiny.

Size.—Length: 0.5 cm. Width: 0.1 cm.

Pistil form.—One style with two stigma branches.

Ovary.—Superior.

Style color.—RHS 83A (Purple).

Stigma color.—RHS 83A (Purple).

Stamens.—5, fused into a single tube.

Anther color.—RHS 83A (Purple).

Pollen color.—RHS 23A (Yellow-orange).

Disease and insect resistance: Very disease resistant, however, plants are susceptible to aphids, thrips, whiteflies and worms.

COMPARISON WITH PARENTAL LINES AND KNOWN CULTIVARS

‘Sakcadnucoop’ is a distinct variety of *Osteospermum* owing to its purple-terracotta-copper flower color, large flower size with flowers that remain open under low light conditions and its compact plant growth habit. ‘Sakcadnucoop’ is distinguished from its parents primarily by ray floret color, flower size and plant growth habit as described in Table 1 below (color references are to The Royal Horticultural Society Colour Chart, 4th edition):

TABLE 1

Characteristic	‘Sakcadnucoop’	Male Parent ‘Sunny Amanda’	Female Parent ‘203005’
Ray floret (ligule) color, upper surface:	RHS N77B (Purple)	Pale yellow terracotta-brown tips	Bright yellow
Flower size	Large	Medium	Medium
Disc florets color	Brown	Blue	

‘Sakcadnucoop’ is most similar to the *Osteospermum* plants named ‘Iringa’ (U.S. Plant Pat. No. 15,632) and ‘Wildside’ (U.S. Plant Pat. No. 10,603); however, there are differences as described in Table 2 below (color references are to The Royal Horticultural Society Colour Chart, 4th edition):

TABLE 2

Characteristic	‘Sakcadnucoop’	‘Iringa’	‘Wildside’
Petal color:	RHS N77B (Purple)	RHS 71A (Red-purple)	RHS 71A (Red Purple)
Inflorescence diameter	7.0 cm to 8.0 cm	6.5 cm to 7.0 cm	5.0 cm to 5.5 cm
Habit	Mounded and compact	Mounded, upright and branching	Erect and less compact

I claim:

1. A new and distinct cultivar of *Osteospermum* plant as shown and described herein.

* * * * *



Fig. 1



Fig. 2