



US00PP19138P2

(12) **United States Plant Patent**  
**Ren**

(10) **Patent No.:** **US PP19,138 P2**  
(45) **Date of Patent:** **Aug. 26, 2008**

(54) **CALIBRACHOA PLANT NAMED**  
**'BALCABLAV'**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(50) Latin Name: *Calibrachoa*×*hybrida*  
Varietal Denomination: **Balcablav**

(52) **U.S. Cl.** ..... **Plt./263**  
(58) **Field of Classification Search** ..... **Plt./263**  
See application file for complete search history.

(75) Inventor: **Jianping Ren**, Geneva, IL (US)

*Primary Examiner*—Kent Bell

(73) Assignee: **Ball Horticultural Company**, West  
Chicago, IL (US)

*Assistant Examiner*—June Hwu

(74) *Attorney, Agent, or Firm*—Audrey Charles

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 104 days.

(57) **ABSTRACT**

A new and distinct cultivar of *Calibrachoa* plant named  
'Balcablav', characterized by its light lavender-colored  
flowers, medium green-colored foliage, and moderately  
vigorous, mounded-trailing growth habit.

(21) Appl. No.: **11/602,624**

**1 Drawing Sheet**

(22) Filed: **Nov. 20, 2006**

**1**

**2**

Latin name of genus and species of plant claimed: *Cali-  
brachoa*×*hybrida*.  
Variety denomination: 'Balcablav'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of *Calibrachoa* plant botanically known as *Calibrachoa*×  
*hybrida* and hereinafter referred to by the cultivar name  
'Balcablav'.

The new cultivar originated in a controlled breeding pro-  
gram in Elburn, Ill. during December 2003. The objective of  
the breeding program was the development of *Calibrachoa*  
cultivars with attractive flower coloration, medium green-  
colored foliage, and a moderately vigorous, mounded-  
trailing growth habit.

The new *Calibrachoa* cultivar is the result of cross-  
pollination. The female (seed) parent of the new cultivar is  
the proprietary *Calibrachoa*×*hybrida* breeding selection  
designated 2085-1A, not patented, characterized by its  
medium lavender-colored flowers, medium green-colored  
foliage, and mounded, trailing growth habit. The male  
(pollen) parent of the new cultivar is a *Calibrachoa*×*hybrida*  
mixed pollen source from plants having various colored  
flowers, medium to dark green-colored foliage, and prostrate  
to mounded growth habits. The new cultivar was discovered  
and selected as a single flowering plant within the progeny  
of the above stated cross-pollination during August 2004 in a  
controlled environment at Elburn, Ill.

Asexual reproduction of the new cultivar by terminal stem  
cuttings since August 2004 at Arroyo Grande, Calif. and  
West Chicago, Ill. has demonstrated that the new cultivar  
reproduces true to type with all of the characteristics, as  
herein described, firmly fixed and retained through succes-  
sive generations of such asexual propagation.

**SUMMARY OF THE INVENTION**

The following characteristics of the new cultivar have  
been repeatedly observed and can be used to distinguish  
'Balcablav' as a new and distinct cultivar of *Calibrachoa*  
plant:

1. Light lavender-colored flowers;
  2. Medium green-colored foliage; and
  3. Moderately vigorous, mounded-trailing growth habit.
- Plants of the new cultivar differ from plants of the female  
parent primarily in flower color.

Of the many commercially available *Calibrachoa* culti-  
vars known to the inventor, the most similar in comparison to  
the new cultivar is Callie Purple 'Cal Pur', U.S. Plant Pat.  
No. 15,426. However, in side by side comparisons, plants of  
the new cultivar differ from plants of 'Cal Pur' in the follow-  
ing characteristics:

1. Plants of the new cultivar are narrower than plants of  
'Cal Pur';
2. Plants of the new cultivar have larger flowers than  
plants of 'Cal Pur'; and
3. Plants of the new cultivar have a flower color different  
from plants of 'Cal Pur'.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographs show, as nearly true as it  
is reasonably possible to make the same in color illustrations  
of this type, typical flower and foliage characteristics of the  
new cultivar. Colors in the photographs differ slightly from  
the color values cited in the detailed description, which  
accurately describes the colors of 'Balcablav'. The plants  
were grown in 4.5 inch pots for 16 weeks in a greenhouse at  
West Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and  
flowering habit of 'Balcablav'.

FIG. 2 illustrates a close-up view of an individual flower  
of 'Balcablav'.

**DETAILED BOTANICAL DESCRIPTION**

The new cultivar has not been observed under all possible  
environmental conditions to date. Accordingly, it is possible  
that the phenotype may vary somewhat with variations in the  
environment, such as temperature, light intensity, and day  
length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2001 edition, except where general color terms of ordinary significance are used. The color values were determined on Mar. 24, 2006 between 9:00 a.m. and 11:00 a.m. under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe plants produced from cuttings taken from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown at West Chicago, Ill. in 4.5 inch pots for 16 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 70° F. to 77° F. (21° C. to 25° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Greenhouse light levels of 2,500 footcandles to 6,000 footcandles were maintained during the day.

Botanical classification: *Calibrachoa x hybrida* cultivar Balcablav.

Parentage:

*Female parent.*—Proprietary *Calibrachoa x hybrida* breeding selection designated 2085-1A.

*Male parent.*—*Calibrachoa x hybrida* mixed pollen source.

Propagation:

*Type cutting.*—Terminal stem.

*Time to initiate roots.*—Approximately 6 to 8 days.

*Time to produce a rooted cutting.*—Approximately 24 to 28 days.

*Root description.*—Fibrous.

*Rooting habit.*—Freely branching.

Plant description:

*Commercial crop time.*—Approximately 6 to 7 weeks from a rooted cutting.

*Growth habit and general appearance.*—Moderately vigorous and trailing, becoming mounded with age.

*Size.*—Height from soil level to top of plant plane: Approximately 11.3 cm. Width: Approximately 50.9 cm.

*Branching habit.*—Freely branching. Quantity of main branches per plant: Approximately 5.

*Branch.*—Strength: Strong. Length: Approximately 32.4 cm. Diameter: Approximately 2.0 mm. Length of central internode: Approximately 1.8 cm. Texture: Densely glandular pubescent. Color of young stem: 144B. Color of mature stem: 144B with overlay of 187B on upward facing surface.

Foliage Description:

*General description.*—Quantity of leaves per main branch: Approximately 42. Fragrance: Slight. Form: Simple. Arrangement on flowering stem: Alternate.

*Leaves.*—Aspect: At acute angle to stem with tip turning downward. Shape: Elliptic. Margin: Entire. Apex: Obtuse. Base: Attenuate. Venation pattern: Pinnate. Length of mature leaf: Approximately 3.1 cm. Width of mature leaf: Approximately 1.3 cm. Texture of upper and lower surfaces: Glandular pubescent. Color of upper surface of young foliage: 137C with venation of 144D. Color of lower surface of young foliage: 138C with venation of 144D. Color of upper surface of mature foliage: 137A with venation of 144D. Color of lower surface of mature foliage: 138B with venation of 144D.

*Petiole.*—Length: Approximately 5.7 mm. Diameter: Approximately 1.0 mm. Texture: Glandular pubescent. Color: 144D.

Flowering Description:

*Flowering habit.*—‘Balcablav’ is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

*Lastingness of individual flower on the plant.*—Approximately 9 to 11 days.

Flower Description:

*General description.*—Type: Single, salverform. Quantity per plant: Approximately 41. Fragrance: None.

*Bud.*—Rate of opening: Generally takes 3 to 4 days for bud to progress from first color to fully open flower. Quantity per plant: Approximately 24.

*Bud just before opening.*—Shape: Oblong. Length: Approximately 1.9 cm. Diameter: Approximately 5.2 mm. Texture: Densely glandular pubescent. Color: Petals of N155A and corolla tube of 145C, both with venation of N144D having an overlay of N186C.

*Corolla.*—Diameter: Approximately 3.9 cm.

*Petals.*—Quantity: 5, fused to form a tube. Shape: Obovate. Margin: Entire. Apex: Obtuse. Length from tube: Approximately 2.1 cm. Length of free portion: Approximately 6.7 mm. Width: Approximately 2.0 cm. Texture of upper surface: Glabrous. Texture of lower surface: Glandular pubescent, dense along the veins. Color of upper surface when first open: Darker than N87A with midveins of N186B. Color of lower surface when first open: N87C with midveins of 144D having a faint overlay of N186C. Color of upper surface when fully open: N87A fading to N87C with age and midveins of N186B. Color of lower surface when fully open: N87D with midveins of 144D having a faint overlay of N186C.

*Corolla tube.*—Length: Approximately 1.6 cm. Diameter at distal end: Approximately 8.8 mm. Diameter at proximal end: Approximately 2.0 mm. Texture of inner surface: Glabrous. Texture of outer surface: Densely glandular pubescent. Color of inner surface: 12B transitioning to 155D at distal end and venation of N186B. Color of outer surface: 145C transitioning to 155D at distal end with venation of 144C having a faint overlay of N186C.

*Peduncle.*—Strength: Strong. Aspect: Acute angle to stem. Length: Approximately 1.2 cm. Diameter: Approximately 1.0 mm. Texture: Densely glandular pubescent. Color: 144B.

*Sepals.*—Quantity per flower: 5, fused along lower half. Shape: Lanceolate. Apex: Acute. Length: Approximately 1.5 cm. Width: Approximately 3.0 mm. Texture of upper and lower surfaces: Densely glandular pubescent. Color of upper surface: 137B. Color of lower surface: 137B with 144B at base.

*Reproductive organs.*—Androecium: Stamen quantity: 5 per flower, fused to inside of corolla tube. Stamen length: Approximately 8.8 mm. Filament length of free portion: Approximately 4.8 mm. Filament color: 145D with a faint overlay of N87B. Anther shape: 2 parted, oval. Anther length: Approximately 1.0 mm. Anther color: 8A. Pollen amount: Abundant. Pollen color: 8C. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 9.0 mm. Stigma shape:

US PP19,138 P2

**5**

Funnel. Stigma length: Approximately 1.0 mm. Stigma color: 143A. Style length: Approximately 6.1 mm. Style color: 144D. Ovary diameter: Approximately 1.9 mm. Ovary color: 144A with spots of 187A at the distal end.

Seed and fruit production: Neither seed nor fruit production has been observed.

**6**

Disease and pest resistance: Resistance to pathogens and pests common to *Calibrachoa* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Calibrachoa* plant named 'Balcablav', substantially as herein shown and described.

\* \* \* \* \*



FIG. 1




FIG. 2