



US00PP19163P2

(12) **United States Plant Patent**  
**Cook et al.**

(10) **Patent No.:** **US PP19,163 P2**  
(45) **Date of Patent:** **Sep. 2, 2008**

(54) **OZOTHAMNUS PLANT NAMED '13-15'**

(50) Latin Name: *Ozothamnus diosmifolius*  
Varietal Denomination: **13-15**

(75) Inventors: **Edward Graham Cook**, Helidon (AU);  
**Esther Ruth Cook**, Helidon (AU)

(73) Assignee: **Amerinova Properties LLC**, Bonsall,  
CA (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/800,771**

(22) Filed: **May 7, 2007**

(51) **Int. Cl.**  
*A01H 5/00* (2006.01)

(52) **U.S. Cl.** ..... **Plt./263.1**

(58) **Field of Classification Search** ..... **Plt./263.1**  
See application file for complete search history.

*Primary Examiner*—Annette H Para

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Ozothamnus* plant named  
'13-15', characterized by its upright plant habit; vigorous  
growth habit; short internodes; dense and bushy habit; freely  
flowering habit; and inflorescences arranged in corymbs  
with white-colored involucre bracts.

**1 Drawing Sheet**

**1**

Botanical designation: *Ozothamnus diosmifolius*.  
Cultivar denomination: '13-15'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of *Ozothamnus* plant, botanically known as *Ozothamnus*  
*diosmifolius*, and hereinafter referred to by the name  
'13-15'.

The new *Ozothamnus* is a product of a planned breeding  
program conducted by the Inventors in Helidon, Queensland,  
Australia. The objective of the breeding pro-  
gram is to create vigorous and upright *Ozothamnus* cultivars  
with numerous and attractive flowers.

The new *Ozothamnus* originated from a self-pollination  
made by the Inventors in Helidon, Queensland, Australia in  
1993, of the *Ozothamnus diosmifolius* cultivar Pom Pom, not  
patented. The new *Ozothamnus* was discovered and selected  
by the Inventors as a single flowering plant from within the  
progeny of the stated self-pollination grown in a controlled  
environment in Helidon, Queensland, Australia in March,  
1993.

Asexual reproduction of the new *Ozothamnus* by vegeta-  
tive cuttings in a controlled environment in Helidon,  
Queensland, Australia since 1994, has shown that the unique  
features of this new *Ozothamnus* are stable and reproduced  
true to type in successive generations.

**SUMMARY OF THE INVENTION**

The cultivar 13-15 has not been observed under all pos-  
sible environmental conditions. The phenotype may vary  
somewhat with variations in environment such as tempera-  
ture and light intensity, without, however, any variance in  
genotype.

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of '13-15'.  
These characteristics in combination distinguish '13-15' as  
a new and distinct cultivar of *Ozothamnus*:

1. Upright plant habit.
2. Vigorous growth habit.

**2**

3. Short internodes; dense and bushy habit.
4. Freely flowering habit.
5. Inflorescences arranged in corymbs with white-colored  
involucre bracts.

Plants of the new *Ozothamnus* differ from plants of the  
parent, the cultivar Pom Pom, in the following characteris-  
tics:

1. Plants of the new *Ozothamnus* are more upright than  
plants of the cultivar Pom Pom.
2. Plants of the new *Ozothamnus* and the cultivar Pom  
Pom differ in involucre bract color as plants of the  
cultivar Pom Pom have pink-colored involucre bracts.

Plants of the new *Ozothamnus* can be compared to plants  
of the *Ozothamnus* cultivar Just Blush, not patented. In side-  
by-side comparisons conducted in Helidon, Queensland,  
Australia, plants of the new *Ozothamnus* differed from  
plants of the cultivar Just Blush in the following characteris-  
tics:

1. Plants of the new *Ozothamnus* were taller and more  
upright than plants of the cultivar Just Blush.
2. Plants of the new *Ozothamnus* and the cultivar Just  
Blush differed in involucre bract color as plants of the  
cultivar Just Blush had pink-colored involucre bracts.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographs illustrate the overall  
appearance of the new *Ozothamnus*. The photographs show  
the colors as true as it is reasonably possible to obtain in  
colored reproductions of this type. Colors in the photographs  
may differ slightly from the color values cited in the detailed  
botanical description which accurately describe the colors of  
the new *Ozothamnus*.

The photograph at the bottom of the sheet comprises a  
side perspective view of a typical flowering plant of '13-15'  
grown in a container.

The photograph at the top of the sheet is a close-up view  
of a typical corymb of '13-15'.

## DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. The aforementioned photographs and following observations and measurements describe plants grown in Bonsall, Calif. during the autumn in an outdoor nursery and under conditions and practices which approximate those generally used in commercial *Ozothamnus* production. During the production of the plants, day temperatures ranged from about 18° C. to 38° C. and night temperatures ranged from 13° C. to 24° C. Measurements and numerical values represent averages for typical flowering plants. Plants were about five months old when the photographs and description were taken.

Botanical classification: *Ozothamnus diosmifolius* cultivar 13-15.

Parentage: Self-pollination of a unnamed selection of *Ozothamnus diosmifolius*, not patented.

Propagation:

*Type*.—By vegetative cuttings.

*Time to initiate roots, summer*.—About two to three weeks.

*Time to produce a rooted young plant, summer*.—About six to eight weeks.

*Root description*.—Fine; fibrous; white in color.

*Rooting habit*.—Moderate branching; moderately dense.

Plant description:

*Plant form/growth habit*.—Upright plant habit; inverted triangle. Vigorous growth habit. Short internodes; dense and bushy habit. Freely branching habit with about five to six lateral branches per plant with numerous secondary and tertiary lateral branches.

*Plant height*.—About 46 cm.

*Plant diameter or spread*.—About 45 cm.

*Lateral branches*.—Length: About 60 cm. Diameter: About 6 mm. Internode length: About 4 cm. Aspect: Upright to somewhat outwardly spreading. Strength: Strong. Texture, developing: Pubescent. Texture, mature: Woody. Color, developing: 146C. Color, mature: 199A.

Foliage description:

*Arrangement*.—Alternate to whorled; simple; sessile.

*Length*.—About 1.8 cm.

*Width*.—About 1 mm.

*Shape*.—Acicular.

*Apex*.—Mucronate.

*Base*.—Attenuate, clasping.

*Margin*.—Entire.

*Texture, upper surface*.—Rough, granular.

*Texture, lower surface*.—Pubescent.

*Venation pattern*.—Single midvein.

*Color*.—Developing foliage, upper surface: 137A. Developing foliage, lower surface: 148B. Fully expanded foliage, upper surface: 147A; venation, 147A. Fully expanded foliage, lower surface: 148B; venation, 148B.

Inflorescence description:

*Appearance*.—Composite inflorescence form without ray florets; inflorescences arranged in terminal and axillary corymbs; corymbs hemispherical. Inflorescences face upright to outward. Freely flowering habit; about 50 inflorescences and inflorescence buds per corymb. Inflorescences persistent. Inflorescences faintly fragrant, straw-like.

*Time to flower*.—Under mild weather conditions, plants will flower year-round in Southern California. Inflorescences last about three to four weeks on the plant.

*Inflorescence bud*.—Height: About 3 mm. Diameter: About 3 mm. Shape: Rounded. Color: 155A.

*Inflorescence size*.—Diameter: About 3.2 cm to 3.8 mm. Depth (height): About 4 cm. Disc diameter: About 3 mm. Receptacle diameter: About 4 mm. Receptacle height: About 4 mm.

*Ray florets*.—Ray floret development has not been observed.

*Disc florets*.—Shape: Tubular; apex dentate. Length: About 4 mm. Diameter: About 1 mm. Number of disc florets per inflorescence: About 38. Color, immature and mature: Apex: Close to 155A. Mid-section and base: Close to 145D.

*Phyllaries*.—Quantity per inflorescence: About 36 arranged in about five to six whorls. Length: About 3 mm. Width: About 2 mm. Shape: Ovate. Apex: Acute. Base: Truncate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, immature, upper and lower surfaces: Close to 155D tinted with 65C. Color, mature, upper and lower surfaces: Close to 155D.

*Peduncles*.—Length: About 1.5 cm to 2 cm. Diameter: About 1 mm. Strength: Strong. Aspect: Upright to about 45°. Texture: Pubescent. Color: Close to 145A.

*Reproductive organs*.—Androecium: Quantity per disc floret: Five. Anther shape: Lanceolate. Anther length: About 1.5 mm. Anther color: Close to 155A. Filament length: About 2 mm. Filament color: Close to 155A. Pollen amount: Scarce. Pollen color: Close to 155A. Gynoecium: Quantity per disc floret: One. Pistil length: About 3 mm. Stigma shape: Bi-parted. Stigma color: Close to 155A. Style length: About 1.5 mm. Style color: Close to 155A. Ovary color: Close to 145C.

*Seed/fruit*.—Seed and fruit development have not been observed.

Disease/pest resistance: Plants of the new *Ozothamnus* have not been shown to be resistant to pathogens and pests common to *Ozothamnus*.

Garden performance: Plants of the new *Ozothamnus* have exhibited good tolerance to rain and wind and have been observed to tolerate temperatures from about -5° C. to about 40° C.

It is claimed:

1. A new and distinct *Ozothamnus* plant named '13-15' as illustrated and described.

\* \* \* \* \*

