



US00PP19230P3

(12) **United States Plant Patent**
Houbraeken(10) **Patent No.:** US PP19,230 P3
(45) **Date of Patent:** Sep. 16, 2008(54) **ARGYRANTHEMUM FRUTESCENS PLANT
NAMED 'ARGYPIFRI'**(50) Latin Name: *Argyranthemum frutescens*
Varietal Denomination: Argypifri(75) Inventor: **Anna M. W. P. Houbraeken**, Enkhuizen
(NL)(73) Assignee: **Syngenta Seeds B.V.**, Enkhuizen (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1 day.

(21) Appl. No.: **11/641,491**(22) Filed: **Dec. 19, 2006**(65) **Prior Publication Data**

US 2008/0184443 P1 Jul. 31, 2008

(51) **Int. Cl.****A01H 5/00** (2006.01)(52) **U.S. Cl.** **Plt./406; Plt./263**(58) **Field of Classification Search** **Plt./406,
Plt./263**

See application file for complete search history.

Primary Examiner—Annette H Para*Assistant Examiner*—S. B. McCormick-Ewoldt(74) *Attorney, Agent, or Firm*—S. Matthew Edwards(57) **ABSTRACT**

A new and distinct cultivar of *Argyranthemum frutescens* plant named 'Argypifri' characterized by its horizontal growing habit, pink, white flower heads with frizzled flower shape, and good flowering performance under hot conditions.

1 Drawing Sheet**2**

The following characteristics have been repeatedly observed and are determined to be basic characteristics of 'Argypifri' and distinguish it from new *Argyranthemum* as a new and distinct cultivar:

1. Horizontal growing character
2. Freely branching habit and short internodes; dense and bushy plant habit.
3. Early flowering habit
4. Freely flowering habit with numerous inflorescence per plant.
5. Frizzled flower shape.

TABLE 1**DIFFERENCES BETWEEN THE NEW CULTIVAR 'ARGYPIFRI'
AND TWO SIMILAR CULTIVARS**

	'Argypifri'	'Summer Stars Pink' (U.S. Plant Patent No 11,939)	'Daisy Crazy Blazer Rose' (Not patented)
Plant height	25 cm	35 cm	35 cm
Plant width	45 cm	40 cm	60 cm
Flower diameter	3.5-4 cm	3.5-4 cm	2.5 cm
Foliage color	137B	137C	146A
Flower color	75A	72D	67A

DESCRIPTION OF THE PHOTOGRAPHS

The accompanying color photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Argyranthemum*.

DESCRIPTION OF THE NEW CULTIVAR

The aforementioned photographs, following detailed observations and averaged measurements describe plants of

SUMMARY OF THE INVENTION

The new *Argyranthemum* has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype.

this new Marguerite Daisy plant. The data were collected from plants from asexual reproductions carried out in Enkhuizen, The Netherlands. The plant history was taken on 28 week old plants, grown in the open field in Enkhuizen, The Netherlands with day temperatures ranging from 16 to 35 degrees C., and night temperatures ranging from 10 to 20 degrees C.

Color references are primarily to the R.H.S. Colour Chart of The Royal Horticultural Society, 1995 Edition, except where general terms of ordinary dictionary significance are used.

The plant:

Classification.—Botanical : *Argyranthemum frutescens* cv. ‘Argypifri’.

Parentage:

Female parent.—Proprietary seedling section of *Argyranthemum frutescens*, identified as number ‘D0269-1’ (not patented). Breeder’s code of the new plant is ‘G0132-4.’

Male parent.—Unidentified section of *Argyranthemum frutescens* (not patented).

Propagation:

Type cuttings.—Terminal vegetative cuttings.

Time to initiated roots.—7–10 days at air temperature of 21 degrees C.

Time to develop roots.—7–14 days at air temperature of 18 degrees C.

Root description.—Fibrous, relatively fine, white in color.

Rooting description.—Freely branching, dense.

Plant description:

Growth habit.—Upright and rounded shape.

Plant height.—About 25 cm.

Vigor.—Vigorous compact.

Spreading area of plant.—About 45 cm.

Strength.—Very good.

Branching character.—Freely branching, plants do not require pinching.

Crop time.—About 9 weeks are required to produce a finished flowering plant in a 10.5 cm container from a good developed cutting.

Lateral branch description:

Length.—18 cm.

Diameter.—4 mm.

Shape.—Round, bit grooved.

Anthocyanin pigmentation.—Absent.

Texture.—Smooth, glabrous.

Length of internode.—1–3 cm.

Pubescence.—Absent.

Color.—144A.

The foliage:

Arrangement.—Alternate, simple.

Shape of leaf.—Bipinnatisect.

Leaf apex.—Broadly acute.

Leaf base.—Attenuate, clasping.

Margin.—Deeply dissected, laciniate.

Texture.—Thick glabrous smooth.

Length.—5–7 cm.

Width.—2–3.5 cm.

Depth of incision.—Lanciniate.

Color.—Upper side : 137A. Lower side : 137C. Old leaf : 137A.

Pubescence.—Absent.

Petiole.—Length : About 2–2.5 cm. Diameter : About 2 mm. Texture : Smooth, glabrous. Color : 137B.

Venation.—Shape : Pinnate. Color : 137A.

Inflorescence description:

Appearance.—Daisy-type inflorescences with ligulate ray florets. Disc and ray florets develop acropetally on a capitulum. Inflorescences held upright and outwardly on terminal and axillary peduncles. Inflorescences positioned perpendicular to the peduncles. Inflorescences persistent.

Flowering response.—Under natural conditions, plant flower from spring to early Fall in N-W Europe; plants flower continuous during this period.

Inflorescence longevity.—Inflorescences last about ten days on the plant.

Quantity of inflorescences.—Freely flowering, about 500 open inflorescences and inflorescence buds per plant.

Flower bud:

Peduncle.—Length : 7–10 cm. Diameter : 1.5 mm. Strength : Strong. Aspect : Upright to slightly outward. Texture : Smooth, glabrous. Color : 137C.

Inflorescence bud.—Height : 5 mm. Diameter : 7 mm. Shape : Ovoid. Color : 199A.

Inflorescence:

Size.—Diameter : 3.5–4 cm. Depth : About 6 mm.

Diameter of disc.—About 0.4–1 cm.

Form.—Semi double flowered.

Receptacle.—Height : About 4 mm. Diameter : About 10 mm.

Involucre/phyllaries.—3 series of 5 bracts, tightly to receptacle. Appearance : Margins, papery and membranous. Shape: Elliptic. Apex : Broad. Base : Truncate. Texture : Smooth glabrous. Margin : Entire. Length : 2 mm. Width : 1 mm. Color : Upper N144C, lower 139B.

Fragrance.—Absent.

Ray florets:

Shape of ray florets.—Ligulate.

Apex.—Emarginate.

Base.—Attenuate.

Margin.—Entire.

Color.—Upper surface : 75A. Lower surface : 75B.

No. of ray florets per inflorescence.—20–30 arranged in many whorls which give a frizzled look.

Size.—Length : 0.5–1.4 cm. Width : 0.2–0.4 cm.

Texture.—Smooth, glabrous; longitudinally ridged.

Disc florets:

Arrangement.—Massed at the center of the inflorescence.

Shape of disc florets.—Tubular, elongated, five-pointed; base, fused.

Color.—Immature : 14B. Mature : 14B.

Diameter of disc floret.—Apex : 1 mm. Base : <1 mm.

Length of disc floret.—4 mm.

Quantity per inflorescence.—Numerous, About 40.

Androecium.—Present only on disc florets.

Stamen quantity per floret.—Five.

Anther length.—About 1 mm.

Anther shape.—Oval.

Anther color.—Yellow 21A.

Pollen color.—Yellow 21A.

Amount of pollen.—Only in disc florets, scarce pollen.

Gynoecium.—Present on ray and disc florets.

Pistil number.—One per floret.

Pistil length.—Average 3 mm.

US PP19,230 P3

5

Style color.—Yellow green 144D.

Style length.—About 1 mm.

Stigma color.—Yellow orange 14A.

Stigma shape.—Bilobed.

Ovary color.—144C.

Seed/fruit: Seed and fruit production is not observed.

Disease resistance: No susceptibility nor resistance to fungal, bacterial or viral pathogens has been noted. Material is free of virus and Chrysanthemum stunt viroid (CSVd).

6

Temperature/weather tolerance: Plants of the new *Argyranthemum* have been observed to be tolerant to rain, wind and to temperatures from 0° C. to 36° C.

What is claimed is:

1. A new and distinct variety of *Argyranthemum frutescens* plant, named 'Argypifri,' substantially as herein illustrated and described.

* * * * *

U.S. Patent

Sep. 16, 2008

US PP19,230 P3



UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : PP 19,230 P3
APPLICATION NO. : 11/641491
DATED : September 16, 2008
INVENTOR(S) : Anna M.W.P. Houbraken

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

At column 2, line 3, after “distinguish”, delete “to” and insert therefor --the--

At column 2, line 27, delete “PHOTOGRAPHS” and insert therefor --PHOTOGRAPH--


At column 2, line 28, delete “photographs” and insert therefor --photograph--

At column 2, line 37, delete “photographs” and insert therefor --photograph--

At column 4, after “Flowering response.-Under natural conditions,” delete “plant” and insert therefor --plants--

Signed and Sealed this

Eleventh Day of November, 2008



JON W. DUDAS
Director of the United States Patent and Trademark Office