

(12) **United States Plant Patent**
Seidler

(10) **Patent No.:** **US PP19,468 P2**
(45) **Date of Patent:** **Nov. 18, 2008**

(54) **HYPERICUM PLANT NAMED 'SEIFIRE'**

(50) Latin Name: *Hypericum androsaemum*
Varietal Denomination: **Seifire**

(76) Inventor: **Thomas Seidler**, Ramskamp 64,
DE-25337, Elmshorn (DE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 4 days.

(21) Appl. No.: **11/809,983**

(22) Filed: **Jun. 4, 2007**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./442**

(58) **Field of Classification Search** Plt./263.1,
Plt./226, 442

See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

European Plant Breeders' Rights No. 17902 granted Jun. 19,
2006. Plant material first became available to the public on
Aug. 11, 2006 in the form of a European sale.

Primary Examiner—Kent L Bell

(74) *Attorney, Agent, or Firm*—Audrey Charles

(57) **ABSTRACT**

A new and distinct cultivar of *Hypericum* plant named
'Seifire', characterized by its yellow-colored flowers, large,
red-colored fruit, dark green-colored foliage, and vigorous,
upright growth habit.

1 Drawing Sheet

1

Latin name of genus and species of plant claimed: *Hypericum androsaemum*.

Variety denomination: 'Seifire'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Hypericum* plant botanically known as *Hypericum*
androsaemum, used commercially as stems with fruit and
hereinafter referred to by the cultivar name 'Seifire'.

The new cultivar originated in a controlled breeding pro-
gram in Elmshorn, Germany during July 2001. The objective
of the breeding program was the development of *Hypericum*
cultivars with unique and large attractive fruit coloration.

The new *Hypericum* cultivar is the result of open-
pollination using a bulk of proprietary female and male
Hypericum androsaemum parent plants, not patented, char-
acterized by large brown-colored berries or small berries
ranging in color from red to pink. The new cultivar was
discovered and selected as a single flowering plant within
the progeny of the above stated open-pollination during June
2002 in a controlled environment at Elmshorn, Germany.

Asexual reproduction of the new cultivar by terminal stem
cuttings since June 2002 at Elmshorn, Germany and Arroyo
Grande, Calif. has demonstrated that the new cultivar repro-
duces true to type with all of the characteristics, as herein
described, firmly fixed and retained through successive gen-
erations of such asexual propagation.

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have
been repeatedly observed and can be used to distinguish
'Seifire' as a new and distinct cultivar of *Hypericum* plant:

1. Yellow-colored flowers;
2. Large, red-colored fruit;
3. Dark green-colored foliage; and
4. Vigorous, upright growth habit.

2

Plants of the new cultivar differ from plants of the female
and male parents primarily in fruit size and coloration.

Of the many commercially available *Hypericum* cultivars
known to the inventor, the most similar in comparison to the
new cultivar is 'Excellent Flair', not patented. However, in
side by side comparisons, plants of the new cultivar differ
from plants of 'Excellent Flair' in the following characteris-
tics:

1. Plants of the new cultivar are shorter than plants of
'Excellent Flair';
2. Plants of the new cultivar have a leaf color different
from plants of 'Excellent Flair'; and
3. Plants of the new cultivar have fruit color different from
plants of 'Excellent Flair'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it
is reasonably possible to make the same in color illustrations
of this type, typical flower and foliage characteristics of the
new cultivar. Colors in the photographs differ slightly from
the color values cited in the detailed description, which
accurately describes the colors of 'Seifire'. The plants were
grown outdoors in the ground for 12 months at Arroyo
Grande, Calif.

FIG. 1 illustrates a side view of flowers, fruits, and the
overall growth habit of 'Seifire'.

FIG. 2 illustrates a close-up view of the flower of 'Seifire'.

FIG. 3 illustrates a close-up view of the fruit of 'Seifire'.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible
environmental conditions to date. Accordingly, it is possible
that the phenotype may vary somewhat with variations in the
environment, such as temperature, light intensity, and day
length, without, however, any variance in genotype.

The chart used in the identification of colors described
herein is The R.H.S. Colour Chart of The Royal Horticul-

tural Society, London, England, 2001 edition, except where general color terms of ordinary significance are used. The color values were determined on Aug. 18, 2004 between 10:00 a.m. and 12:00 p.m. under natural light conditions in Elmshorn, Germany when outdoor temperatures typically range from 82° F. (28° C.) during the day to 53° F. (12° C.) at night.

The following descriptions and measurements describe plants grown in an outdoor nursery in Elmshorn, Germany and under commercial production practices. Plants were about two years old.

Botanical classification: *Hypericum androsaemum* cultivar Seifire.

Parentage:

Female parent.—A proprietary mix of *Hypericum androsaemum* plants, not patented.

Male parent.—A proprietary mix of *Hypericum androsaemum* plants, not patented.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 14 to 21 days.

Time to produce a rooted cutting.—Approximately 2 months.

Root description.—Fine, white.

Rooting habit.—Freely branching, dense. Height of entire root structure: Approximately 50 cm. Diameter of entire root structure: Approximately 100 cm.

Plant description:

Form.—Erect, perennial, woody shrub; broad, inverted triangle.

Growth habit and general appearance.—Vigorous, dense bushy.

Size.—Height from soil level to top of plant plane: Approximately 57 cm. Width: Approximately 50 cm to 60 cm.

Branching habit.—Freely basal branching. Quantity of lateral branches per plant: Approximately 25. Quantity of harvestable stems per year per plant: Approximately 19; production of stems increases with age of plant and remains constant after three years of growth.

Branch.—Shape: Rounded. Strength: Strong. Length: Approximately 57 cm. Diameter: Approximately 5.0 mm. Length of central internode: Approximately 7.4 cm. Texture: Smooth, glabrous. Color of mature stem: 166B.

Foliage description:

General description.—Quantity of leaves per main branch: Approximately 50. Form: Simple, sessile. Arrangement: Opposite.

Leaves.—Shape: Oval to lanceolate. Margin: Entire. Apex: Acute. Base: Cordate. Venation pattern: Pinnate. Length of mature leaf: Approximately 8.0 cm. Width of mature leaf: Approximately 3.9 cm. Texture of upper surface: Smooth, glabrous. Texture of lower surface: Smooth, glabrous. Color of upper surface of young foliage: 144A with venation of 144C. Color of lower surface of young foliage: 138B with venation of 144C. Color of upper surface of mature foliage: 139A with venation of 144C. Color of lower surface of mature foliage: Between 147B and 147C with venation of 144C.

Flowering description:

Flowering habit.—‘Seifire’ flowers throughout the summer under outdoor growing conditions and pro-

duces harvestable cut stems with fruit beginning in August; flowering period typically lasts about 60 days.

Lastingness of individual flower on the plant.—Approximately 14 days.

Lastingness of individual flower as a cut flower.—Approximately 12 days.

Inflorescence description:

General description.—Type: Compound umbel, not persistent. Quantity of flowers per inflorescence: Approximately 9. Fragrance: None. Length or height: Approximately 6.0 cm. Width: Approximately 4.5 cm.

Peduncle.—Strength: Strong. Aspect: Approximately 45° angle from stem. Length: Approximately 1.8 cm. Diameter: Approximately 1.5 mm. Texture: smooth, glabrous. Color: 144A with an overlay of 183C.

Flower description:

General description.—Type: Single. Aspect: Upright. Quantity per lateral stem: Approximately 50.

Bud.—Shape: Ovoid. Quantity per lateral stem: Approximately 50. Length: Approximately 9.0 mm. Diameter: Approximately 6.0 mm. Color: 14A.

Corolla.—Shape: Round. Diameter: Approximately 2.5 cm. Depth: Approximately 2.0 cm.

Petals.—Quantity: 5, nonimbricate. Shape: Oval. Appearance: Flat. Aspect: Slightly concave. Margin: Entire. Apex: Rounded. Base: Attenuate. Length: Approximately 1.5 cm. Width: Approximately 7.8 mm. Texture of upper and lower surfaces: Smooth, glabrous. Color of upper surface when first open: 13A. Color of lower surface when first open: 12A. Color of upper surface when fully open: 13A. Color of lower surface when fully open: 13B.

Calyx.—Shape: Divergate. Length: Approximately 4.0 mm. Width: Approximately 1.3 cm.

Sepals.—Quantity per flower: 5, nonimbricate. Shape: Oval. Appearance: Flat. Margin: Entire. Apex: Apiculate. Base: Broadly cuneate. Length: Approximately 7.4 mm. Width: Approximately 4.0 mm. Texture of upper and lower surfaces: Smooth, glabrous. Color of upper surface of immature sepals: 144A. Color of lower surface of immature sepals: 144B. Color of upper surface of mature sepals: 144A with an overlay of 59A. Color of lower surface of mature sepals: 144B with an overlay of 59A.

Pedicels.—Strength: Strong. Aspect: Approximately 45° angle from stem. Length: Approximately 2.6 cm. Diameter: 1.5 mm. Texture: Smooth, glabrous. Color: 144A with an overlay of 183C.

Reproductive organs.—Androecium: Stamen quantity: Approximately 85. Anther shape: Oval. Anther length: Approximately 0.5 mm. Anther color: 17C. Pollen amount: Moderate. Pollen color: 13B. Gynoecium: Pistil quantity: 3 per flower. Pistil length: Approximately 7.0 mm. Stigma shape:

Globular. Stigma color: 59A. Style length: Approximately 6.0 mm. Style color: 150C. Ovary color: 150B.

Fruit.—Type: Capsule. Taste: Bitter. Shape: Ellipsoidal. Quantity per lateral branch: Approximately 50. Quantity per plant: Greater than 500. Days to ripening: Approximately 10 days to color and 30 days to seed. Longevity on plant: Approximately 30 days. Length: Approximately 1.5 cm. Diameter: Approximately 1.1 cm. Texture: Smooth, glabrous. Color: 46B.

Seed.—Quantity per fruit: Typically more than 100.
Length: Approximately 1.0 mm. Diameter: Approximately 0.5 mm. Texture: Smooth, glabrous. Color 200D.
Disease and pest resistance: Plants of the new *Hypericum* have been observed to be resistant to rust.
Weather tolerance: Plants of the new *Hypericum* have been observed to tolerate temperatures ranging from −10° C. to

35° C., and to demonstrate tolerance to rain and wind effects.
What is claimed is:
1. A new and distinct cultivar of *Hypericum* plant named ‘Seifire’, substantially as herein shown and described.

* * * * *



FIG. 1




FIG. 2




FIG. 3