



US00PP19614P2

(12) **United States Plant Patent**  
**Heinrich**(10) **Patent No.:** US PP19,614 P2  
(45) **Date of Patent:** Jan. 6, 2009(54) **PHLOX PLANT NAMED 'WESPOSKY'**(50) Latin Name: *Phlox hybrida*  
Varietal Denomination: Wesposky(75) Inventor: **Westhoff Heinrich**, Südlohn (DE)(73) Assignee: **Gartenbau und Spezialkulturen**  
**Westhoff GbR**, Südlohn (DE)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/983,724**(22) Filed: **Nov. 12, 2007**(51) **Int. Cl.**  
**A01H 5/00** (2006.01)(52) **U.S. Cl.** ..... **Plt./320**(58) **Field of Classification Search** ..... Plt./320  
See application file for complete search history.*Primary Examiner*—Kent L. Bell*Assistant Examiner*—June Hwu(74) *Attorney, Agent, or Firm*—C. A. Whealy**(57) ABSTRACT**

A new and distinct cultivar of *Phlox* plant named 'Wesposky', characterized by its compact, upright and mounding plant habit; freely branching growth habit; freely and continuous flowering habit; long flowering period; white colored flowers with a light violet-colored central star pattern; and good garden performance.

**1 Drawing Sheet****1**

Botanical Designation: *Phlox hybrida*.  
Cultivar Denomination: 'Wesposky'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of *Phlox*, botanically known as *Phlox hybrida* and hereinafter referred to by the name 'Wesposky'. 5

The new *Phlox* is a product of a planned breeding program conducted by the Inventor in Südlohn, Germany. The objective of the breeding program was to create new compact and freely branching *Phlox* cultivars with numerous attractive flowers.

The new *Phlox* originated from a cross-pollination made by the Inventor in 2004, in Südlohn, Germany, of a proprietary selection of *Phlox hybrida* identified as code number 04P005, not patented, as the female, or seed, parent with a proprietary selection of *Phlox hybrida* identified as code number 04P010, not patented, as the male, or pollen, parent. The new *Phlox* was discovered and selected by the Inventor as a single flowering plant within the progeny of the stated cross-pollination in a controlled environment in Südlohn, Germany in 2005. 15

Asexual reproduction of the new *Phlox* by vegetative cuttings in a controlled environment in Südlohn, Germany since 2005 has shown that the unique features of this new *Phlox* are stable and reproduced true to type in successive generations. 20

**SUMMARY OF THE INVENTION**

The cultivar Wesposky has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment and cultural practices such as temperature and light intensity without, however, any variance in genotype. 35

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Wesposky'. These characteristics in combination distinguish 'Wesposky' as a new and distinct cultivar of *Phlox*: 40

**2**

1. Compact, upright and mounding plant habit.
2. Freely branching growth habit.
3. Freely and continuous flowering habit; long flowering period.
4. White-colored flowers with a light violet-colored central star pattern.
5. Good garden performance.

Plants of the new *Phlox* differ from plants of the female parent selection in the following characteristics:

1. Plants of the new *Phlox* are more freely branching than plants of the female parent selection.
2. Plants of the new *Phlox* have larger and lighter green-colored leaves than plants of the female parent selection.

Plants of the new *Phlox* differ from plants of the male parent selection in the following characteristics:

1. Plants of the new *Phlox* are more compact than plants of the male parent selection.
2. Plants of the new *Phlox* and the male parent selection differ in flower color as plants of the male parent selection have white-colored flowers.

Plants of the new *Phlox* can be compared to plants of the *Phlox* cultivar USPHL304, disclosed in U.S. Plant Pat. No. 17,899. In side-by-side comparisons conducted in Südlohn, Germany, plants of the new *Phlox* differed from plants of the cultivar USPHL304 in the following characteristics:

1. Plants of the new *Phlox* were more compact than plants of the cultivar USPHL304.
2. Plants of the new *Phlox* had stronger lateral branches than plants of the cultivar USPHL304.
3. Plants of the new *Phlox* had smaller flowers than plants of the cultivar USPHL304.
4. Plants of the new *Phlox* and the cultivar USPHL304 differed in flower color as plants of the USPHL304 had lavender-colored flowers with a white and purple-colored star pattern at the center.

Plants of the new *Phlox* can also be compared to plants of the *Phlox* cultivar Astoria White, not patented. In side-by-

side comparisons conducted in Südlohn, Germany, plants of the new *Phlox* differed from plants of the cultivar Astoria White in the following characteristics:

1. Plants of the new *Phlox* had stronger lateral branches than plants of the cultivar Astoria White.
2. Plants of the new *Phlox* had larger leaves than plants of the cultivar Astoria White.
3. Plants of the new *Phlox* and the cultivar Astoria White differed in flower color as plants of the Astoria White had white-colored flowers

#### BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Phlox*, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type.

Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Phlox*. The photograph comprises a side perspective view of a typical flowering plant of 'Wesposky' grown in a container.

#### DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown in Südlohn, Germany, under commercial practice during the summer in a glass-covered greenhouse with day temperatures ranging from 20° C. to 25° C., night temperatures ranging from 16° C. to 18° C. and light levels ranging from 3,000 lux 50,000 lux. Rooted young plants were grown for about 20 weeks when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used.

**Botanical classification:** *Phlox hybrida* cultivar Wesposky.

**Parentage:**

*Female, or seed, parent.*—Proprietary selection of *Phlox hybrida* identified as code number 04P005, not patented.

*Male, or pollen, parent.*—Proprietary selection of *Phlox hybrida* identified as code number 04P010, not patented.

**Propagation:**

*Type.*—By vegetative cuttings.

*Time to initiate roots, summer.*—About 10 to 14 days at temperatures of 20° C.

*Time to initiate roots, winter.*—About 16 to 18 days at temperatures of 20° C.

*Time to produce a rooted young plant, summer.*—About 21 to 24 days at temperatures of 20° C.

*Time to produce a rooted young plant, winter.*—About 24 to 26 days at temperatures of 20° C.

*Root description.*—Fibrous; color, close to 158A.

*Rooting habit.*—Freely branching; moderately dense to dense.

**Plant description:**

*Plant form/habit.*—Compact, upright and mounded plant habit; outwardly spreading; vigorous growth habit. Freely branching habit with lateral branches developing at every node; pinching enhances branching potential.

*Plant height.*—About 20 cm to 21 cm.

*Plant width (spread).*—About 36 cm.

*Lateral branches.*—Length: About 11.2 cm. Diameter: About 2.4 mm. Internode length: About 1.7 cm. Strength: Strong. Texture: Pubescent; viscid. Color: 144A to 144B.

**Foliage description:**

*Arrangement.*—Alternate, simple, sessile.

*Length.*—About 2.4 cm.

*Width.*—About 0.8 cm.

*Shape.*—Lanceolate.

*Apex.*—Acute.

*Base.*—Obtuse.

*Margin.*—Entire.

*Texture, upper and lower surfaces.*—Pubescent.

*Venation pattern.*—Pinnate.

*Color.*—Developing leaves, upper surface: 137A. Developing leaves, lower surface: 146B. Fully expanded leaves, upper surface: 146A; venation, 146A. Fully expanded leaves, lower surface: 146B; venation, 146B.

**Flower description:**

*Flower type/habit.*—Single salverform flowers arranged in terminal cymes; flowers face upward to outward. Freely flowering habit with about nine flowers per cyme.

*Fragrance.*—None detected.

*Natural flowering season.*—Long flowering period; continuously flowering from April to autumn in Germany. Flowers not persistent.

*Postproduction longevity.*—Flowers last about five days on the plant.

*Flower buds.*—Height: About 1.3 cm. Diameter: About 2.5 mm. Shape: Oblong. Color: 150C.

*Inflorescence height.*—About 6.9 cm.

*Inflorescence diameter.*—About 6.55 cm.

*Flower diameter.*—About 2.6 cm.

*Flower depth.*—About 1.64 cm.

*Petals.*—Quantity per flower: Typically five in a single whorl; petals fused at the base into a narrow tube. Lobe length: About 1.3 cm. Lobe width: About 1.2 cm. Shape: Obovate. Apex: Cuspidate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: Developing petal lobes, upper surface: 155A; central star, N92C and 94C. Developing petal lobes, lower surface: 11D. Fully expanded petal lobes, upper surface: 155A; central star, N92C and 94C; color becoming closer to 155D with central star, 71A and N74A with development. Fully expanded petal lobes, lower surface: Close to 155. Eye: Between N92C and 94C. Throat: 145C to 145D. Tube: 160D slightly flushed with N79B.

*Sepals.*—Quantity per flower: Typically five in a single whorl, fused at base; tubular in shape. Length: About 8.1 mm. Width: About 0.95 mm. Shape: Acicular. Apex: Acute. Margin: Entire. Texture, upper and lower surfaces: Densely pubescent. Color, upper and lower surfaces: 138A.

*Peduncles.*—Length: About 0.9 cm to 1.7 cm. Diameter: About 1.9 mm. Strength: Strong, flexible. Texture: Pubescent. Color: 144A to 144B.

*Pedicels.*—Length: About 2 mm to 9 mm. Diameter: About 1 mm. Strength: Strong, flexible. Texture: Pubescent. Color: 144B to 144C.

*Reproductive organs.*—Stamens: Quantity per flower: Typically five. Filament length: Less than 1 mm. Anther shape: Lanceolate. Anther length: About 2.2

US PP19,614 P2

**5**

mm. Anther width: About 0.1 mm. Anther color: 9A. Pollen amount: Abundant. Pollen color: 13A. Pistils: Quantity per flower: One. Pistil length: About 5 mm. Stigma shape: Three-parted. Stigma color: 145B; color becoming closer to 151C. Style length: Less than 1 mm. Style color: 144C. Ovary color: 144C.  
*Seeds.*—Quantity per flower: Typically three. Length: About 3.1 mm. Diameter: About 2 mm. Color: N200A.

**6**

Disease/pest resistance: Plants of the new *Phlox* have not been noted to be resistant to pathogens and pests common to *Phlox*.

Garden performance: Plants of the new *Phlox* have been observed to have good garden performance and tolerate rain, wind and temperatures from about 5° C. to about 30° C.

It is claimed:

1. A new and distinct *Phlox* plant named ‘Wesposky’ as illustrated and described.

\* \* \* \* \*

**U.S. Patent**

**Jan. 6, 2009**

**US PP19,614 P2**

