



US00PP19622P2

(12) **United States Plant Patent**
Kawashima et al.

(10) **Patent No.:** **US PP19,622 P2**
(45) **Date of Patent:** **Jan. 13, 2009**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED
'SAKIMPOO9'**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(50) Latin Name: ***Impatiens*×*hybrida* (hort)**
Varietal Denomination: **SAKIMP009**

(52) **U.S. Cl.** **Plt./318.7**

(58) **Field of Classification Search** **Plt./318.7**
See application file for complete search history.

(75) Inventors: **Moriya Kawashima**, Yokohama (JP);
Yoneo Kobayashi, Kakegawa (JP)

Primary Examiner—Annette H Para
Assistant Examiner—Georgia Helmer
(74) *Attorney, Agent, or Firm*—Jondle & Associates, P.C.

(73) Assignee: **Sakata Seed Corporation**, Yokohama
(JP)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(57) **ABSTRACT**

A New Guinea *Impatiens* cultivar particularly distinguished
by having red flowers with red-purple eyes, a strong root
system and a fairly compact plant habit, is disclosed.

(21) Appl. No.: **12/011,183**

(22) Filed: **Jan. 24, 2008**

1 Drawing Sheet

1

Genus and species: *Impatiens*×*hybrida* (hort).
Variety denomination: 'SAKIMP009'.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new and distinct culti-
var of New Guinea *impatiens*, botanically known as
Impatiens×*hybrida* (hort), and hereinafter referred to by the
cultivar name 'SAKIMP009'. 'SAKIMP009' originated
from an interspecific hybridization occurring in Misato,
Japan between the female parent, 'NC-1H1', an unpatented
proprietary *Impatiens* breeding line with light-orange flow-
ers and the male parent, 'NC-229', an unpatented propri-
etary *Impatiens* breeding line with pink-lilac flowers.

In January 2004, the female parent line, 'NC-1H1', and
male parent line, 'NC-229', were crossed and a population
of F₁ plants was created. The F₁ plants were evaluated in
Misato, Japan in an open field trial. The criteria for plant
selection included flower color, strong root system and a
compact plant growth habit. At the completion of the trial,
one single-plant selection was made based on the above cri-
teria and vegetatively propagated. From May to August
2005, the selection was evaluated in an open field in Misato,
Japan. Shoot-tip cuttings of the variety were then shipped to
Salinas, Calif., where the plants were regenerated and
reevaluated for stability of traits. The selection subsequently
was named 'SAKIMP009' and found to have its unique char-
acteristics reproduce true to type in successive generations
of asexual propagation.

SUMMARY OF THE INVENTION

The following are the most outstanding and distinguishing
characteristics of this new cultivar when grown under nor-
mal horticultural practices in Salinas, Calif.

1. Red flowers with red-purple eyes;
2. A fairly compact plant habit; and
3. A strong root system.

2

DESCRIPTION OF THE PHOTOGRAPHS

This new *impatiens* plant is illustrated by the accompany-
ing photographs which show overall plant habit including
blooms, buds, and foliage of the plant; the colors shown are
as true as can be reasonably obtained by conventional photo-
graphic procedures. The photographs are of plants that are 5
months from their transplanting date and 6 months from the
stick date.

FIG. 1 shows the overall plant habit including blooms,
buds and foliage.

FIG. 2 shows a close-up of the mature inflorescence.

**DETAILED DESCRIPTION OF THE NEW
CULTIVAR**

The following detailed descriptions set forth the distinc-
tive characteristics of 'SAKIMP009'. The data which define
these characteristics were collected from asexual reproduc-
tions carried out in Salinas, Calif. The plant history was
taken on plants grown for about four months from propaga-
tion by terminal cuttings under greenhouse conditions. Color
references are primarily to the R.H.S. Colour Chart of The
Royal Horticultural Society of London (R.H.S.), 4th edition
(2001). Anatomic labels are from The Cambridge Illustrated
Glossary of Botanical Terms, by M. Hickey and C. King,
Cambridge University Press.

DETAILED BOTANICAL DESCRIPTION

Classification:
Botanical.—*Impatiens* interspecific cross, *Impatiens*×
hybrida (hort).

Common name.—*Impatiens*.

Parentage:

Female parent.—'NC-1H1', an unpatented proprietary
Impatiens breeding line with light-orange flowers

Male parent.—'NC-229', an unpatented proprietary
Impatiens breeding line with pink-lilac flowers.

Growth:

Time to produce a rooted cutting.—The terminal 1.0 to
1.5 inches of an actively growing stem was excised.

The vegetative cuttings were propagated in five to six weeks. The base of the cuttings were dipped for 1 to 2 seconds in a 1:9 solution of Dip 'N Grow (1 solution:9 water) root inducing solution immediately prior to sticking into the cell trays. Cuttings were stuck into plastic cell trays having 98 cells, and containing a moistened peat moss-based growing medium. The cuttings were misted with water from overhead for 10 seconds every 30 minutes until sufficient roots were formed.

Environmental conditions for plant growth.—Rooted cuttings were transplanted and grown in 6-inch plastic pots in a glass greenhouse located in Salinas, Calif. Pots contained a peat moss-based growing medium. Soluble fertilizer containing 20% nitrogen, 10% phosphorus and 20% potassium was applied once a day or every other day by overhead irrigation. Plants were fertilized every 2–3 days, 2 times in consecutive applications and then given one clear water application. Pots were top-dressed with a dry, slow release fertilizer containing 14% nitrogen, 14% phosphorus and 14% potassium. The typical average air temperature was 24° C.

Time to bloom from propagation.—6 to 8 weeks.

Plant Description:

Habit.—Fairly compact.

Life cycle.—Tender perennial.

Height.—26.0 cm to 28.0 cm from soil line to top of foliage.

Spread.—48.0 cm to 50.0 cm.

Time to produce a rooted cutting.—About 4 weeks.

Flowering requirements.—Will flower so long as temperature is above 5° C.

Temperature tolerance.—Plants have been observed to continuously flower at a temperature range of 5° C.–36° C., with ability to withstand high heat and humidity.

Branches:

Number.—8 total with 5 main branches.

Length.—3.0 cm from soil line to first node; 20.0 cm to 22.0 cm total.

Diameter (main branch).—1.0 cm to 1.2 cm.

Color.—RHS 187B (greyed-purple).

Stems:

Length.—8.0 cm from first to second node, 12.0 cm to 14.0 cm total.

Diameter.—0.5 cm to 0.6 cm.

Internode length.—4.0 cm to 4.5 cm.

Color.—RHS 187C (greyed-purple).

Stem description.—Strong; circular cross-section, smooth and shiny.

Pubescence.—Absent.

Anthocyanin color.—RHS 187C (greyed-purple).

Leaves:

Arrangement.—Whorled with up to 5 leaves per node; opposite if only two leaves at one node.

Length.—9.0 cm to 10.0 cm.

Width.—4.0 cm.

Shape.—Lanceolate, curled.

Apex.—Acuminate.

Base.—Attenuate.

Margin.—Ciliate.

Texture.—Dull, waxy.

Color.—Upper surface: RHS 147A (yellow-green).

Lower surface: RHS 147B (yellow-green).

Variation.—Absent.

Fragrance.—Absent.

Pubescence.—Absent.

Venation.—Pinnate.

Venation color.—Upper surface: RHS 147D (yellow-green). Lower surface: RHS 185D (greyed-purple).

Petioles.—Length: 1.0 cm to 1.5 cm. Diameter: 0.5 cm to 0.6 cm. Color: RHS 185C (greyed-purple). Texture: Smooth, glabrous.

Flower buds:

Shape.—Deltoid, longitudinal cross-section.

Length.—1.3 cm to 1.5 cm.

Diameter.—1.5 cm.

Color.—RHS 52B (red).

Texture.—Glabrous.

Inflorescence:

Inflorescence type.—Single flower with spur.

Number of flowers per node.—1 to 2 in bloom at one time; about 5 flower buds per node.

Number of flowers per plant.—25 at one time.

Lastingness of individual blooms on the plant.—14 days.

Fragrance.—Absent.

Peduncles:

Length.—5.5 cm to 6.0 cm.

Diameter.—0.2 cm to 0.22 cm.

Color.—RHS 185D (greyed-purple).

Texture.—Smooth, glabrous.

Corolla:

Shape.—Roughly circular with 5 radial petals.

Diameter.—About 6.3 cm to 6.5 cm.

Depth.—0.2 cm to 0.3 cm.

Petals:

Length.—3.5 cm.

Width.—3.3 cm to 3.5 cm.

Shape.—Obovate.

Apex.—Emarginate (cleaved).

Base.—Attenuate.

Margin.—Entire.

Texture.—Glabrous.

Color.—Upper surface: RHS 52C (red). Lower surface: RHS 58D (red-purple). Eye zone: RHS 61A (red-purple).

Spur:

Shape.—Tubular and curved downward.

Color.—RHS 52B (red).

Length.—6.5 cm.

Diameter.—0.2 cm.

Sepals:

Shape.—Lanceolate.

Number.—Two.

Color.—RHS 144D (yellow-green).

Length.—1.0 cm to 1.2 cm.

Diameter.—1.0 cm.

Apex.—Caudate.

Base.—Subcordate.

Margin.—Entire.

Texture.—Glabrous.

Reproductive organs:

Stamens.—Form: Fused; split into 4 lobes. Number: Many. Filament length: 0.4 cm. Filament color: RHS 52D (red). Anther length: 0.5 cm. Anther color: RHS N155A (white). Pollen amount: Abundant. Pollen color: RHS N155A (white). Pollen description: Powdery.

5

Pistil.—Number: 5. Stigma color: RHS 137A (green).

Style color: RHS 137A (green).

Ovary arrangement.—Parietal.

Ovary surface color.—RHS 144B (yellow-green).

Fruit and seed set: No seed set observed.

Disease and insect resistance: No particular resistance or susceptibility has been observed.

COMPARISON WITH PARENTAL AND KNOWN CULTIVARS

‘SAKIMP009’ differs from the female parent, ‘NC-1H1’, an unpatented proprietary *Impatiens* plant, in that ‘SAKIMP009’ has red (upper surface) and red-purple (lower surface) petals with red-purple eye zones, while ‘NC-1H1’ has light orange flowers. Additionally, ‘SAKIMP009’ has a fairly compact plant habit, while ‘NC-1H1’ has a branching plant habit.

‘SAKIMP009’ differs from the male parent, ‘NC-229’, an unpatented proprietary *Impatiens* plant in that ‘SAKIMP009’ has red (upper surface) and red-purple (lower surface) petals with red-purple eye zones, while ‘NC-229’

6

has pink-lilac flowers. Additionally, ‘SAKIMP009’ has a fairly compact plant habit, while ‘NC-229’ has a branching plant habit.

‘SAKIMP009’ is similar to the commercial *Impatiens* variety ‘Misato FG2’ (U.S. Plant Pat. No. 17,663) however, there are differences as listed in the table below:

TABLE 1

Comparison of Characteristics between ‘SAKIMP009’ and ‘Misato FG2’		
Characteristic	‘SAKIMP009’	‘Misato FG2’
Growth habit	Fairly compact	Upright and branching
Petal color, upper surface	RHS 52C (red)	RHS N30C (orange-red)
Spur color	RHS 52B (red)	RHS 63A (red-purple) at the base and fading to RHS 62D (red-purple) at the tip

I claim:

1. A new and distinct cultivar of New Guinea *Impatiens* plant as shown and described herein.

* * * * *



Fig. 1



Fig. 2