



US00PP19673P3

(12) **United States Plant Patent**
Ferguson et al.

(10) **Patent No.:** **US PP19,673 P3**
(45) **Date of Patent:** **Feb. 3, 2009**

(54) **STRAWBERRY PLANT NAMED**
'DRISSTRAWTHREE'

(58) **Field of Classification Search** Plt./209
See application file for complete search history.

(50) Latin Name: *Fragaria×ananassa*
Varietal Denomination: **DrisStrawThree**

(56) **References Cited**

(75) Inventors: **Michael D. Ferguson**, Moorpark, CA
(US); **Bruce D. Mowrey**, Watsonville,
CA (US)

U.S. PATENT DOCUMENTS

PP11,548 P * 10/2000 Amorao et al. Plt./209
PP12,889 P2 * 8/2002 Lamb et al. Plt./364
PP14,771 P3 * 5/2004 Amorao et al. Plt./209
PP18,878 P2 * 6/2008 Mowrey et al. Plt./209

(73) Assignee: **Driscoll Strawberry Associates, Inc.**,
Watsonville, CA (US)

OTHER PUBLICATIONS

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 47 days.

UPOV Rom Gtitm Compuyer Database, GTI Jouve
Reteieval Software, Feb. 2008 Citations for 'DrisStraw-
Three'.*

* cited by examiner

(21) Appl. No.: **11/821,016**

Primary Examiner—Wendy C. Haas

(22) Filed: **Jun. 21, 2007**

(74) *Attorney, Agent, or Firm*—Jondle & Associates, P.C.

(65) **Prior Publication Data**

US 2008/0320625 P1 Dec. 25, 2008

(57) **ABSTRACT**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

This invention relates to a new and distinct cultivar of straw-
berry plant named 'DrisStrawThree'. The new cultivar is
primarily characterized by its large fruit size, heavy fruit
production, and resistance to powdery mildew.

(52) **U.S. Cl.** **Plt./209**

3 Drawing Sheets

1

2

Genus and species: *Fragaria×ananassa*.
Variety denomination: 'DrisStrawThree'.

DESCRIPTION OF THE NEW CULTIVAR

BACKGROUND OF THE NEW PLANT

The present invention relates to a new and distinct straw-
berry cultivar designated 'DrisStrawThree' and botanically
known as *Fragaria×ananassa*. This new strawberry cultivar
was discovered in October, 2002 and originated from a cross
between the female parent '5F205', a proprietary strawberry
plant (unpatented) and the male parent 'San Juan' (U.S.
Plant Pat. No. 12,899). The original seedling of the new
cultivar was asexually propagated at a nursery in Shasta
County, Calif. 'DrisStrawThree' was subsequently asexually
propagated and underwent further testing at a nursery in
Ventura County, Calif. for four years. The present invention
has been found to retain its distinctive characteristics
through successive asexual propagations.

The following description of 'DrisStrawThree' is based on
observations taken from the 2002 to 2006 growing seasons
in Ventura County, Calif. This description is in accordance
with UPOV terminology. Color designations, color
descriptions, and other phenotypical descriptions may deviate
from the stated values and descriptions depending upon
variation in environmental, seasonal, climatic and cultural
conditions. 'DrisStrawThree' has not been observed under
all possible environmental conditions. Color terminology
follows the Royal Horticultural Society Colour Chart, Lon-
don (R.H.S.) (2001).

DETAILED BOTANICAL DESCRIPTION

DESCRIPTION OF THE PHOTOGRAPHS

The accompanying color photographs show typical speci-
mens of the new cultivar at various stages of development as
nearly true as it is possible to make in color reproductions.

FIG. 1 shows overall plant habit including fruit at various
stages of development.

FIG. 2 shows leaves of the plant with three leaflets.

FIG. 3 shows both the upperside and underside of several
of the flowers.

FIG. 4 shows the whole fruit.

FIG. 5 shows the fruit in longitudinal cross-section.

Table 1 shows plant characteristics of the new variety
compared with plant characteristics of the parent 'San Juan'.
Plant characteristics include plant height, leaf shape in cross
section, fruit hollow center, fruit texture when tasted, and
reaction to aphids.

TABLE 1

Characteristic	'DrisStrawThree'	'San Juan'
Plant height (cm)	20.5	28.0
Leaf shape in cross	Slightly concave	Flat to slightly convex
Fruit hollow center	Small	Medium
Fruit texture when tasted	Very fine	Medium
<i>Aphis</i> spp. (Aphids)	Moderately resistant	Susceptible

Table 2 shows plant characteristics of the new variety compared with plant characteristics of the commercial varieties, 'Driscoll Camarillo' (U.S. Plant Pat. No. 14,771) and 'Baeza' (U.S. Plant Pat. No. 11,548). Plant characteristics include plant height, diameter, number of crowns per plant, habit, density of individual plant and vigor.

TABLE 2

Characteristic	'Driscoll Camarillo'	'Baeza'
Plant height (cm)	23.3	22.4
Plant diameter (cm)	43.9	44.7
Number of crowns/plant	3	3
Habit	Globose	Globose
Density of individual plant	Medium to Dense	Medium to Dense
Vigor	Strong	Medium

Table 3 shows leaf characteristics of the new cultivar compared with leaf characteristics of 'Driscoll Camarillo' and 'Baeza'. Leaf characteristics include terminal leaflet length and width in centimeters, length to width ratio, number of teeth per terminal leaflet, shape of teeth, color of upper and underside of leaf, leaf shape in cross section, leaf blistering, leaf glossiness, number of leaflets, leaflet margin, and shape of leaf base.

TABLE 3

Leaf Characteristic	'Driscoll Camarillo'	'Baeza'
Terminal leaflet length (cm)	0.90	0.83
Terminal leaflet width (cm)	0.97	0.91
Terminal leaflet length/width ratio	0.93	0.91
No. teeth/terminal leaflet	27	29
Shape of teeth	Rounded	Rounded
Color of upper side of leaf	RHS N189B	RHS 147A
Color of underside of leaf	RHS 137C	RHS 137C
Leaf shape in cross section	Slightly concave	Concave
Leaf blistering	Medium	Medium
Leaf glossiness	Weak	Weak
No. leaflets	Three only	Three only
Terminal leaflet margin	Revolute	Revolute
Terminal leaflet base shape	Rounded	Rounded

Table 4 shows information about the petiole, the petiolule, the bract, and the stipule of the new cultivar compared to 'Driscoll Camarillo' and 'Baeza'. This includes petiole length in centimeters, petiole diameter in centimeters, petiolule length in centimeters, petiolule diameter in centimeters, bract frequency per petiole, stipule length in centimeters, stipule width in centimeters, stipule pubescence, petiole pubescence, pose of hairs on the petiole, color of the petiole and color of the petiolule.

TABLE 4

Characteristic	'Driscoll Camarillo'	'Baeza'
Petiole length (cm)	12.7	13.4
Petiole diameter (cm)	0.408	0.420
Petiole pubescence	Sparse	Medium to dense
Petiole pose of hairs	Outwards	Outwards
Petiole color	RHS 144B	RHS 144B
Petiolule color	RHS 145B	RHS 145B
Petiolule length (cm)	1.255	1.456
Petiolule diameter (cm)	0.254	0.232
Bract frequency	1	1
Stipule length (cm)	4.0	3.7
Stipule width (cm)	1.070	1.256
Stipule pubescence	Sparse	Sparse

Table 5 shows stolon characteristics of the new cultivar compared to 'Driscoll Camarillo' and 'Baeza'. These characteristics include the number of stolons, the anthocyanin coloration of the stolons, the thickness of the stolons, and the pubescence of the stolons.

TABLE 5

Characteristic	'Driscoll Camarillo'	'Baeza'
Stolon Number	Few to medium	Few to medium
Stolon Anthocyanin	Strong	Weak to medium
Stolon Thickness	Medium	Thick
Stolon Pubescence	Sparse	Medium to dense

Table 6 shows inflorescence characteristics of the new cultivar compared to 'Driscoll Camarillo' and 'Baeza'. These characteristics include inflorescence position relative to foliage, relative flower size, flower diameter in centimeters (measured from petal tip to petal tip), relative spacing of petals, petal length in centimeters, petal width in centimeters, petal length to width ratio, petal color, calyx diameter in centimeters (measured on back of flower from sepal tip to sepal tip), diameter of calyx relative to corolla, diameter of inner calyx relative to outer, sepal length in centimeters (measured from sepal tip to point of attachment to receptacle), sepal width in centimeters, receptacle color and anther color.

TABLE 6

Characteristic	'Driscoll Camarillo'	'Baeza'
Inflorescence position relative to foliage	Above	Above
Flower size	Medium	Medium
Flower diameter (cm)	2.263	2.490
Petal spacing	Overlapping	Overlapping
Petal length (cm)	1.262	1.496

TABLE 6-continued

Characteristic	'DrisStrawThree'	'Driscoll Camarillo'	'Baeza'
Petal width (cm)	1.531	1.263	1.511
Petal length/width ratio	0.98	1.00	0.99
Petal color	RHS 155B	RHS 155C	RHS 155B
Calyx diameter (cm)	4.088	2.550	2.925
Calyx diameter relative to corolla	Larger	Smaller	Same size
Inner calyx diameter relative to outer	Same size	Same size	Same size
Sepal length (cm)	1.582	0.964	1.087
Sepal width (cm)	0.817	0.544	0.505
Receptacle color	RHS N144A	RHS 2B	RHS 1A
Anther color	RHS 17A	RHS 15B	RHS 17B

Table 7 shows fruit characteristics of the new cultivar compared to 'Driscoll Camarillo' and 'Baeza'.

TABLE 7

Characteristic	'DrisStrawThree'	'Driscoll Camarillo'	'Baeza'
Fruiting truss length (cm)	13.4	26.5	21.7
Fruiting truss attitude	Prostrate	Erect	Erect
Fruiting truss length	Medium	Long	Long
Fruit length (cm)	4.93	3.76	4.35
Fruit width (cm)	4.36	3.73	3.92
Fruit length/width ratio	1.13	1.01	1.11
Fruit weight (g)	27.5	21.0	22.3
Relative fruit size	Very large	Medium	Medium
Predominant fruit shape	Almost cylindrical	Conical	Conical
Difference in shape between primary & secondary fruits	Slight	None or very slight	None or very slight
Band without achenes	Narrow	Absent or very narrow	Absent or very narrow
Unevenness of fruit surface	Medium	Medium	Medium
Fruit skin color	RHS 45B Red	RHS 46A Red	RHS 46B Red
Evenness of fruit color	Uneven	Even	Even
Fruit glossiness	Medium	Medium	Medium
Insertion of achenes	Level with surface	Level with surface	Level with surface
Achene coloration-sunward side of berry	RHS 165A Red to purple, possibly greyed	RHS 166B Red to purple, possibly greyed	RHS 163B Red to purple, possibly greyed
Achene coloration-shaded side of berry	RHS 151C yellow to green	RHS 151C yellow to green	RHS 151A Yellow to green
Achenes per berry	243.6	316.2	361.3
Achene weight	0.000664409	0.000570367	0.000466731
Insertion of calyx	Set above the fruit	Level	Level
Pose of calyx segments	Reflexed	Reflexed	Reflexed

TABLE 7-continued

Characteristic	'DrisStrawThree'	'Driscoll Camarillo'	'Baeza'
Size of calyx in relation to fruit	Smaller to larger	Same size	Same size
Adherence of calyx	Strong	Strong	Strong
Firmness of flesh	Firm	Firm	Firm
Color of the flesh	RHS 155A (White) to RHS 44D (light red)	RHS 155A (White) to RHS 44D (light red)	RHS 155A (White) to RHS 38B (pale rose)
Evenness of flesh color	Slightly uneven	Even	Even
Distribution of flesh color	Marginal and central	Only marginal	Only marginal
Hollow center	Small	Absent	Medium
Sweetness	Strong	Medium	Medium
Acidity	Weak	Medium	Medium
Texture when tasted	Very fine	Medium	Medium
Time of flowering	Early	Early	Late
Harvest maturity	Mid-season, early June to late December	Mid-season, early June to late December	Late early July to late December
Type of bearing	Fully everbearing	Fully everbearing	Fully everbearing
Grams of fruit/plant	500	286	289

Table 8 shows the resistance to different forms of stress of the new cultivar compared to 'Driscoll Camarillo' and 'Baeza'. These forms of stress include drought, high temperatures, wind, high pH, and high soil salt level.

TABLE 8

Reaction to Stress	'DrisStrawThree'	'Driscoll Camarillo'	'Baeza'
Drought	Moderately resistant	Moderately resistant	Moderately resistant
High temperatures	Moderately resistant	Moderately resistant	Moderately resistant
Wind	Moderately resistant	Moderately resistant	Moderately resistant
High pH	Moderately resistant	Moderately resistant	Moderately resistant
High soil salt level	Moderately resistant	Moderately resistant	Susceptible

Table 9 shows pest and disease characteristics of the new cultivar compared to 'Driscoll Camarillo' and 'Baeza'.

TABLE 9

Pest or Disease	'DrisStrawThree'	'Driscoll Camarillo'	'Baeza'
<i>Tetranychus urticae</i> (2-spotted spider mite)	Moderately resistant	Moderately resistant	Moderately resistant
<i>Tarsonemus pallidus</i>	Moderately resistant	Moderately resistant	Moderately resistant
<i>Aphelencoides fragariae</i>	Moderately resistant	Moderately resistant	Moderately resistant
<i>Pratylenchus penetrans</i>	Moderately resistant	Moderately resistant	Moderately resistant
<i>Ditylenchus dipsac</i>	Moderately resistant	Moderately resistant	Moderately resistant

TABLE 9-continued

Pest or Disease	'DrisStrawThree'	'Driscoll Camarillo'	'Baeza'
<i>Anthonomus rubi</i>	Moderately resistant	Moderately resistant	Moderately resistant
<i>Aphis</i> spp. (Aphids)	Moderately resistant	Moderately resistant	Moderately resistant
<i>Lygus hesperus</i> (Lygus bug)	Moderately resistant	Moderately resistant	Moderately resistant
Botrytis fruit rot	Moderately Susceptible	Moderately Susceptible	Moderately Susceptible

TABLE 9-continued

Pest or Disease	'DrisStrawThree'	'Driscoll Camarillo'	'Baeza'
Powdery mildew	Resistant	Highly susceptible	Susceptible

We claim:

1. A new and distinct cultivar of strawberry plant as described and shown herein.

* * * * *



FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5