



US00PP19770P2

(12) **United States Plant Patent**
Bergman(10) **Patent No.:** US PP19,770 P2
(45) **Date of Patent:** Feb. 24, 2009

- (54) **CHrysanthemum PLANT NAMED 'CURRANT YOIRVINE'**
- (50) Latin Name: *Chrysanthemum×morifolium*
Varietal Denomination: **Currant Yorvine**
- (75) Inventor: **Wendy R. Bergman**, Lehigh Acres, FL
(US)
- (73) Assignee: **Yoder Brothers, Inc.**, Barberton, OH
(US)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **11/983,744**
- (22) Filed: **Nov. 12, 2007**
- (51) **Int. Cl.**
A01H 5/00 (2006.01)

- (52) **U.S. Cl.** **Plt./298**
- (58) **Field of Classification Search** Plt./298
See application file for complete search history.

Primary Examiner—Annette H Para
Assistant Examiner—June Hwu
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Chrysanthemum* plant named 'Currant Yorvine', characterized by its upright and uniformly mounded plant habit; freely branching and vigorous growth habit; dark green-colored foliage; uniform flowering response; early and freely flowering habit; daisy-type inflorescences with cherry coral-colored ray florets; and excellent postproduction longevity with inflorescences maintaining good substance and color for about four weeks in an interior environment.

1 Drawing Sheet**1**

Botanical designation: *Chrysanthemum×morifolium*.
Cultivar denomination: 'CURRANT YOIRVINE'.

CROSS-REFERENCE TO RELATED APPLICATIONS

Chrysanthemum Plant Named Dark Bronze Yorvine U.S. Plant Pat. application Ser. No. 11/983,733, Wendy R. Bergman, applicant; filed concurrently.

Chrysanthemum Plant Named Regal Yorvine U.S. Plant patent application Ser. No. 11/983,732; Wendy R. Bergman, applicant; filed concurrently.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Chrysanthemum* plant, botanically known as *Chrysanthemum×morifolium*, commercially grown as a pot-type *Chrysanthemum* and hereinafter referred to by the name 'Currant Yorvine'.

The new *Chrysanthemum* is a naturally-occurring whole plant mutation of the *Chrysanthemum×morifolium* cultivar Yorvine, not patented. The new *Chrysanthemum* was discovered and selected by the Inventor as a single flowering plant within a population of plants in a greenhouse environment of the cultivar Yorvine in December, 2003, in Fort Myers, Fla. The selection of this plant was based on its uniform plant growth habit, vigor, freely branching habit, desirable inflorescence form and floret colors, fast response time and excellent postproduction longevity.

Asexual reproduction of the new *Chrysanthemum* by vegetative tip cuttings was first conducted in Fort Myers, Fla. in March, 2004. Asexual reproduction by cuttings has shown that the unique features of this new *Chrysanthemum* are stable and reproduced true to type in successive generations.

2**SUMMARY OF THE INVENTION**

Plants of the cultivar Currant Yorvine have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature, daylength and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Currant Yorvine'. These characteristics in combination distinguish 'Currant Yorvine' as a new and distinct pot-type *Chrysanthemum* cultivar:

1. Upright and uniformly mounded plant habit.
2. Freely branching and vigorous growth habit.
3. Dark green-colored foliage.
4. Uniform flowering response.
5. Typically grown as a natural spray type.
6. Early flowering habit, eight-week response time.
7. Freely flowering habit.
8. Daisy-type inflorescences with cherry coral-colored ray florets.
9. Excellent postproduction longevity with inflorescences maintaining good substance and color for about four weeks in an interior environment.

Plants of the new *Chrysanthemum* differ from plants of the parent, the cultivar Yorvine, in the following characteristics:

1. Plants of the new *Chrysanthemum* flower a few days later than plants of the cultivar Yorvine.
2. Plants of the new *Chrysanthemum* and the cultivar Yorvine differ in ray floret color as plants of the cultivar Yorvine have light purple-colored ray florets.

Plants of the new *Chrysanthemum* differ from plants of the cultivars Dark Bronze Yorvine and Regal Yorvine primarily in ray floret color.

Plants of the new *Chrysanthemum* can be compared to plants of the *Chrysanthemum* cultivar Raspberry Yolompoc, disclosed in U.S. Plant Pat. No. 12,276. In side-by-side comparisons conducted in Fort Myers, Fla., plants of the new *Chrysanthemum* differed from plants of the cultivar Raspberry Yolompoc in the following characteristics:

1. Plants of the new *Chrysanthemum* were more vigorous than plants of the cultivar Raspberry Yolompoc.
2. Plants of the new *Chrysanthemum* had larger inflorescences than plants of the cultivar Raspberry Yolompoc.
3. Inflorescences of plants of the new *Chrysanthemum* and the cultivar Raspberry Yolompoc differed in ray floret color as plants of the cultivar Raspberry Yolompoc had darker-colored ray florets.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying photograph illustrates the overall appearance of the new *Chrysanthemum*. This photograph shows the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Chrysanthemum*. The photograph comprises a side perspective view of typical flowering plants of 'Currant Yoirvine' grown in a container.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations and measurements describe plants grown in Leamington, Ontario, Canada during the summer in a glass-covered greenhouse and under conditions and practices which approximate those generally used in commercial pot-type *Chrysanthemum* production. During the production of the plants, day temperatures ranged from about 21° C. to 27° C., night temperatures ranged from about 17° C. to 17.9° C. and light levels ranged from 4,000 to 6,000 foot candles. Four unrooted cuttings were directly stuck in 15-cm containers, exposed to long day/short night conditions, and pinched about three weeks later. At the time of the pinch, the photoinductive short day/long night treatments were started. Plants used in the photographs and for the description were center-budded and were about eleven weeks old. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Chrysanthemum* × *morifolium* cultivar Currant Yoirvine.

Parentage: Naturally-occurring whole plant mutation of the *Chrysanthemum* × *morifolium* cultivar Yoirvine, not patented.

Propagation:

Type.—Terminal vegetative cuttings.

Time to initiate roots.—About four days at temperatures of about 21° C.

Time to produce a rooted young plant.—About ten days at temperatures of about 21° C.

Root description.—Fine to thick, fibrous; white in color.

Rooting habit.—Freely branching; moderately dense.

Plant description:

Appearance.—Herbaceous daisy pot-type *Chrysanthemum* typically grown as a center-budded or as a natural spray type. Stems upright and outwardly spread-

ing giving a uniformly mounded appearance to the plant. Freely branching habit, about four lateral branches develop after removal of terminal apex (pinching); dense and full plant habit. Strong and vigorous growth habit.

Plant height.—About 28.5 cm.

Plant width.—About 42 cm.

Lateral branches.—Length: About 25 cm. Diameter: About 6 mm. Internode length: About 2.2 cm. Strength: Strong. Texture: Pubescent. Color: Between 144A and 146B.

Foliage description:

Arrangement.—Alternate, simple.

Length.—About 7.75 cm.

Width.—About 5.3 cm.

Shape.—Palmately lobed.

Apex.—Palmately lobed.

Apex.—Cuspidate to mucronate.

Base.—Attenuate with truncate tendencies.

Margin.—Palmately lobed, sinuses between lateral lobes parallel to divergent.

Texture, upper and lower surfaces.—Fine pubescence; veins prominent on lower surface.

Color.—Developing and fully expanded leaves, upper surface: Darker than 147A; venation, close to 147A to 147B. Developing and fully expanded leaves, lower surface: Close to 147B; venation, close to 147B.

Petiole.—Length: About 1.2 cm. Diameter: About 4 mm. Texture, upper and lower surfaces: Pubescent. Color, upper surface: Close to 147B. Color, lower surface: Close to 147B to 147C.

Inflorescence description:

Appearance.—Daisy-type inflorescence form with elongated oblong-shaped ray florets. Inflorescences borne on terminals above foliage. Disk and ray florets arranged acropetally on a capitulum. Inflorescence not fragrant. Typically grown as a natural spray type.

Flowering response.—Under natural conditions, plants flower in the autumn/winter in the Northern Hemisphere. At other times of the year, inflorescence initiation and development can be induced under short day/long night conditions (at least 13.5 hours of darkness). Early flowering habit; plants exposed to three weeks of long day/short night conditions followed by photoinductive short day/long night conditions flower about eight weeks later.

Postproduction longevity.—Inflorescences maintain good color and substance for about four weeks in an interior environment.

Quantity of inflorescences.—Freely flowering, about 12 inflorescences develop per lateral stem.

Inflorescence bud.—Height: About 5 mm. Diameter: About 1 cm. Shape: Oblate. Color: Close to 147A.

Inflorescence size.—Diameter: About 7.2 cm. Depth (height): About 2.5 cm. Diameter of disc: About 1 cm. Receptacle height: About 7 mm. Receptacle diameter: About 6 mm. Receptacle color: Close to 144B.

Ray florets.—Shape: Elongated oblong. Orientation: Initially upright, then with development, about 45° from vertical. Aspect: Initially incurved, then mostly flat. Length: About 3.7 cm. Width: About 1.1 cm. Apex: Acute. Base: Attenuate; short corolla tube. Margin: Entire. Texture, upper and lower surfaces:

Smooth, glabrous; satiny. Number of ray florets per inflorescence: About 42 arranged in about three whorls. Color: When opening and fully opened, upper surface: Close to 6A overlain with close to 59A. When opening and fully opened, lower surface: Close to 6A to 6B underlain with close to between 59A and 71A.

Disc florets.—Arrangement: Massed at center of receptacle. Shape: Tubular, elongated. Apex: Five-pointed. Length: About 4 mm. Width: About 1 mm. Number of disc florets per inflorescence: About 162. Color, immature: Apex: Close to 154B. Mid-section and base: Close to 155D. Color, mature: Apex: Close to 6A. Mid-section: Close to 154A. Base: Close to 155D.

Phyllaries.—Number of phyllaries per inflorescence: About 18 arranged in about two whorls. Length: About 1 cm. Width: About 4 mm. Shape: Lanceolate. Apex: Acute. Base: Truncate. Margin: Entire. Texture, upper surface: Smooth, waxy. Texture, lower surface: Pubescent. Color, upper surface: Close to 146A. Color, lower surface: Close to 147A.

Peduncles.—Length: First peduncle: About 6 cm. Fourth peduncle: About 8.5 cm. Seventh peduncle: About 12 cm. Diameter (first peduncle): About 2 mm. Angle: About 45° from vertical. Strength:

Strong, flexible. Texture: Pubescent. Color: Close to between 144A to 146B.

Reproductive organs.—Androecium: Present on disc florets only. Stamen length: About 4 mm. Filament length: About 3 mm. Filament color: Close to 154D. Anther shape: Narrowly oblong. Anther length: Less than 1 mm. Anther color: Close to 6A. Pollen amount: None observed. Gynoecium: Present on both ray and disc florets. Pistil length: About 5 mm. Stigma shape: Bi-parted. Stigma color: Close to 4A. Style length: About 4 mm. Style color: Close to 4A. Ovary color: Close to 155D.

Seed/fruit.—Seed and fruit production has not been observed.

Disease/pest resistance: Resistance to pathogens and pests common to *Chrysanthemums* has not been observed on plants grown under commercial conditions.

Temperature tolerance: Plants of the new *Chrysanthemum* tolerate temperatures ranging from about 5° C. to about 38° C.

It is claimed:

1. A new and distinct *Chrysanthemum* plant named ‘Currant Yoirvine’ as illustrated and described.

* * * * *

