

(12) **United States Plant Patent**
Walker

(10) **Patent No.:** **US PP19,993 P2**
(45) **Date of Patent:** **May 12, 2009**

(54) **NEMATODE RESISTANT GRAPE
ROOTSTOCK '9363-16'**

(50) Latin Name: *Vitis rufotomentosa* × (*V. champinii*)
Varietal Denomination: **9363-16**

(75) Inventor: **Andrew Walker**, Woodland, CA (US)

(73) Assignee: **The Regents of the University of
California**, Oakland, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/009,064**

(22) Filed: **Jan. 15, 2008**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./174**

(58) **Field of Classification Search** Plt./174,
Plt./205

See application file for complete search history.

Primary Examiner—Annette H Para

(74) *Attorney, Agent, or Firm*—Morrison & Foerster LLP

(57) **ABSTRACT**

Rootstock '9363-16' is a cross of 'L514-30' (*Vitis rufotomentosa* × (*V. champinii* 'Dog Ridge' × *V. riparia* 'Riparia Gloire')) × *V. riparia* 'Riparia Gloire'. The rootstock is a new and distinct variety of grape rootstock with resistance to a broad array of soil-borne nematodes including root-knot, dagger and lesion nematodes. The rootstock has moderate resistance to citrus nematodes but is susceptible to ring nematodes. The rootstock has strong resistance to grape phylloxera.

2 Drawing Sheets

1

Botanical/commercial classification: 'L514-30' (*Vitis rufotomentosa* × (*V. Champinii* 'Dog Ridge' × *V. riparia* 'Riparia Gloire')) × *V. riparia* "Riparia Gloire".
Variety Denomination: "'9363-16'".

BACKGROUND OF THE INVENTION

Nematodes are important pests in vineyards around the world, and these soil-borne pests can be particularly problematic in California vineyards. Two factors intensify the impact of nematodes — the high value of grapes and of vineyard land. These factors force growers to ignore the steps of leaving land fallow and rotating crops, both of which reduce nematode build up and delay the selection of adapted strains. Nematicides and fumigants help control nematodes, but the use of these pesticides has been greatly restricted and their future use in doubt because they must be persistent and penetrate deeply through the soil profile to be highly effective. In addition, grape rootstocks were bred to resist grape phylloxera (a devastating root-feeding aphid), and were not selected for nematode resistance. The grape rootstocks 'Freedom' and 'Harmony' were released by the USDA/Fresno to provide resistance to nematodes, but they are not resistant to grape phylloxera. More recently, McKenry at UC Riverside released two rootstocks, 'RS3' and 'RS9' (siblings of a 'Schwarzmann' × 'Ramsey' cross), designed to resist multiply nematode species and provide growers with nematode resistant rootstock alternatives. However, they have nematode resistance from a relatively narrow genetic base, which may promote the evolution of strains capable of feeding on them. Therefore, there is a need to develop nematode resistant grape rootstock.

SUMMARY OF THE INVENTION

This invention relates to a new and distinct variety of grape rootstock with resistance to a broad array of soil-borne nematodes including root-knot, dagger and lesion nematodes. The variety has moderate resistance to citrus nema-

2

todes but is susceptible to ring nematodes. The variety has strong resistance to grape phylloxera. The variety is a cross of 'L514-30' (*Vitis rufotomentosa* × (*V. champinii* 'Dog Ridge' × *V. riparia* 'Riparia Gloire')) × *V. riparia* 'Riparia Gloire'. The variety is recommended for vineyard sites with severe nematode infestations.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1: number of ring nematodes recovered off rootstock selections growing in 1 gal pots with three soils collected from the Gallo Livingston Ranch, known to have severe and chronic nematode pressure. Results per soils are means of three replicate pots.

FIG. 2: a photograph showing rootstock '9363-16'.

DETAILED DESCRIPTION OF THE INVENTION

A breeding program was initiated in 1990 at UC Davis to provide a group of rootstocks with broad and durable resistance to the nematodes found in California vineyards. This program commenced with an evaluation of selections that remained from breeding efforts in the late 1960s and early 1970s. These selections from the late 1960s and early 1970s were previously screened against two root knot nematode species (*Meloidogyne incognita acrita* and *M. arenaria thamesi*), the lesion nematode (*Pratylenchus vulnus*), and the dagger nematode (*Xiphinema index*). In 1990, these selections were evaluated for their ability to root and for growth habits such as bushy growth, internode length and the degree of lateral shoot production. In 1993 and 1994, the best of these selections were crossed to species chosen for their ability to reduce scion vigor or improve the rooting of the progeny. The parentage and species composition of '9363-16' and other crosses are listed in Table 1. About 5,000 progeny were planted in the vineyard and their evaluation for nematode resistance was initiated in 1996.

The first phase of the selection process examined the progeny for their general vigor and horticultural characters such as internode length and the degree of lateral shoot formation. The best 1,000 progeny, selected from a large number of, were advanced to a rooting assay. Ten 2-node dormant cuttings of the best 1,000 were taken in December 1996 and tested for their ability to form roots. Rootstocks that root well generally graft well, thus this evaluation was a key indicator of their future success as rootstocks. One hundred of the progeny were selected, again with an effort to get a broad representation from the large number of families.

The second phase of the selection process involved testing these 100 selections for resistance to *M. incognita* 13. This root-knot nematode isolate is capable of feeding on many rootstocks, but does not feed on rootstocks that derive their resistance from *V. champinii* ('Freedom', 'Harmony', 'Dog Ridge' and 'Ramsey'). All of the nematode testing in this breeding program utilized potted plants under greenhouse conditions with optimized soils and irrigation techniques to promote nematode feeding. Resistance to this isolate of root-knot nematode was evaluated by assaying the number of galls on the roots after inoculation with 1,000 J2 larvae (the free-living infectious stage of this nematode). Resistance was also evaluated by extracting the J2 nematodes in the pots after the root galls were counted. Later phases of the root-knot nematode screening evaluated resistance by counting the number of egg masses formed using a technique developed in the Walker lab (Cousins and Walker 2001 Plant Disease 85:1052–1054). There were no root galls on 33 of the 100 selections and the 33 selections were advanced to the next phase of screening.

The third phase of the selection process tested the 33 root-knot nematode resistant selections against two aggressive strains of root-knot nematode and against the dagger nematode, *X. index*. The two aggressive strains were selected in the Walker lab from soils provided by McKenry, from a declining 'Harmony' vineyard. Root-knot nematodes were extracted from this soil and the larvae were placed onto tomato plants with high susceptibility to root-knot nematodes. Two egg masses were collected from these infested plants and J2 larvae from each mass were put on a separate uninfested tomato plant to create new strains from a single egg mass (root-knot nematodes are parthenogenic). These new strains were multiplied on tomato and were then used to inoculate potted plants of 'Harmony' rootstock to verify their ability to feed aggressively on this resistant rootstock. These two strains were named 'HarmA' and 'HarmC'. Later investigations determined that 'HarmA' was a strain of *M. arenaria* while 'HarmC' was a strain of *M. incognita*. The dagger nematodes were collected from several vineyards in the Napa Valley of California, which were known to have fanleaf degeneration.

The 33 selections were then inoculated with each of the three nematode strains independently. The root-knot nematode screens used 1,000 J2 larvae to inoculate plants growing in 1,000 cm³ plastic pots with a coarse sand/clay loam soil mix. The dagger nematode screens were done in the same pots and soils, but used 200 adult *X. index* as the inoculum. Fourteen selections did not produce egg masses when inoculated with the three root-knot nematode strains nor did they produce root tip galls after inoculation with *X. index* (Table 2).

In the fourth phase, the 14 selections were subjected to a series of tests. The first was a combined inoculum of all four nematodes at once to determine the impact of simultaneous nematode feeding on resistance (Table 3). They were also

tested against the four nematodes over a range of temperatures, 24, 27, 30 and 32° C. There was some erosion of resistance to 'HarmA' (the most aggressive strain of root-knot nematode) at this temperature, but 6 of the selections performed very well (Table 4). Root-knot nematode resistance is known to fail at about 28° C. a wide range of species including tomato, pepper and plum. The 14 selections were also evaluated for resistance to lesion (*Pratylenchus vulnus*), citrus (*Tylenchulus semipenetrans*) and ring (*Mesocriconema xenoplax*) nematodes (Table 3). From this series of tests a number of six rootstock selections were made, including '9363 16'. A summary of the characteristics of variety '9363-16' is presented below.

Grape phylloxera are capable of feeding and producing galls (nodosities) on the young roots of virtually all grape rootstocks and species. In order to gauge the phylloxera hosting ability of the final six selections, they were tested for the ability to support phylloxera on nodosities and compared to a set of commercial rootstocks. Three sets of young root pieces from each selection were inoculated with 10 phylloxera eggs collected from '101-14 Mgt' rootstock roots. Over a 21-day period, the number of eggs and juveniles that were produced were summed and divided by 10 (the original inoculum) to produce the average rate of increase. Table 5 presents these results for the six selections. Nodosity galling on young roots does not appear to damage grapevines, only feeding and galling on mature roots (tuberosities) leads to vine death. The high rate of phylloxera feeding, galling and reproduction on '101-14 Mgt' (7.98 average rate of increase) has been observed in past tests. Most of the values were very low, although this '101-14 Mg' strain is well adapted to '9407-14'. Three of the selections were very resistant: '8909-05', '9363-16' and '9449-27'. The low values for 'AXR#1' demonstrate that the results of this test do not reflect field level or tuberosity level feeding and damage, and that phylloxera adapt independently to rootstock hosts.

It will take years to determine which sites each of these rootstock selections are best suited to, but they have unparalleled levels of resistance to nematodes and should excel in sites with single and mixed nematode species infestations. Four of the selections ('9365-43', '9365-85', '9407-14' and '9449-27') were grafted to 'Fiesta Seedless' and planted in a Fresno rootstock trial in 2004. That year two of these selections ('9365-43' and '9365-85') were also included in a Chardonnay rootstock trial in Santa Maria. The six rootstock selections in large pots using soils from the Gallo Livingston Ranch where nematode pressure from root-knot, lesion, ring and *Xiphinema americanum*, is known to be severe and chronic. This test was conducted to evaluate these selections under "field conditions" using infested soil without added inoculations. All of the selections performed very well against root-knot nematodes (Table 6) and two, '9407-14' and '8909-05', also performed very well against ring nematode (FIG. 1).

DETAILED BOTANICAL DESCRIPTION

Rootstock '9363-16' acquires its nematode resistance from *V. rufotomentosa* (highly resistant to *X. index*) and *V. champinii* 'Dog Ridge' (strong resistance to root-knot and dagger nematodes) and roots and grafts easily because of its *V. riparia* parentage. The rootstock has slightly lobed leaves, acquired from *V. rufolomentos*, that are relatively glabrous, acquired from *V. riparia*. '9363-16' is a good mothervine with staminate flowers, long shoots, long internodes and few laterals. Preliminary propagation tests have found it to

have a relatively shallow rooting angle. The rootstock is susceptible to citrus and ring nematodes, but has excellent resistance to root-knot and dagger nematodes (Table 3).

The following is a further description of rootstock ‘9363-16’. All color descriptions are from the Munsell Color Chart for Plant Tissues with Hue Value/Chroma values. Measurements are averages from 5 mid cane leaves and presented as length of the main vein from the petiole to the end of the apical lobexthe width from lateral lobe to lateral lobe. Grape flowers are inconspicuous and never used to distinguish varieties or species except in the case of being staminate, pistillate or hermaphroditic. The size of the young leaves is strongly influenced by the environment and was not reported.

Shoot tips.—The variety is slightly exposed, light green (5GY 6/8) and sparsely covered with arachnose tomentum.

Young leaves.—The variety has young leaves which are light green (5GY 6/8) mostly distinctly three-lobed with sparse arachnose tomentum and relatively sharp pointed teeth.

Mature leaves.—The mature leaves are green upper (5Y 4/6) and lower (5GY 5/6) surfaces) medium sized (8.7×10.7 cm) some with distinct three-lobing and others three-lobed to cuneiform, with short angular to convex teeth. The upper surface is glabrous, slightly rugose and mostly flat. The lower surface is mostly glabrous with sparse bristles and arachnose tomentum along the main veins, with limited tufting of bristles (domatia) at the intersection of the main veins. The petiolar sinus is a narrow, to pinched closed, U-shape.

Cane.—The variety has light brown (5YR 4/6) canes with long internodes and relatively few lateral shoots, nodes have medium sized slightly pointed buds. Canes are round in cross-section and have very thin diaphragms and medium pith.

Flowers.—The flowers are staminate.

TABLE 1

Parentage of the five nematode resistant grape rootstock selections.	
Selection	Parentage
‘8909-05’	<i>V. rupestris</i> × <i>M. rotundifolia</i>
‘9363-16’	(<i>V. rufotomentosa</i> × (<i>V. champinii</i> ‘Dog Ridge’ × ‘Riparia Gloire’)) × ‘Riparia Gloire’
‘9365-43’	(<i>V. rufotomentosa</i> × (<i>V. champinii</i> ‘Dog Ridge’ × ‘Riparia Gloire’)) × <i>V. champinii</i> ‘c9038’ (probably <i>V. candicans</i> × <i>V. monticola</i>)
‘9365-85’	(<i>V. rufotomentosa</i> × (<i>V. champinii</i> ‘Dog Ridge’ × ‘Riparia Gloire’)) × <i>V. champinii</i> ‘c9038’ (probably <i>V. candicans</i> × <i>V. monticola</i>)
‘9407-14’	(<i>V. champinii</i> ‘Ramsey’ × ‘Riparia Gloire’ × <i>V. champinii</i> ‘c9021’ (probably <i>V. candicans</i> × <i>V. berlandieri</i>))

TABLE 2

Selections with broad resistance to four nematodes when inoculated individually.				
Selection	X. index	<i>M. incognita</i> I3	<i>M. arenaria-HarmA</i>	<i>M. incognita-HarmC</i>
‘8909-05’	R	R	R	R
‘9317-06’	R	R	R	R

TABLE 2-continued

Selections with broad resistance to four nematodes when inoculated individually.				
Selection	X. index	<i>M. incognita</i> I3	<i>M. arenaria-HarmA</i>	<i>M. incognita-HarmC</i>
‘9332-43’	R	R	R	R
‘9344-03’	R	R	R	R
‘9363-16’	R	R	R	R
‘9365-43’	R	R	R	R-
‘9365-62’	R	R	R	R
‘9365-85’	R-	R	R	R
‘9403-35’	R	R	R-	R-
‘9403-107’	R	R	R	R
‘9407-14’	R	R	R	R
‘9449-23’	R	R	R	R
‘9449-25’	R	R	R	R
‘9449-27’	R	R	R	R
Control Group:				
‘1616C’	S	R	R-	R-
‘Harmony’	S	R	S	S
‘Colombard’	S	S	S	S

R = Resistant, no gall symptoms or egg masses observed
R- = Trace infection
S = Susceptible, symptoms present, nematode reproduction supported

TABLE 3

Summary results for 14 nematode resistant selections when tested against combined inoculum with three root-knot nematode (RKN) strains and dagger nematode (Xi). Results of testing against citrus, lesion and ring are also reported.						
Genotypes	Xi Galls in Combined Testing		RKN Egg Masses in Combined Testing		Citrus	
‘8909-05’	R	0	R	0	R	<100
‘9317-06’	MS	<1	S	<5	R	<100
‘9332-43’	S	<5	S	<5	R	<100
‘9344-03’	S	<5	MS	<1	S	>400
‘9363-16’	R	0	R	0	S	>400
‘9365-43’	R	0	R-	<1	R	<100
‘9365-62’	MS	<1	S	<5	R	<100
‘9365-85’	MS	<1	R-	<1	R	<100
‘9403-107’	R	0	S	<5	R	<100
‘9403-35’	S	<5	S	<5		
‘9407-14’	R	0	R	0	R	<100
‘9449-23’	MS	<1	R	0	R	<100
‘9449-25’	MS	<1	R	0	R	<100
‘9449-27’	MS	<1	R-	<1	R	<100
‘1616C’	S	>50	S	<5	S	>400
‘Freedom’	S	>10	S	<10	R	<100
‘Colombard’	S	>100	S	>100	S	<1000
‘Harmony’	S	>10	S	<50	S	<1000
‘St. George’	S	<100	S	<50	S	>400

Genotypes	Lesion		Ring	
‘8909-05’	R	<10	R	<1,000
‘9317-06’	R	<50	HS	<20,000
‘9332-43’	R	<50	S	<10,000
‘9344-03’	S	<200	S	<10,000
‘9363-16’	R	<50	S	<10,000
‘9365-43’	R	<50	MS	<5,000
‘9365-62’	R	<50	MS	<5,000
‘9365-85’	R	<50	S	<10,000
‘9403-107’	R	<50	MS	<5,000
‘9403-35’	R	<50	MS	<5,000
‘9407-14’	R	<50	LS	<3,000
‘9449-23’	R	<50	S	<10,000
‘9449-25’	R	<50	S	<10,000
‘9449-27’	R	<50	S	<10,000
‘1616C’	S	<200	HS	<20,000
‘Freedom’	R	<50	HS	<20,000

TABLE 3-continued

Summary results for 14 nematode resistant selections when tested against combined inoculum with three root-knot nematode (RKN) strains and dagger nematode (Xi). Results of testing against citrus, lesion and ring are also reported.					
	'Colombard'	S	<300	HS	<30,000
	'Harmony'	S	<200	HS	<30,000
	'St. George'	R	<50	S	<10,000

TABLE 4

Number of egg masses per potted plant and per gram of root after inoculation with 1,000 <i>M. arenaria</i> 'HarmA' nematodes and testing at 32 C.			
Genotype	Egg Mass/Plant	Egg Mass/g Root	
'Colombard'	442.50 a	155.86	a
'Harmony'	156.00 b	123.28	b
'9365-85'	32.25 c	16.49	c
'9365-43'	7.25 c	6.04	cd
'9363-16'	6.75 c	5.74	cd
'9449-27'	0.25 c	0.22	d
'9317-06'	0.00 c	0	d
'8909-05'	0.00 c	0	d

TABLE 5

The average rate of population increase of phylloxera on nodosities formed on young root tips of advanced rootstock selections and a set of standard rootstocks. The phylloxera were selected from a vineyard planted on '101-14 Mgt' rootstock	
'8909-05'	0.59
'9363-16'	0.89
'9365-43'	1.86

TABLE 5-continued

The average rate of population increase of phylloxera on nodosities formed on young root tips of advanced rootstock selections and a set of standard rootstocks. The phylloxera were selected from a vineyard planted on '101-14 Mgt' rootstock	
'9365-85'	1.99
'9407-14'	9.83
'9449-27'	0.30
'1103P'	2.10
'101-14 Mgt'	7.98
'Teleki 5C'	1.38
'AxR#1'	2.15
'O39-16'	0.43

TABLE 6

Number of root-knot nematode egg masses recovered from rootstock selections growing in 1 gal pots with soils collected from sites at the Gallo Livingston vineyards. This vineyard is known to have severe and chronic nematode pressure. Results per soils are means of three replicate pots.			
Selection	Soil 1	Soil 2	Soil 3
'Colombard'	69.02	89.8	74.6
'Harmony'	31.2	0	2.8
'St. George'	9.8	26.2	16.0
'9365-85'	1	0	1.4
'9407-14'	0	0	0
'9363-16'	0	0	1.4
'9365-43'	0	0	0.2
'9449-27'	0	0	0.8
'8909-05'	0	0	0

What is claimed is:

1. A novel and distinct variety of grape rootstock designated '9363-16' having the characteristics described and illustrated herein.

* * * * *




Figure 1




Figure 2