



(12) **United States Plant Patent**
Wood

(10) **Patent No.:** **US PP20,025 P2**
(45) **Date of Patent:** **May 26, 2009**

(54) **WEIGELA PLANT NAMED ‘CARLTON’**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(50) Latin Name: ***Weigela florida***
Varietal Denomination: **Carlton**

(52) **U.S. Cl.** **Plt./226**
(58) **Field of Classification Search** **Plt./226**
See application file for complete search history.

(75) Inventor: **Timothy D. Wood**, Spring Lake, MI
(US)

Primary Examiner—S. B McCormick Ewoldt
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(73) Assignee: **Spring Meadow Nursery Inc.**, Grand
Haven, MI (US)

(57) **ABSTRACT**

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

A new and distinct cultivar of *Weigela* plant named
‘Carlton’, characterized by its uniform plant habit; freely
branching habit; developing leaves that are yellow green in
color and fully expanded leaves that become yellow green in
color during the mid-summer through autumn; dark red
purple-colored flowers; and good garden performance.

(21) Appl. No.: **12/075,453**

(22) Filed: **Mar. 11, 2008**

1 Drawing Sheet

1

Botanical designation: *Weigela florida*.
Cultivar denomination: ‘Carlton’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Weigela* plant, botanically known as *Weigela florida*, and
hereinafter referred to by the name ‘Carlton’.

The new *Weigela* is a product of a planned breeding pro-
gram conducted by the Inventor in Grand Haven, Mich. The
objective of the breeding program was to develop new
Weigelas with unique leaf shapes and coloration, uniform
plant habit and hardiness.

The new *Weigela* originated from an open-pollination in
2003 of the *Weigela florida* cultivar Rubidor, not patented.
The cultivar Carlton was discovered and selected by the
Inventor in 2004 as a flowering plant within the progeny of
the stated open-pollination in a controlled environment in
Grand Haven, Mich. Asexual reproduction of the new culti-
var by cuttings in Grand Haven, Mich. since the summer of
2004 has shown that the unique features of this new *Weigela*
plant are stable and reproduced true to type in successive
generations of asexual reproduction.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Carlton’.
These characteristics in combination distinguish ‘Carlton’ as
a new and distinct cultivar of *Weigela*:

1. Uniform plant habit.
2. Freely branching habit.
3. Developing leaves that are yellow green in color and
fully expanded leaves that become yellow green in
color during the mid-summer through autumn.
4. Dark red purple-colored flowers.
5. Good garden performance.

In side-by-side comparisons in Grand Haven, Mich.,
under commercial practice, plants of the new *Weigela* differ

2

from plants of the female parent, the cultivar Rubidor, in the
following characteristics:

1. Plants of the new *Weigela* and the cultivar Rubidor
differ in leaf color as plants of the cultivar Rubidor have
developing leaves that are yellow in color.
2. Under high light conditions, leaves of plants of the new
Weigela do not burn whereas leaves of plants of the
cultivar Rubidor burn.

Plants of the new *Weigela* can be compared to plants of
the *Weigela florida* cultivar Variegata, not patented. In side-
by-side comparisons in Grand Haven, Mich. plants of the
new *Weigela* differ from plants of the cultivar Variegata in
the following characteristics:

1. Plants of the new *Weigela* and the cultivar Variegata
differ in leaf coloration as plants of the cultivar Varie-
gata have variegated leaves.
2. Plants of the new *Weigela* and the cultivar Variegata
differ in flower color as plants of the cultivar Variegata
have light pink and white-colored flowers.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the
overall appearance of the new *Weigela*, showing the colors
as true as it is reasonably possible to obtain in colored repro-
ductions of this type. Colors in the photographs may differ
slightly from the color values cited in the detailed botanical
description which accurately describe the colors of the new
Weigela.

The photograph at the bottom of the sheet comprises a
side perspective view of a typical plant of the cultivar Carl-
ton.

The photograph at the top of the sheet is a close-up view
of typical flowers and leaves of the cultivar Carlton.

DETAILED BOTANICAL DESCRIPTION

The new *Weigela* has not been observed under all possible
environmental conditions. The phenotype may vary some-
what with variations in environment such as temperature and

light intensity, without, however, any variance in genotype. The aforementioned photographs, following observations and measurements describe plants grown during the summer and autumn in Grand Haven, Mich., in an outdoor nursery and under commercial production practices. Plants were about three years old when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Weigela florida* cultivar Carlton.

Parentage:

Female or seed, parent.—*Weigela florida* cultivar Rubidor, not patented.

Male or pollen, parent.—Unknown selection of *Weigela florida*, not patented.

Propagation:

Type.—By softwood cuttings.

Time to initiate roots.—About 15 days at temperatures of 22° C.

Time to produce a rooted young plant.—About two months at temperatures of 22° C.

Root description.—Fine.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Perennial shrub. Upright, rounded and uniform plant habit. Vigorous growth habit. Freely branching habit with about 20 lateral branches developing per plant; pinching enhances lateral branch development.

Plant height.—About 2 meters.

Plant diameter.—About 1.7 meters.

Lateral branch description:

Length.—About 1.9 meters.

Diameter.—About 4 mm.

Internode length.—About 9 cm.

Strength.—Strong.

Aspect.—Erect to about 45° from vertical.

Texture.—Pubescent.

Color, developing stems.—145B.

Color, mature stems.—197B.

Foliage description:

Arrangement.—Opposite; simple.

Length.—About 10 cm.

Width.—About 5 cm.

Shape.—Elliptic.

Apex.—Acute to acuminate.

Base.—Cuneate.

Margin.—Serrulate; slightly undulating.

Texture, upper surface.—Smooth, glabrous.

Texture, lower surface.—Pubescence at veins.

Venation pattern.—Pinnate.

Color.—Developing foliage, upper surface: 151C. Developing foliage, lower surface: 154C. Fully expanded foliage, upper surface: During the spring, 144A; mid-summer to autumn, color becoming closer to 151C; venation, 151B. Fully expanded foliage, lower surface: During the spring, 138B; mid-summer to autumn, color becoming closer to 151C; venation, 151B.

Petiole.—Length: About 6 mm. Diameter: About 2 mm.

Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: 151B.

Flower description:

Flower arrangement and habit.—Funnelform flowers arranged in clusters of about nine flowers. Flowers not persistent. Flowers face upright, outwardly or are drooping.

Fragrance.—None detected.

Natural flowering season.—Plants of the new *Weigela* flower from May to June and will typically reflower in October in Grand Haven, Mich.

Flower longevity.—Flowers last about six to eight weeks on the plant.

Flower diameter.—About 3 cm.

Flower length (height).—About 4 cm.

Flower bud.—Shape: Narrowly ovate. Length: About 1.2 cm. Diameter: About 4 mm. Color: 60A.

Corolla.—Arrangement: Funnelform, five petals fused into a long tube with separate petal lobes. Petal lobe length: About 1.3 cm. Petal lobe width: About 9 mm. Petal lobe shape: Roughly orbicular. Petal lobe apex: Obtuse. Petal lobe margin: Mostly entire. Petal texture, upper and lower surfaces: Smooth, glabrous. Color: Petal, when opening, upper and lower surfaces: 60A. Petal, fully opened, upper and lower surfaces: 60B; color becoming closer to 60C with development.

Sepals.—Quantity per flower: Five. Length: About 1 cm. Width: About 1 mm. Shape: Lanceolate. Apex: Acuminate. Margin: Entire. Sepal texture, upper and lower surfaces: Smooth; glabrous. Color, upper and lower surfaces: 155B; towards the apex, speckled with 60C.

Peduncles.—Length: About 8 cm. Diameter: About 2 mm. Angle: Erect to about 30° from stem axis. Strength: Moderately strong; flexible. Texture: Smooth, glabrous. Color: 138B.

Pedicels.—Length: About 1.8 cm. Diameter: About 1.5 mm. Angle: About 10° to about 60° from stem axis. Strength: Moderately strong; flexible. Texture: Smooth, glabrous. Color: 138B.

Reproductive organs.—Stamens: Quantity/arrangement: Five per flower. Anther shape: Oblong. Anther length: About 4 mm. Anther color: 196C. Pollen amount: Moderate. Pollen color: Close to 156A. Pistils: Quantity: One per flower. Pistil length: About 3.5 cm. Style length: About 2 cm. Style color: 196A. Stigma shape: Rounded to globular. Stigma color: 196C. Seeds: Length: Less than 0.5 mm. Diameter: Less than 0.5 mm. Color: Close to 200A.

Garden performance: Plants of the new *Weigela* have exhibited good tolerance to rain and wind and have been observed to tolerate temperatures from about -29° C. to about 38° C.

Pathogen/pest resistance: Plants of the new *Weigela* have not been observed to be resistant to pests and pathogens common to *Weigela*.

It is claimed:

1. A new and distinct *Weigela* plant named 'Carlton' as illustrated and described.

* * * * *

