



(12) **United States Plant Patent**
Goetz

(10) **Patent No.:** **US PP20,084 P2**
(45) **Date of Patent:** **Jun. 9, 2009**

(54) **FUCHSIA PLANT NAMED ‘GOETZIMPRO’**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(50) Latin Name: ***Fuchsia×hybrida***
Varietal Denomination: **Goetzimpro**

(52) **U.S. Cl.** **Plt./300**
(58) **Field of Classification Search** **Plt./300**
See application file for complete search history.

(75) Inventor: **Wolfram Goetz**, Herbrechtingen (DE)

Primary Examiner—June Hwu
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(73) Assignee: **Amerinova Properties LLC**, Bonsall,
CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(57) **ABSTRACT**

A new and distinct cultivar of *Fuchsia* plant named
‘Goetzimpro’, characterized by its compact, upright to out-
wardly arching growth habit; freely branching plant habit;
light and darker pink-colored flowers; early and freely flow-
ering habit; and good garden performance.

(21) Appl. No.: **12/070,571**

(22) Filed: **Feb. 19, 2008**

1 Drawing Sheet

1

Botanical designation: *Fuchsia×hybrida*.
Cultivar denomination: ‘GOETZIMPRO’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Fuchsia*, botanically known as *Fuchsia×hybrida* and
hereinafter referred to by the name ‘Goetzimpro’.

The new *Fuchsia* is a product of a planned breeding pro-
gram conducted by the Inventor in Herbrechtingen, Ger-
many. The objective of the breeding program is to create new
Fuchsia cultivars that flower early and have good garden
performance.

The new *Fuchsia* originated from a cross-pollination
made by the Inventor in 2000 in Herbrechtingen, Germany
of a proprietary selection of *Fuchsia×hybrida* identified as
code number 58/97, not patented, as the female, or seed,
parent with the *Fuchsia×hybrida* cultivar Goetzrose, dis-
closed in U.S. Plant Pat. No. 14,492, as the male, or pollen,
parent. The new *Fuchsia* was discovered and selected by the
Inventor as a single flowering plant within the progeny of the
stated cross-pollination in a controlled environment in
Herbrechtingen, Germany in 2001.

Asexual reproduction of the new *Fuchsia* by terminal cut-
tings in a controlled environment in Herbrechtingen, Ger-
many since the autumn of 2001, has shown that the unique
features of this new *Fuchsia* are stable and reproduced true
to type in successive generations.

SUMMARY OF THE INVENTION

The cultivar Goetzimpro has not been observed under all
possible environmental conditions. The phenotype may vary
somewhat with variations in environment and cultural prac-
tices such as temperature, daylength and light intensity
without, however, any variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of ‘Goetzim-
pro’. These characteristics in combination distinguish ‘Goet-
zimpro’ as a new and distinct cultivar of *Fuchsia*:

1. Compact, upright to outwardly arching growth habit.
2. Freely branching plant habit.

2

3. Light and darker pink-colored flowers.
4. Early and freely flowering habit.
5. Good garden performance.

Plants of the new *Fuchsia* differ primarily from plants of
the female patent selection in the following characteristics:

1. Plants of the new *Fuchsia* have smaller leaves than
plants of the female parent selection.
2. Plants of the new *Fuchsia* flower earlier than plants of
the female parent selection.

Plants of the new *Fuchsia* differ primarily from plants of
the male patent, the cultivar Goetzrose, in the following
characteristics:

1. Plants of the new *Fuchsia* are more compact than plants
of the cultivar Goetzrose.
2. Plants of the new *Fuchsia* flower earlier than plants of
the cultivar Goetzrose.
3. Plants of the new *Fuchsia* are more weather-tolerant
than plants of the cultivar Goetzrose.

Plants of the new *Fuchsia* can be compared to plants of
the cultivar Kiecandiro, disclosed in U.S. Plant Pat. No.
15,410. Plants of the new *Fuchsia* and the cultivar Kiecand-
iro differ in the following characteristics:

1. Plants of the new *Fuchsia* are more compact than plants
of the cultivar Kiecandiro.
2. Plants of the new *Fuchsia* are not as upright as plants of
the cultivar Kiecandiro.
3. Plants of the new *Fuchsia* and the cultivar Kiecandiro
differ in flower color.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the
overall appearance of the new *Fuchsia*, showing the colors
as true as it is reasonably possible to obtain in colored repro-
ductions of this type. Colors in the photographs may differ
slightly from the color values cited in the detailed botanical
description which accurately describe the colors of the new
Fuchsia.

The photograph at the bottom of the sheet comprises a
side perspective view of a typical flowering plant of ‘Goetz-
impro’ grown in a container.

The photograph at the top of the sheet is a close-up of a typical flowering stem of 'Goetzimpro'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown in Bonsall, Calif., under commercial practice during the spring and early summer in a polyethylene-covered greenhouse with day temperatures ranging from 16° C. to 35° C. and night temperatures ranging from 13° C. to 21° C. Plants were grown for about 18 weeks in 15-cm containers and were pinched. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Fuchsia*×*hybrida* cultivar Goetzimpro.

Parentage:

Female, or seed, parent.—Proprietary selection of *Fuchsia*×*hybrida* identified as code number 58/97, not patented.

Male, or pollen, parent.—*Fuchsia*×*hybrida* cultivar Goetzrose, disclosed in U.S. Plant Pat. No. 14,492.

Propagation:

Type.—By terminal cuttings.

Time to initiate roots, summer.—About two to three weeks.

Time to initiate roots, winter.—About three weeks.

Time to produce a rooted young plant, summer.—About six weeks.

Time to produce a rooted young plant, winter.—About eight weeks.

Root description.—Fine, fleshy; white to light brown in color.

Rooting habit.—Freely branching; moderately dense.

Plant description:

Plant and growth habit.—Initially upright, then outwardly spreading; compact plant habit; rounded in overall shape. Freely branching; about six to seven primary branches per plant. Moderately vigorous growth habit.

Plant height.—About 24 cm.

Plant diameter.—About 22 cm.

Lateral branch description:

Length.—About 16 cm.

Diameter.—About 4 mm.

Internode length.—About 1.3 cm.

Strength.—Strong.

Aspect.—Initially upright to outwardly arching.

Texture.—Pubescent; minute hairs.

Color, young.—146C.

Color, mature.—177A to 177B.

Foliage description:

Arrangement.—Opposite, simple.

Length.—About 4.3 cm.

Width.—About 2.3 cm.

Shape.—Ovate to elliptic.

Apex.—Acute.

Base.—Obtuse to slightly attenuate.

Margin.—Entire with irregular shallow points.

Texture, upper and lower surfaces.—Smooth, glabrous.

Venation pattern.—Pinnate; arcuate.

Color.—Developing foliage, upper surface: 146A.

Developing foliage, lower surface: 146B. Fully expanded foliage, upper surface: 147A; venation,

146A. Fully expanded foliage, lower surface: 146A; venation, 182B.

Petiole.—Length: About 1.7 cm. Diameter: About 2 mm. Texture, upper and lower surfaces: Pubescent; minute hairs. Color, upper surface: 199C. Color, lower surface: 174C.

Flower description:

Flower arrangement and habit.—Single bi-colored axillary flowers. Flowers initially upright and then pendulous. Flowers not fragrant. Freely flowering habit with typically about 70 open flowers and flower buds per plant.

Natural flowering season.—In southern California, plants flower from early spring to fall; flowering continuous during this period. Plants begin flowering about four to six weeks after planting. Flowers last about five to six days on the plant. Flowers not persistent.

Flower diameter.—About 1.7 cm; corolla diameter, about 1 cm.

Flower height.—About 5 cm.

Flower buds.—Shape: Elliptic. Length: About 3.2 cm. Diameter: About 1.1 cm. Color: 157C tinted with 56B.

Petals.—Arrangement: Four in a single whorl. Length: About 1.4 cm. Width: About 9 mm. Shape: Obovate. Apex: Rounded. Base: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening, upper and lower surfaces: 85D. Fully opened, upper surface: 155A. Fully opened, lower surface: 155C.

Sepals.—Arrangement: Calyx star-shaped with four sepals fused at the base. Length: About 2.1 cm. Width: About 7 mm. Shape: Narrowly elliptic. Apex: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening and fully opened, upper surface: 158C tinted with 56B to 56C. When opening and fully opened, lower surface: 55B to 55C.

Peduncles.—Length: About 2.8 cm. Diameter: About 1 mm. Angle: About 45° to 60° from vertical. Strength: Strong. Texture: Pubescent; minute hairs. Color: 145A.

Reproductive organs.—Stamens: Quantity: Eight per flower. Filament length: About 2 cm. Filament color: 158C. Anther shape: Oval. Anther size: About 1 mm by 2 mm. Anther color: 177A. Pollen amount: Sparse. Pollen color: 164B. Pistils: Quantity: One per flower. Pistil length: About 5 cm. Style length: About 4.2 cm. Style color: 56D. Stigma shape: Oval. Stigma color: 160B. Ovary color: 146C.

Seed/fruit.—Seed and fruit development have not been observed on plants of the new *Fuchsia*.

Garden performance. —Plants of the new *Fuchsia* have been observed to have good garden performance and to tolerate wind, rain and temperatures from about 10° C. to about 32° C.

Pathogen/pest resistance: Plants of the new *Fuchsia* have not been observed to be resistant to pests and pathogens common to *Fuchsia*.

It is claimed:

1. A new and distinct *Fuchsia* plant named 'Goetzimpro' as illustrated and described.

