



US00PP20095P2

(12) **United States Plant Patent**  
**Schröder**

(10) **Patent No.:** **US PP20,095 P2**  
(45) **Date of Patent:** **Jun. 16, 2009**

(54) ***OSTEOSPERMUM* PLANT NAMED ‘SUMIPAS 03’**

(50) Latin Name: ***Osteospermum ecklonis***  
Varietal Denomination: **Sumipas 03**

(76) Inventor: **Ralf Schröder**, Karl-Leisner-Str. 15,  
D-59348 Lüdinghausen (DE)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 454 days.

(21) Appl. No.: **11/323,155**

(22) Filed: **Dec. 30, 2005**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(52) **U.S. Cl.** ..... **Plt./360**

(58) **Field of Classification Search** ..... **Plt./360**  
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

Upov CD–Rom Plant Variety Database 2006/04 search for  
cultivar Sumipas 03.\*

\* cited by examiner

*Primary Examiner*—Kent Bell  
*Assistant Examiner*—Annette H Para  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Osteospermum* plant named  
‘Sumipas 03’, characterized by its upright, outwardly  
spreading, compact and mounded plant habit; freely branch-  
ing habit; freely, early and continuous flowering habit; and  
inflorescences with purple-colored ray florets.

**1 Drawing Sheet**

**1**

Botanical designation: *Osteospermum ecklonis*.  
Botanical denomination: ‘Sumipas 03’.

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct cultivar  
of *Osteospermum* plant, botanically known as *Osteosper-  
mum ecklonis*, and hereinafter referred to by the name  
‘Sumipas 03’.

The new *Osteospermum* is a product of a planned breed-  
ing program conducted by the Inventor in Lüdinghausen,  
Germany. The objective of the breeding program was to cre-  
ate new compact and early flowering *Osteospermum* culti-  
vars with large and attractive inflorescences.

The new *Osteospermum* originated from a cross-  
pollination in April, 2001 of a proprietary *Osteospermum*  
*ecklonis* selection identified as code number 24, not  
patented, as the female, or seed, parent with the *Osteosper-  
mum ecklonis* cultivar Sumpink, not patented, as the male, or  
pollen, parent. The new *Osteospermum* was discovered and  
selected by the Inventor as a single flowering plant within  
the progeny of the stated cross-pollination grown in a con-  
trolled environment in Lüdinghausen, Germany in April,  
2002.

Asexual reproduction of the new *Osteospermum* by termi-  
nal vegetative cuttings was first conducted in Lüdinghausen,  
Germany in May, 2002. Asexual reproduction by cuttings  
has shown that the unique features of this new *Osteosper-  
mum* are stable and reproduced true to type in successive  
generations.

**SUMMARY OF THE INVENTION**

The cultivar Sumipas 03 has not been observed under all  
possible environmental conditions. The phenotype may vary  
somewhat with variations in environment such as tempera-

**2**

ture and light intensity without, however, any variance in  
genotype.

The following traits have been repeatedly observed and  
are determined to be the unique characteristics of ‘Sumipas  
03’. These characteristics in combination distinguish ‘Sumi-  
pas 03’ as a new and distinct *Osteospermum*:

1. Upright, outwardly spreading, compact and mounded  
plant habit.
2. Freely branching habit.
3. Freely, early and continuous flowering habit.
4. Inflorescences with purple-colored ray florets.

Plants of the new *Osteospermum* differ primarily from  
plants of the female parent selection in flowering time as  
plants of the new *Osteospermum* flower about 10 days ear-  
lier than plants of the female parent selection. Plants of the  
new *Osteospermum* differ primarily from plants of the male  
parent, the cultivar Sumpink in plant size as plants of the  
new *Osteospermum* are more compact than plants of the  
cultivar Sumpink.

Plants of the new *Osteospermum* can be compared to  
plants of the cultivar Osoutis, disclosed in U.S. Plant Pat.  
No. 15,491. In side-by-side comparisons, plants of the new  
*Osteospermum* differed from plants of the cultivar Osoutis in  
the following characteristics:

1. Plants of the new *Osteospermum* were larger than  
plants of the cultivar Osoutis.
2. Plants of the new *Osteospermum* had thicker stems than  
plants of the cultivar Osoutis.
3. Plants of the new *Osteospermum* had larger inflores-  
cences than plants of the cultivar Osoutis.
4. Plants of the new *Osteospermum* and the cultivar Osou-  
tis differed in ray floret coloration.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying colored photograph illustrates the  
overall appearance of the new *Osteospermum* showing the



colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ from the color values cited in the detailed botanical description which accurately describe the colors of the new *Osteospermum*. The photograph comprises a top perspective view of a typical flowering plant of 'Sumipas 03' grown in a container.

#### DETAILED BOTANICAL DESCRIPTION

The new *Osteospermum* has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype.

The aforementioned photograph, following observations and measurements describe plants grown in Lüdinghausen, Germany in an outdoor nursery during the spring and summer and under commercial production practices. Plants were about three months from planting rooted young plants when the photograph and description were taken. During the production of the plants, day temperatures ranged from 16° C. to 30° C. and night temperatures ranged from 8° C. to 14° C. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Osteospermum ecklonis* cultivar Sumipas 03.

Parentage:

*Female, or seed, parent.*—Proprietary selection of *Osteospermum ecklonis* identified as code number 24, not patented.

*Male, or pollen, parent.*—*Osteospermum ecklonis* cultivar Sumpink, not patented.

Propagation:

*Type.*—Terminal vegetative cuttings.

*Time to initiate rooting, summer.*—About 14 days at 16° C.

*Time to initiate rooting, winter.*—About 18 days at 16° C.

*Time to produce a rooted young plant, summer.*—About 30 days at 16° C.

*Time to produce a rooted young plant, winter.*—About 32 days at 16° C.

*Root description.*—Fine, fibrous; color, 162C.

*Rooting habit.*—Freely branching; moderately dense.

Plant description:

*Plant form and growth habit.*—Perennial herbaceous container and garden plant. Upright, somewhat outwardly spreading, compact and mounded plant habit; broad inverted triangle. Freely branching habit, about 18 primary lateral branches and numerous secondary lateral branches.

*Plant height.*—About 25 cm.

*Plant width or area of spread.*—About 20 cm.

*Lateral branches.*—Length: About 16 cm. Diameter: About 5 mm. Internode length: About 8 cm to 10 cm. Aspect: Upright and outwardly spreading. Strength: Strong. Texture: Smooth, glabrous. Color: 145B.

*Foliage description.*—Arrangement: Alternate; simple. Length: About 4 cm to 5.5 cm. Width: About 2.5 cm to 3.5 cm. Shape: Lanceolate. Apex: Emarginate. Base: Acute to truncate. Margin: Slightly serrate. Venation pattern: Pinnate. Texture, upper and lower surfaces: Smooth, glabrous. Color: Developing

foliage, upper and lower surfaces: 137A. Fully expanded foliage, upper surface: 146A; venation, 146B. Fully expanded foliage, lower surface: 146B; venation, 146B. Petiole: Length: About 1 cm to 1.5 cm. Diameter: About 2 mm to 3 mm. Texture, upper and lower surfaces: Glabrous, smooth. Color, upper and lower surfaces: 146D.

Inflorescence description:

*Appearance.*—Solitary terminal and axillary inflorescences held above and beyond the foliage on strong peduncles. Composite inflorescence form, radially symmetrical, with elliptic to ligulate-shaped ray florets and disc florets massed at the center; ray and disc florets develop acropetally on a capitulum. Inflorescences not persistent. Inflorescences face mostly upright to outwardly.

*Flowering response.*—Plants flower continuous and freely from the early spring through the fall in Germany. Early flowering habit; plants grown in containers begin flowering about six to eight weeks after pinching.

*Postproduction longevity.*—Inflorescences maintain good color and substance for about four to six days on the plant.

*Quantity of inflorescences.*—Freely flowering; at one time, about eight to twelve open inflorescences and inflorescence buds per lateral stem.

*Fragrance.*—Faint.

*Inflorescence bud.*—Length: About 1 cm to 1.5 cm. Diameter: About 7 mm to 10 mm. Shape: Ovoid. Color: 145C. Inflorescence size: Diameter: About 5.5 cm to 6 cm. Depth (height): About 5 cm. Disc diameter: About 2 cm. Ray florets: Length: About 2.5 cm to 2.8 cm. Width: About 5 mm to 6 mm. Shape: Elliptic to ligulate. Apex: Obtuse. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Number of ray florets per inflorescence: About 16 to 18 in a single whorl. Color: When opening, upper surface: 77A. When opening, lower surface: Close to N79A. Fully opened, upper surface: 77A to 77B; color becoming closer to 79C with development. Fully opened, lower surface: N79C to N79D.

*Disc florets.*—Shape: Tubular, elongated. Apex: Five-pointed. Length: About 1 cm. Width: At apex: About 5 mm. At base: About 1 mm. Number of disc florets per inflorescence: About 60. Color: 79A.

*Phyllaries.*—Length: About 1.5 mm to 2 mm. Diameter: About 1 mm. Shape: Lanceolate. Apex: Acute. Base: Truncate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Number per inflorescence: About 16 to 20. Color, upper and lower surfaces: 137C.

*Peduncles.*—Length: About 8 cm to 10 cm. Diameter: About 1 mm. Angle: Terminal peduncles, mostly upright; axillary peduncles about 35° to 45° from stem axis. Strength: Strong. Texture: Smooth, glabrous. Color: 144A.

*Reproductive organs.*—Androecium: Present on disc florets only. Stamen number: Five per floret; fused around style. Anther shape: Oblong. Anther length: Less than 0.5 mm. Anther color: 25B. Pollen amount: Scarce. Pollen color: 25B. Gynoecium: Present on both ray and disc florets. Pistil number: One per floret. Pistil length: About 4 mm. Stigma shape: Two-parted. Stigma color: N89D. Style length: About 1 mm. Style color: 144C. Ovary color: 144C.

**5**

*Seeds*.—Length: About 3 mm. Diameter: About 1.5 mm. Color: 200D.

Disease/pest resistance: Resistance to pathogens and pests common to *Osteospermums* has not been observed on plants grown under commercial greenhouse or outdoor conditions.

**6**

Temperature tolerance: Plants of the new *Osteospermum* have been observed to tolerate temperatures from 0° C. to 40° C.

It is claimed:

1. A new and distinct cultivar of *Osteospermum* plant named ‘Sumipas 03’, as illustrated and described.

\* \* \* \* \*



