

### US00PP20276P2

# (12) United States Plant Patent Hanes

(10) Patent No.:

US PP20,276 P2

(45) Date of Patent:

Sep. 8, 2009

# (54) PELARGONIUM PLANT NAMED 'AMRI WHIT09'

(50) Latin Name: *Pelargonium hortorum (L.) L' Herit. ex Ait* 

Varietal Denomination: Amri Whit09

(75) Inventor: Mitchell Hanes, Morgan Hill, CA (US)

(73) Assignee: Goldsmith Seeds, Inc., Gilroy, CA (US)

(\*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: 12/220,486

(22) Filed: **Jul. 24, 2008** 

(51) Int. Cl. A01H 5/00 (2006.01)

(52) U.S. Cl. ...... Plt./326

See application file for complete search history.

Primary Examiner—Susan B McCormick Ewoldt (74) Attorney, Agent, or Firm—S. Matthew Edwards

# (57) ABSTRACT

A new *Pelargonium* plant named 'Amri Whit09,' particularly distinguished by the clean white flower color, light to medium green foliage, round and semi-compact habit, with good branching and garden performance.

### 1 Drawing Sheet

# 1

Latin name of the genus and species of the plant claimed:  $Pelargonium\ hortorum\ (L.)\ L'Herit.\ ex\ Ait.$ 

Varietal denomination: 'Amri Whit09'.

#### BACKGROUND OF THE NEW PLANT

The present invention comprises a new *Pelargonium*, botanically known as *Pelargonium hortorum*, and hereinafter referred to by the variety name 'Amri Whit09'.

'Amri Whit09' is a product of a planned breeding program. <sup>10</sup> The new cultivar 'Amri Whit09' has a clean white flower color, light to medium green foliage, round and semi-compact habit, with good branching and garden performance.

'Amri Whit09' originated from a sibling cross in a controlled breeding program in Gilroy, Calif. USA. The female 15 parent was an unpatented hybrid seedling identified as '9912-3' with white flower color. '9912-3' has smaller flowers and is less vigorous than 'Amri Whit09'.

The male parent of 'Amri Whit09' was an unpatented hybrid seedling identified as '9912-2' with white color. 20 '9912-2' has fewer petals, less vigor, and is earlier to flower than 'Amri Whit09'.

'Amri Whit09' was selected as one flowering plant within the progeny of the stated cross in 2005 in a controlled environment in Gilroy, Calif. USA.

The first act of asexual reproduction of 'Amri Whit09' was accomplished when vegetative cuttings were taken from the initial selection in the November 2005 in a controlled environment in Gilroy, Calif. USA.

Horticultural examination of plants grown from cuttings of the plant initiated in November 2005 in Gilroy, Calif., USA and continuing thereafter, has demonstrated that the combination of characteristics as herein disclosed for 'Amri Whit09' are firmly fixed and are retained through successive generations of asexual reproduction.

'Amri Whit09' has not been observed under all possible environmental conditions. The phenotype may vary significantly with variations in environment such as temperature, light intensity, and day length.

Plant Breeder's Rights for this cultivar were applied for in Canada on Dec. 24, 2007 and in Germany on Jul. 2, 2007. 'Amri Whit09' has not been made publicly available more than one year prior to the filing of this application.

# 2

#### DESCRIPTION OF THE DRAWING

The accompanying photographic drawing shows typical flower and foliage characteristics of 'Amri Whit09' with colors being as true as possible with an illustration of this type.

The photographic drawing shows a container of three flowering potted plants of the new variety, and a close-up of the flowers.

# DETAILED BOTANICAL DESCRIPTION

The measurements were taken in Gilroy, Calif. USA, in August 2007 on plants that were growing in 14 inch containers containing 3 plants each. Cultures of these plants started in the greenhouse in April 2007 and were transferred to the outdoor containers in early June 2007. The plants were approximately 4 months old.

Color chart used: The Royal Horticultural Society Colour Chart (R.H.S.) 2001

### BRIEF SUMMARY OF INVENTION

The following observations, measurements, and comparisons describe plants grown outside in Gilroy, Calif., USA. The following traits have been repeatedly observed and are determined to be basic characteristics of the new variety. The combination of these characteristics distinguishes this *Pelargonium* as a new and distinct variety.

## TABLE 1

#### DIFFERENCES BETWEEN THE NEW VARIETY AND A SIMILAR VARIETY 'Fisroweiss' (U.S. Plant 'Amri Whit09' Pat. No. 16,974) More vigorous Less vigorous Vigor Peduncle length Shorter Longer Foliage color Darker green Lighter green Bud length Shorter Longer

40 Plant:

Form, growth and habit.—Upright and rounded growth habit. Well-branched.

3

Plant height.—12–15 cm. Color upper and lower petals, lower surface.—RHS N155B but whiter. Plant height (inflorescence included).—22–24 cm. Plant width.—13–16 cm. Length of upper petals.—2.5 cm. Foliage: Width of upper petals.—2.0 cm. *Petal shape.*—Obovate. Arrangement.—Alternate, simple. Immature, leaf color, upper surface.—Between RHS *Apex shape.*—Rounded. Margin.—Entire. 144A and B. Lower surface: RHS 144B. Mature, leaf color, upper surface.—Closest to RHS Petal texture.—Papillose, both sides. 146A but more green. Lower surface: Closest to RHS *Number of petaloids.*—Sometimes 2–3 smaller in vari-146B. ous sizes. 10 *Length.*—5.3–5.8 cm. Color of petaloids.—RHS N155B but whiter. *Width.*—6.8–7.1 cm. Duration of flowering.—Continuous flowering through-Shape.—Orbicular. out the Summer. Bud (when sepals first divide): Base shape.—Reniform. Color.—RHS N155B but whiter. *Apex shape.*—Rounded. *Margin.*—Slightly lobed; irregularly crenate. *Length.*—0.7-0.8 cm. Texture.—Hirsute, glandular hairs. Width.—1.8—2.3 cm. Color of zone.—No zone. *Shape.*—Oblong to elliptic. Color of veins, upper surface.—Indistinct. *Number of sepals.*—5, fused at the base. Color of veins, lower surface.—RHS 146B. Color of sepals.—RHS 144B. Petioles color.—RHS 146B. Length of sepals.—1.3–1.4 cm. Petioles length.—2.6–2.9 cm. Width of sepals.—0.5–0.6 cm. Diameter of petiole.—0.2 cm. Sepal shape.—Lanceolate. *Texture*.—Hirsute. *Apex shape.*—Acute. Stem: *Margins*.—Entire. Color of stem.—RHS 146B. *Texture.*—Long hirsute/villous; glandular hairs. Length of stem.—8–10 cm. Reproductive organs: *Diameter.*—0.6–0.8 cm. Pistil.—1. Length of internodes.—0.7–1.0 cm. *Length.*—1.0–1.1 cm. *Texture*.—Hirsute. Style color.—RHS N155B but whiter. Inflorescence: Style length.-0.2-0.3 cm. *Type.*—An umbel composed of 15–30 flowers and buds. Stigma color.—RHS N155B but whiter. *Umbel diameter.*—6.5–8.0 cm. *Number of anthers.*—About 5. *Umbel depth.*—5.5–6.5 cm. Length filaments.—0.5–0.7 cm. Color of peduncle.—RHS 144A-B. Color of filaments.—RHS N155B but whiter. Length of peduncle.—8.6–10.5 cm. *Pollen amount.*—Moderate. Peduncle diameter.—0.3–0.4 cm. Color of pollen.—RHS N25A. *Texture.*—Short hirsute; glandular hairs. *Fertility/seed set.*—Not observed on this hybrid. Pedicel color.—RHS 146B. Disease/pest resistance: Disease resistance and/or suscepti-Length of pedicel.—3.5–3.8 cm. bility has not been observed on this hybrid. Diameter of pedical.—0.2. *Texture.*—Glandular hairs. What is claimed is: Corolla: *Form.*—Polypetalous, overlapping.

*Number of petals.*—5.

N155B but whiter.

Width of flower.—4.2–4.5 cm.

Color upper and lower petals, upper surface.—RHS

1. A new and distinct variety of *Pelargonium* plant named 'Amri Whit09', substantially as illustrated and described herein.

\* \* \* \* \*

