



US00PP20727P3

(12) **United States Plant Patent**  
**Skelton**(10) **Patent No.:** US PP20,727 P3  
(45) **Date of Patent:** Feb. 9, 2010

- (54) **ACTINIDIA CHINENSIS PLANT NAMED 'S600'**
- (50) Latin Name: *Actinidia chinensis*  
Varietal Denomination: S600
- (76) Inventor: **Donald Skelton**, R D 1, Huntly (NZ)
- (\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **12/072,479**
- (22) Filed: **Feb. 26, 2008**
- (65) **Prior Publication Data**  
US 2009/0217427 P1 Aug. 27, 2009
- (51) **Int. Cl.**  
**A01H 5/00** (2006.01)

- (52) **U.S. Cl.** ..... **Plt./156**
- (58) **Field of Classification Search** ..... Plt./156  
See application file for complete search history.

*Primary Examiner*—June Hwu**(57) ABSTRACT**

A new and distinct *Actinidia chinensis* cultivar named 'S600' is disclosed, characterized by having distinctive yellow flesh fruit, heavy cropping and a seasonal harvest time of late April. Additionally, the new variety produces fruit with very soft, slight pubescence and with a distinctive blunt stylar end shape and square shoulders. The new variety is suitable for commercial production of kiwi fruit.

**3 Drawing Sheets****1**

Latin name of the genus and species: *Actinidia chinensis*.  
Variety denomination: 'S600'.

**BACKGROUND OF THE INVENTION**

The new cultivar is a product of a planned breeding program, begun in 1975 under the direction of the inventor, Donald Alfred Skelton, a citizen of New Zealand. The seed parent is the unpatented, proprietary seedling variety referred to as *Actinidia chinensis* 'A124'. The pollen parent is the unpatented, proprietary seedling variety referred to as *Actinidia chinensis* 'RY'.

Fruit of the new variety was first evaluated in 2000 with favorable results. After the first evaluation, semi-hardwood cuttings were made of 'S600' and were grafted onto 5 seedling rootstocks 3; of *A. chinensis* and 2 of *A. deliciosa*. Evaluation, asexual propagation and grafting all first took place at the inventor's commercial nursery in Rangiriri, New Zealand in 2000. Subsequent evaluations of the variety have shown the characteristics to be true to type.

**SUMMARY OF THE INVENTION**

The cultivar 'S600' has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature, day length, and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'S600.' These characteristics in combination distinguish 'S600' as a new and distinct *Actinidia chinensis* cultivar:

1. Distinctive yellow flesh fruit.
2. Seasonal harvest time of late April.
3. Very soft, downy pubescence on fruit, with weak adherence to the fruit skin.
4. Oblong fruit shape.
5. Fruit with a distinctive blunt stylar end shape and square shoulders.

**COMPARISON TO PARENT**

Plants of the new cultivar 'S600' are similar to plants of the seed parent, *Actinidia chinensis* 'A124' in most horticultural

**2**

characteristics, however, plants of the new cultivar 'S600' produce fruit that is smaller and of a different shape than the seed parent variety. Additionally, fruit of the new variety 'S600' reaches maturity for harvest four to six weeks later than 'A124.'

Plants of the new cultivar 'S600' are similar to plants of the male parent, *Actinida chinensis* 'RY' in most horticultural characteristics, however, plants of the new cultivar 'S600' produce significantly more female flowers than the male parent.

**COMMERCIAL COMPARISON**

The new variety is best compared to the commercial variety, 'Hort16A,' U.S. Plant Pat. No. 11,066. 'S600' is similar to 'Hort16A' in many horticultural characteristics, however, 'S600' produces mature fruit two to three weeks earlier than 'Hort16A.' Additionally, the new variety produces an oblong shaped fruit, compared to ovoid. Fruit of the new variety 'S600' are distinctively blunt on the stylar end, compared to the strong pointed stylar end shape of 'Hort16A.' Fruit shape differences can also be noted in the shoulder shape. Whereas 'S600' produces fruit with strongly squared shoulder, 'Hort16A' fruit have rounded shoulders.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying photograph in FIG. 1 illustrates in full color typical foliage and flowers on plants of 'S600.'

FIG. 2 illustrates in full color typical flowers of 'S600.'

FIG. 3 shows examples of typical fruit harvested from 'S600.'

All photographs were taken of an approximately 8 year old plant. The photographs were taken using conventional techniques and although colors may appear different from actual colors due to light reflectance it is as accurate as possible by conventional photographic techniques.

**DETAILED BOTANICAL DESCRIPTION**

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used.

The following observations and measurements describe 'S600' plants grown under outdoor commercial trial conditions in Rangiriri New Zealand. The growing temperature ranged from 10° C. to 25° C. with no precipitation. Measurements and numerical values represent averages of typical plant types.

Botanical classification: *Actinidia chinensis* 'S600.'

#### PROPAGATION

'S600' can be successfully grafted onto rootstocks of *Actinidia chinensis* or *Actinidia deliciosa*.

#### PLANT

10 Peduncle:

*Length*.—Average 5.1 cm.  
*Color*.—Near RHS Yellow-Green 145A.  
*Texture*.—Smooth.

#### FRUIT

*Color outer pericarp*.—Near RHS Yellow 1C.

*Color inner pericarp*.—Near RHS Yellow 1B.

*Core color*.—Near RHS White 158D.

*Brix at consumption*.—15.8–16.0.

*Brix at harvest*.—8.50%.

*Average weight*.—105 grams.

*Minimum weight*.—87 grams.

*Maximum weight*.—115 grams.

*Length*.—Avg. 59 mm.

*Width*.—Avg. 51 mm.

*Core diameter (maximum)*.—14.0 mm.

*Core diameter (minimum)*.—7.0 mm.

*Locule number*.—30.

*Fruit peduncle length*.—46 mm.

*Fruit peduncle width*.—3.1 mm.

*General shape*.—Oblong.

*Median cross section*.—Circular.

*Stylar end shape*.—Slightly blunt protruding.

*Shoulder shape*.—Squared.

*Calyx ring*.—Present.

*Calyx ring expression*.—Moderate.

*Skin color at harvest*.—Near RHS Grey-Orange 165D.

*Hair on fruit skin*.—Downy.

*Hair adherence to skin*.—Weak.

*Skin adherence to flesh at maturity*.—Moderate.

*Lenticels on fruit*.—Not present.

*Fruit core shape*.—Elliptic.

*Core-woody spike*.—Small.

*Mature seed color*.—Black.

*Dried seed*.—Brown.

*Harvest time*.—Late April.

*Overall cropping quantity*.—Heavy.

#### OTHER CHARACTERISTICS

*Storage life*: Storage life is a minimum of 3 months at 2° C.

*Disease/pest resistance*: Neither resistance nor susceptibility to pathogens and pests common to *Actinidia chinensis* have been observed.

*Temperature tolerance*: Tolerates low temperatures to approximately -8° C. without negative effects, tolerates high temperatures to approximately 35° C. without negative effects.

What is claimed is:

1. A new and distinct cultivar of *Actinidia chinensis* plant named 'S600' as herein illustrated and described.






Fig. 2

