



(12) **United States Plant Patent**
Jacobs

(10) **Patent No.:** **US PP20,824 P2**
(45) **Date of Patent:** **Mar. 9, 2010**

(54) **ALSTROEMERIA PLANT NAMED**
'ZALSAMIN'

(50) Latin Name: *Alstroemeria hybrida*
Varietal Denomination: **Zalsamin**

(75) Inventor: **Henricus Cornelius Maria Jacobs,**
Rijsenhout (NL)

(73) Assignee: **Van Zanten Plants b.v.,** Hillegom (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/291,388**

(22) Filed: **Nov. 7, 2008**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./309**

(58) **Field of Classification Search** **Plt./309**
See application file for complete search history.

Primary Examiner—June Hwu
Assistant Examiner—Louanne C Krawczewicz Myers
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Alstroemeria* plant named
'Zalsamin', characterized by its erect and strong flowering
stems; vigorous growth habit; dark red-colored flowers; and
excellent postproduction longevity.

1 Drawing Sheet

1

Botanical designation: *Alstroemeria hybrida*.
Cultivar denomination: 'Zalsamin'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Alstroemeria* plant, botanically known as *Alstroemeria*
hybrida, commercially used as a cut flower *Alstroemeria*, and
hereinafter referred to by the name 'Zalsamin'.

The new *Alstroemeria* plant is a product of a planned
breeding program conducted by the Inventor in Rijsenhout,
The Netherlands. The objective of the breeding program is to
create new cut flower *Alstroemeria* cultivars with desirable
flower and plant qualities, attractive foliage and flower col-
oration and excellent postproduction longevity.

The new *Alstroemeria* plant originated from a cross-pollina-
tion made by the Inventor in Rijsenhout, The Netherlands
in May, 2003, of a proprietary *Alstroemeria hybrida* selection
identified as code number 02-0107-15, not patented, as the
female, or seed, parent with a proprietary *Alstroemeria*
hybrida selection identified as code number 02-2001n-os, not
patented, as the male, or pollen, parent. The new *Alstroemeria*
was discovered and selected by the Inventor as a flowering
plant from within the progeny of the stated cross-pollination
in a controlled greenhouse environment in Rijsenhout, The
Netherlands in June, 2004.

Asexual reproduction of the new *Alstroemeria* plant by
rhizome divisions in a controlled greenhouse environment in
Rijsenhout, The Netherlands since September, 2004, has
shown that the unique features of this new *Alstroemeria* plant
are stable and reproduced true to type in successive genera-
tions.

SUMMARY OF THE INVENTION

Plants of the new *Alstroemeria* have not been observed
under all possible environmental conditions. The phenotype
may vary somewhat with variations in environment and cul-
tural practices such as temperature and light intensity with-
out, however, any variance in genotype.

The following traits have been repeatedly observed and are
determined to be the unique characteristics of 'Zalsamin'.

2

These characteristics in combination distinguish 'Zalsamin'
as a new and distinct cultivar of *Alstroemeria* plant:

1. Erect and strong flowering stems.
2. Vigorous growth habit.
3. Dark red-colored flowers.
4. Excellent postproduction longevity.

Plants of the new *Alstroemeria* can be compared to plants
of the female parent selection. Plants of the new *Alstroemeria*
differ from plants of the female parent selection primarily in
flower color as plants of the female parent have lighter red-
colored flowers than plants of the new *Alstroemeria*.

Plants of the new *Alstroemeria* can be compared to plants
of the male parent selection. Plants of the new *Alstroemeria*
differ from plants of the male parent selection primarily in
plant height as plants of the male parent selection are shorter
than plants of the new *Alstroemeria*.

Plants of the new *Alstroemeria* can be compared to plants
of *Alstroemeria hybrida* 'Tiesto', not patented. In side-by-
side comparisons conducted in Rijsenhout, The Netherlands,
plants of the new *Alstroemeria* differed primarily from plants
of 'Tiesto' in flower color as plants of 'Tiesto' had flowers that
had larger stripes and a distinct white-colored band.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the over-
all appearance of the new *Alstroemeria*, showing the colors as
true as it is reasonably possible to obtain in colored reproduc-
tions of this type. Colors in the photograph may differ slightly
from the color values cited in the detailed botanical descrip-
tion which accurately describe the colors of the new *Alstro-*
emeria.

The photograph comprises a side perspective view of a
typical flowering stem of 'Zalsamin'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observa-
tions and measurements describe plants of the new *Alstro-*
emeria grown in Rijsenhout, The Netherlands in a glass-
covered greenhouse in ground beds. During the production of
the plants, day temperatures ranged from 15° C. to 20° C.,

night temperatures ranged from 10° C. to 15° C. and light levels averaged 5,000 lux. Plants used for the photograph and description had been growing for one year. The photograph and the description were taken in the late summer. Color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Alstroemeria hybrida* 'Zalsamin'.

Parentage:

Female, or seed, parent.—Proprietary *Alstroemeria hybrida* selection identified as code number 02-0107-15, not patented.

Male or pollen parent.—Proprietary *Alstroemeria hybrida* selection identified as code number 02-2001n-os, not patented.

Propagation:

Type.—By tissue culture.

Time to produce a rooted young plant, summer.—About 40 days.

Time to produce a rooted young plant, winter.—About 60 days.

Root description.—Fibrous, fleshy, thick; close to 155D in color.

Rooting habit.—Freely branching; moderately dense.

Rhizomes.—Shape: Elongate; rounded. Length: About 10 cm to 30 cm. Diameter: About 3 mm to 10 mm. Texture: Smooth. Color: Close to 155D.

Plant description:

Plant and growth habit.—Upright; freely branching, bushy appearance; vigorous growth habit.

Time from planting to harvest of cut flowers.—About 80 to 90 days.

Number of flowering stems produced per year.—About 200 to 240.

Plant height.—About 135 cm to 160 cm.

Plant diameter (spread).—About 35 cm.

Flowering stem description:

Aspect.—Erect.

Length.—About 120 cm to 147 cm.

Diameter.—About 6 mm to 11 mm.

Internode length.—About 2 cm to 6.5 cm.

Strength.—Strong.

Texture.—Smooth, glabrous.

Color.—Close to 147B.

Foliage description:

Appearance.—Leaves asymmetrical; sessile.

Length.—About 19.6 cm to 24 cm.

Width.—About 3.2 cm to 4.6 cm.

Shape.—Lanceolate.

Apex.—Acute.

Base.—Attenuate.

Margin.—Entire; slightly undulate.

Texture, upper and lower surfaces.—Smooth, glabrous.

Venation pattern.—Parallel.

Color.—Developing and fully developed leaves, upper surface: Close to 147A; venation, close to 147A. Developing and fully developed leaves, lower surface: Close to 147B; venation, close to 147B.

Flower description:

Flower type and habit.—Single cup-shaped flowers arranged in compound umbels. Flowers face mostly outwardly. Perianth segments separate. Freely and continuously flowering.

Natural flowering season.—Flowering continuous during the spring in The Netherlands.

Fragrance.—Not detected.

Flower longevity on the plant.—About four weeks; flowers not persistent.

Flower longevity as a cut flower.—About 12 to 16 days.

Flower buds (showing color).—Length: About 5.5 cm to 6 cm. Diameter: About 1.3 cm to 1.8 cm. Shape: Roughly ovoid. Color: Close to 187B.

Umbel height.—About 14 cm to 16 cm.

Umbel diameter.—About 20 cm to 23 cm.

Number of flowers per umbel.—About 10 to 24.

Flower diameter.—About 6.5 cm to 7.5 cm.

Flower length (height).—About 7.5 cm to 8.5 cm.

Flower depth.—About 7.5 cm to 8 cm.

Perianth.—Arrangement: Six arranged in two whorls, each whorl with two lateral and one median segments. Outer perianth, lateral segments: Length: About 6.2 cm to 6.6 cm. Width: About 3.4 cm to 3.7 cm. Shape: Obovate. Apex: Embedded pointed. Base: Attenuate. Margin: Entire; weakly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 46B; towards the apex, greenish overtones. Color, when opening and fully opened, lower surface: Close to 46B; towards the apex and venation, greenish overtones. Outer perianth, median segment: Length: About 7 cm to 7.4 cm. Width: About 3.4 cm to 3.8 cm. Shape: Obovate. Apex: Embedded pointed. Base: Attenuate. Margin: Entire; weakly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 46B; towards the apex, greenish overtones. Color, when opening and fully opened, lower surface: Close to 46B; towards the apex and venation, greenish overtones. Inner perianth, lateral segments: Length: About 6.9 cm to 7.5 cm. Width: About 1.8 cm to 2 cm. Shape: Elliptical. Apex: Wishbone-shaped. Base: Attenuate. Margin: Entire; weakly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 46B; center, close to 162B; stripes, close to 183A. Color, when opening and fully opened, lower surface: Close to 46B. Inner perianth, median segment: Length: About 5.8 cm to 6.2 cm. Width: About 1.7 cm to 2 cm. Shape: Elliptical. Apex: Wishbone-shaped. Base: Attenuate. Margin: Entire; weakly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 46B; stripes, close to 183A. Color, when opening and fully opened, lower surface: Close to 46B.

Pedicels.—Length: About 6 cm to 10 cm. Diameter: About 2 mm to 3 mm. Strength: Strong. Angle: Erect to about 30° C. from vertical. Texture: Smooth, glabrous. Color: Close to 147B.

Reproductive organs.—Stamens: Quantity per flower: Six. Anther shape: Elliptical. Anther length: About 8 mm. Anther diameter: About 2 mm. Anther color: Close to 177B. Pollen amount: Abundant. Pollen color: Close to 152B. Pistils: Quantity per flower: One. Style length: About 4.5 cm to 5 cm. Style color: Close to 59D. Stigma color: Close to 60B. Ovary color: Close to 146C.

Fruit/seed.—Fruit and seed development has not been observed.

Disease/pest resistance: Plants of the new *Alstroemeria* have not been observed to be resistant to pathogens and pests common to *Alstroemeria*.
Temperature tolerance: Plants of the new *Alstroemeria* have been observed to tolerate temperatures from about -5° C. to about 40° C.

It is claimed:
1. A new and distinct *Alstroemeria* plant named 'Zalsamin' as illustrated and described.

* * * * *

