



US00PP20836P2

(12) **United States Plant Patent**
Jacobs

(10) **Patent No.:** **US PP20,836 P2**
(45) **Date of Patent:** **Mar. 16, 2010**

(54) **ALSTROEMERIA PLANT NAMED**
'ZALSANAO'
(50) Latin Name: *Alstroemeria hybrida*
Varietal Denomination: **Zalsanao**
(75) Inventor: **Henricus Cornelius Maria Jacobs,**
Rijsenhout (NL)
(73) Assignee: **Van Zanten Plants b.v.,** Hillegom (NL)
(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.
(21) Appl. No.: **12/291,387**
(22) Filed: **Nov. 7, 2008**
(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./309**
(58) **Field of Classification Search** **Plt./309**
See application file for complete search history.

Primary Examiner—June Hwu
Assistant Examiner—Luanne C Krawczewicz Myers
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Alstroemeria* plant named
'Zalsanao', characterized by its erect and strong flowering
stems; vigorous growth habit; red-colored flowers; and excel-
lent postproduction longevity.

1 Drawing Sheet

1

Botanical designation: *Alstroemeria hybrida*.
Cultivar denomination: 'Zalsanao'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Alstroemeria* plant, botanically known as *Alstroemeria*
hybrida, commercially used as a cut flower *Alstroemeria*, and
hereinafter referred to by the name 'Zalsanao'.

The new *Alstroemeria* plant is a product of a planned
breeding program conducted by the Inventor in Rijsenhout,
The Netherlands. The objective of the breeding program is to
create new cut flower *Alstroemeria* cultivars with desirable
flower and plant qualities, attractive foliage and flower col-
oration and excellent postproduction longevity.

The new *Alstroemeria* plant originated from a cross-pollin-
ation made by the Inventor in Rijsenhout, The Netherlands
in May, 2003, of a proprietary *Alstroemeria hybrida* selection
identified as code number 02-0030-03, not patented, as the
female, or seed, parent with a proprietary *Alstroemeria*
hybrida selection identified as code number 02-2001n-os, not
patented, as the male, or pollen, parent. The new *Alstroemeria*
was discovered and selected by the Inventor as a flowering
plant from within the progeny of the stated cross-pollination
in a controlled greenhouse environment in Rijsenhout, The
Netherlands in June, 2004.

Asexual reproduction of the new *Alstroemeria* plant by
rhizome divisions in a controlled greenhouse environment in
Rijsenhout, The Netherlands since September, 2004, has
shown that the unique features of this new *Alstroemeria* plant
are stable and reproduced true to type in successive genera-
tions.

SUMMARY OF THE INVENTION

Plants of the new *Alstroemeria* have not been observed
under all possible environmental conditions. The phenotype
may vary somewhat with variations in environment and cul-
tural practices such as temperature and light intensity with-
out, however, any variance in genotype.

The following traits have been repeatedly observed and are
determined to be the unique characteristics of 'Zalsanao'.
These characteristics in combination distinguish 'Zalsanao'
as a new and distinct cultivar of *Alstroemeria* plant:

1. Erect and strong flowering stems.
2. Vigorous growth habit.
3. Red-colored flowers.
4. Excellent postproduction longevity.

2

Plants of the new *Alstroemeria* can be compared to plants
of the female parent selection. Plants of the new *Alstroemeria*
differ from plants of the female parent selection primarily in
flower color as flowers of plants of the female parent have a
distinct white-colored band.

Plants of the new *Alstroemeria* can be compared to plants
of the male parent selection. Plants of the new *Alstroemeria*
differ from plants of the male parent selection primarily in
plant height as plants of the male parent selection are taller
than plants of the new *Alstroemeria*.

Plants of the new *Alstroemeria* can be compared to plants
of *Alstroemeria hybrida* 'Tiesto', not patented. In side-by-
side comparisons conducted in Rijsenhout, The Netherlands,
plants of the new *Alstroemeria* differed primarily from plants
of 'Tiesto' in flower color as plants of 'Tiesto' had flowers that
had a distinct white-colored band.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the over-
all appearance of the new *Alstroemeria*, showing the colors as
true as it is reasonably possible to obtain in colored reproduc-
tions of this type. Colors in the photograph may differ slightly
from the color values cited in the detailed botanical descrip-
tion which accurately describe the colors of the new *Alstro-*
emeria.

The photograph comprises a side perspective view of a
typical flowering stem of 'Zalsanao'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observa-
tions and measurements describe plants of the new *Alstro-*
emeria grown in Rijsenhout, The Netherlands in a glass-
covered greenhouse in ground beds. During the production of
the plants, day temperatures ranged from 15° C. to 20° C.,
night temperatures ranged from 10° C. to 15° C. and light
levels averaged 5,000 lux. Plants used for the photograph and
description had been growing for one year. The photograph
and the description were taken in the late summer. Color
references are made to The Royal Horticultural Society
Colour Chart, 1995 Edition, except where general terms of
ordinary dictionary significance are used.

Botanical classification: *Alstroemeria hybrida* 'Zalsanao'.
Parentage:

Female, or seed, parent.—Proprietary *Alstroemeria hybrida* selection identified as code number 02-0030-03, not patented.

Male or pollen parent.—Proprietary *Alstroemeria hybrida* selection identified as code number 02-2001n-os, not patented.

Propagation:

Type.—By tissue culture.

Time to produce a rooted young plant, summer.—About 40 days.

Time to produce a rooted young plant, winter.—About 60 days.

Root description.—Fibrous, fleshy, thick; close to 155D in color.

Rooting habit.—Freely branching; moderately dense.

Rhizomes.—Shape: Elongate; rounded. Length: About 10 cm to 30 cm. Diameter: About 3 mm to 10 mm. Texture: Smooth. Color: Close to 155D.

Plant description:

Plant and growth habit.—Upright; freely branching, bushy appearance; vigorous growth habit.

Time from planting to harvest of cut flowers.—About 80 to 90 days.

Number of flowering stems produced per year.—About 220 to 260.

Plant height.—About 120 cm to 140 cm.

Plant diameter (spread).—About 35 cm.

Flowering stem description:

Aspect.—Erect.

Length.—About 106 cm to 124 cm.

Diameter.—About 6 mm to 11 mm.

Internode length.—About 1.5 cm to 7.5 cm.

Strength.—Strong.

Texture.—Smooth, glabrous.

Color.—Close to 146B.

Foliage description:

Appearance.—Leaves asymmetrical; sessile.

Length.—About 17.8 cm to 20.6 cm.

Width.—About 3.2 cm to 4.6 cm.

Shape.—Lanceolate.

Apex.—Acute.

Base.—Attenuate.

Margin.—Entire; slightly undulate.

Texture, upper and lower surfaces.—Smooth, glabrous.

Venation pattern.—Parallel.

Color.—Developing and fully developed leaves, upper surface: Close to 147A; venation, close to 147A. Developing and fully developed leaves, lower surface: Close to 147A; venation, close to 147A.

Flower description:

Flower type and habit.—Single cup-shaped flowers arranged in compound umbels. Flowers face mostly outwardly. Perianth segments separate. Freely and continuously flowering.

Natural flowering season.—Flowering continuous during the spring in The Netherlands.

Fragrance.—Not detected.

Flower longevity on the plant.—About four weeks; flowers not persistent.

Flower longevity as a cut flower.—About 12 to 16 days.

Flower buds (showing color).—Length: About 4.5 cm to 5 cm. Diameter: About 2 cm to 2.5 cm. Shape: Roughly ovoid. Color: Close to 187B.

Umbel height.—About 14 cm to 17 cm.

Umbel diameter.—About 18 cm to 21 cm.

Number of flowers per umbel.—About 8 to 32.

Flower diameter.—About 6 cm to 6.5 cm.

Flower length (height).—About 6 cm to 6.5 cm.

Flower depth.—About 6 cm to 7 cm.

Perianth.—Arrangement: Six arranged in two whorls, each whorl with two lateral and one median segments. Outer perianth, lateral segments: Length: About 5.4 cm to 6.4 cm. Width: About 3.1 cm to 3.5 cm. Shape: Obovate. Apex: Embedded pointed. Base: Attenuate. Margin: Entire; weakly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 45B; towards the apex, greenish overtones; occasional stripes, close to 183A. Color, when opening and fully opened, lower surface: Close to 45B; towards the apex and venation, greenish overtones; occasional stripes, close to 183A. Outer perianth, median segment: Length: About 6 cm to 6.7 cm. Width: About 3.1 cm to 3.3 cm. Shape: Obovate. Apex: Embedded pointed. Base: Attenuate. Margin: Entire; weakly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 45B; towards the apex, greenish overtones; stripes, close to 183A. Color, when opening and fully opened, lower surface: Close to 45B; towards the apex and venation, greenish overtones; occasional stripes, close to 183A. Inner perianth, lateral segments: Length: About 6.2 cm to 6.5 cm. Width: About 1.7 cm to 2 cm. Shape: Oblanceolate. Apex: Wishbone-shaped. Base: Attenuate. Margin: Entire; weakly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 45B; center, close to 36C; stripes, close to 183A. Color, when opening and fully opened, lower surface: Close to 45B; center, close to 36C; occasional stripes, close to 183A. Inner perianth, median segment: Length: About 5.3 cm to 5.5 cm. Width: About 1.8 cm to 2 cm. Shape: Elliptical. Apex: Wishbone-shaped. Base: Attenuate. Margin: Entire; weakly undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 45B; stripes, close to 183A. Color, when opening and fully opened, lower surface: Close to 45B.

Pedicels.—Length: About 4 cm to 8 cm. Diameter: About 2 mm to 3 mm. Strength: Strong. Angle: Erect to about 30° C. from vertical. Texture: Smooth, glabrous. Color: Close to 146A.

Reproductive organs.—Stamens: Quantity per flower: Six. Anther shape: Elliptical. Anther length: About 8 mm. Anther diameter: About 2 mm. Anther color: Close to 187A. Pollen amount: Abundant. Pollen color: Close to 187A. Pistils: Quantity per flower: One. Style length: About 4.5 cm to 5 cm. Style color: Close to 47A; towards the apex, close to 46A. Stigma color: Close to 46A. Ovary color: Close to 146B.

Fruit/seed.—Fruit and seed development has not been observed.

Disease/pest resistance: Plants of the new *Alstroemeria* have not been observed to be resistant to pathogens and pests common to *Alstroemeria*.

Temperature tolerance: Plants of the new *Alstroemeria* have been observed to tolerate temperatures from about -5° C. to about 40° C.

It is claimed:

1. A new and distinct *Alstroemeria* plant named 'Zalsanao' as illustrated and described.

* * * * *

