

(12) **United States Plant Patent**
Laughner

(10) **Patent No.:** **US PP21,122 P2**
(45) **Date of Patent:** **Jun. 29, 2010**

(54) **OSTEOSPERMUM PLANT NAMED**
'BALVOYELO'

(50) Latin Name: *Osteospermum ecklonis*
Varietal Denomination: **Balvoyelo**

(75) Inventor: **Linda Laughner**, Ventura, CA (US)

(73) Assignee: **Ball Horticultural Company**, West
Chicago, IL (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/460,609**

(22) Filed: **Jul. 22, 2009**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./360**

(58) **Field of Classification Search** **Plt./360**
See application file for complete search history.

Primary Examiner—Annette H Para

Assistant Examiner—Louanne C Krawczewicz Myers

(74) *Attorney, Agent, or Firm*—Audrey Charles

(57) **ABSTRACT**

A new and distinct cultivar of *Osteospermum* plant named
'Balvoyelo', characterized by its bright yellow-colored flow-
ers, medium green-colored foliage, and moderately vigorous
and mounded growth habit that becomes spreading in the
garden, is disclosed.

1 Drawing Sheet

1

Latin name of genus and species of plant claimed:
Osteospermum ecklonis.

Variety denomination: 'Balvoyelo'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Osteospermum* plant botanically known as *Osteospermum*
ecklonis and hereinafter referred to by the cultivar name 'Bal-
voyelo'.

The new cultivar originated in a controlled breeding pro-
gram in Santa Paula, Calif. during August 2006. The objective
of the breeding program was the development of *Osteosper-*
mum cultivars that are freely flowering with unique flower
coloration and a freely branching, mounded growth habit that
becomes spreading in the garden.

The new *Osteospermum* cultivar is the result of cross-
pollination. The female (seed) parent of the new cultivar is the
proprietary *Osteospermum ecklonis* breeding selection des-
ignated 10512-1, not patented, characterized by its light yel-
low-colored flowers, medium green-colored foliage, and
moderately vigorous, upright growth habit. The male (pollen)
parent of the new cultivar is the proprietary *Osteospermum*
ecklonis breeding selection designated 10013-1, not patented,
characterized by its bright yellow-colored flowers, medium
green-colored foliage, and moderately vigorous, trailing
growth habit. The new cultivar was discovered and selected as
a single flowering plant within the progeny of the above stated
cross-pollination during February 2007 in a controlled envi-
ronment at Santa Paula, Calif.

Asexual reproduction of the new cultivar by terminal stem
cuttings since February 2007 at Santa Paula, Calif. and West
Chicago, Ill. has demonstrated that the new cultivar repro-
duces true to type with all of the characteristics, as herein
described, firmly fixed and retained through successive gen-
erations of such asexual propagation.

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have been
repeatedly observed and can be used to distinguish 'Bal-
voyelo' as a new and distinct cultivar of *Osteospermum* plant:

1. Bright yellow-colored flowers;
2. Medium green-colored foliage; and

2

3. Moderately vigorous, and mounded growth habit that
becomes spreading in the garden.

Plants of the new cultivar differ from plants of the female
parent primarily in flower color, flower size, floriferousness
and growth habit. Plants of the new cultivar have a darker and
brighter flower color, larger and more flowers, and more
spreading growth habit. Plants of the new cultivar differ from
plants of the male parent primarily in having a more mounded
growth habit.

Of the many commercially available *Osteospermum* culti-
vars, the most similar in comparison to the new cultivar is
Symphony Lemon 'Seikilrem', U.S. Plant Pat. No. 13,407.
However, in side by side comparisons, plants of the new
cultivar differ from plants of 'Seikilrem' in at least the fol-
lowing characteristics:

1. Plants of the new cultivar have a larger inflorescence
diameter than plants of 'Seikilrem';
2. Plants of the new cultivar have a ray floret color different
from plants of 'Seikilrem';
3. Plants of the new cultivar are taller, as measured to the
top of the plant plane, than plants of 'Seikilrem'; and
4. Plants of the new cultivar have longer peduncles than
plants of 'Seikilrem'.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it is
reasonably possible to make the same in color illustrations of
this type, typical flower and foliage characteristics of the new
cultivar. Colors in the photographs differ slightly from the
color values cited in the detailed description, which accu-
rately describes the colors of 'Balvoyelo'. The plants were
grown in 4.5 inch pots for 7 weeks in a greenhouse at West
Chicago, Ill.

FIG. 1 illustrates a side view of the overall growth and
flowering habit of 'Balvoyelo'.

FIG. 2 illustrates a close-up view of an individual inflores-
cence of 'Balvoyelo'.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible
environmental conditions to date. Accordingly, it is possible

that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2001 edition, except where general color terms of ordinary significance are used. The color values were determined in May 2009 under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe plants produced from cuttings from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown at West Chicago, Ill. in 4.5 inch pots for 7 weeks utilizing a soilless growth medium. Greenhouse temperatures were maintained at approximately 70° F. to 77° F. (21° C. to 25° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Greenhouse light levels of 2,500 footcandles to 6,000 footcandles were maintained during the day. Measurements and numerical values represent averages of typical plants.

Botanical classification: *Osteospermum ecklonis* cultivar Balvoyelo.

Parentage:

Female parent.—Proprietary *Osteospermum ecklonis* breeding selection designated 10512-1, not patented.

Male parent.—Proprietary *Osteospermum ecklonis* breeding selection designated 10013-1, not patented.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 9 to 12 days.

Time to produce a rooted cutting.—Approximately 28 to 32 days.

Root description.—Fibrous.

Rooting habit.—Freely branching.

Plant description:

Commercial crop time.—Approximately 10 to 13 weeks from a rooted cutting to finish in a 10 cm pot.

Growth habit and general appearance.—Moderately vigorous and mounded growth habit that becomes spreading in the garden.

Size.—Height from soil level to top of plant plane: Approximately 23.1 cm. Width: Approximately 22.3 cm.

Branching habit.—Freely branching, pinching enhances branching. Quantity of main branches per plant: Approximately 4.

Branch.—Strength: Strong. Length to base of peduncle: Approximately 5.1 cm. Diameter: Approximately 3.0 mm. Length of central internode: Approximately 3.3 mm. Texture: Glabrous. Color of young and mature stems: 144B.

Foliage description:

General description.—Quantity of leaves per main branch: Approximately 10. Fragrance: Pungent. Form: Simple. Arrangement: Alternate.

Leaves.—Aspect: Acute angle to stem becoming perpendicular to an obtuse angle with age. Shape: Spatulate. Margin: Widely dentate. Apex: Acute. Base: Attenuate. Venation pattern: Pinnate. Length of mature leaf: Approximately 6.0 cm. Width of mature leaf: Approximately 1.9 cm. Texture of upper surface: Sparsely pubescent. Texture of lower surface: Sparsely glandular pubescent. Gland color: Colorless. Color of upper surface of young foliage: 137A

with venation of 145C. Color of lower surface of young foliage: 138B with venation of 144D. Color of upper surface of mature foliage: 137A with venation of 145B. Color of lower surface of mature foliage: Between 138A and 138B with venation of 144D.

Petiole.—Length: Approximately 2.5 cm. Diameter: Approximately 3.0 mm.

Texture: Mixture of glandular and non-glandular pubescence. Gland color: Colorless. Color: 145B.

Flowering description:

Flowering habit.—‘Balvoyelo’ is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

Lastingness of individual inflorescence on the plant.—Approximately 8 to 10 days.

Inflorescence description:

General description.—Type: Solitary, composite. Persistent. Shape: Round. Aspect: Facing upward. Arrangement: Terminal, held above the foliage, arising from leaf axils. Disc and ray florets develop acropetally on a capitulum. Fragrance: Pungent. Quantity per plant: Approximately 3. Diameter: Approximately 7.2 cm. Depth: Approximately 1.2 cm.

Peduncle.—Strength: Strong. Aspect: Erect. Length: Approximately 17.2 cm. Diameter: Approximately 2.0 mm. Texture: Glandular pubescent. Gland color: Colorless. Color: 144B.

Bud.—Rate of opening: Generally takes 4 to 6 days for bud to progress from first color to fully open flower. Quantity per plant: Approximately 4.

Bud just before opening.—Shape: Ovoid. Length: Approximately 1.6 cm. Diameter: Approximately 1.1 cm. Color: 154B.

Ray florets.—Quantity per inflorescence: Approximately 21. Arrangement: Imbricate in a single whorl. Aspect: Initially at acute angle to disc, becoming perpendicular at maturity. Shape: Oblanceolate. Margin: Entire. Apex: Emarginate. Base: Attenuate. Length: Approximately 3.6 cm. Width: Approximately 9.0 mm. Texture of upper surface: Glabrous. Texture of lower surface: Glabrous with a densely pubescent base. Color of upper surface when first open: Darker than 5A. Color of lower surface when first open: Darker than 9B. Color of upper surface when fully open: 5A. Color of lower surface when fully open: 9B.

Disc florets.—Quantity per inflorescence: Approximately 97. Arrangement: Massed in center of inflorescence. Shape: Tubular. Margin: Entire. Apex: Five acute tips. Base: Attenuate, fused. Length: Approximately 9.0 mm. Diameter at apex: Approximately 2.0 mm. Diameter at base: Approximately 1.0 mm. Texture: Glabrous with a sparsely pubescent outer surface. Color when fully open: 12A at tips with center of pure white and base of 144D.

Disc.—Diameter: Approximately 1.4 cm. Depth: Approximately 9.0 mm.

Receptacle.—Shape: Conical. Height: Approximately 2.0 mm. Diameter: Approximately 4.0 mm. Color: 155D.

Phyllaries.—Quantity per inflorescence Approximately 21. Arrangement: In a single whorl. Shape: Linear to lanceolate. Margin: Entire. Apex: Acute. Base: Truncate. Length: Approximately 1.2 cm. Width: Approximately 2.0 mm. Texture of upper or inner surface:

Densely pubescent at tip. Texture of lower or outer surface: Glandular pubescent. Gland color: Colorless. Color of upper or inner surface: 138B. Color of lower or outer surface: 138A.

Reproductive organs.—Androecium: Present on disc florets only. Stamen quantity: 5 per flower, fused around the style. Stamen length: Approximately 5.0 mm. Anther shape: Linear. Anther length: Approximately 2.0 mm. Anther color: 21B. Pollen amount: Abundant. Pollen color: 21A. Gynoecium: Present on ray and disc florets. Pistil quantity: 1 per floret. Pistil length: Approximately 9.0 mm. Stigma shape: Rounded. Stigma length: Less than 1 mm. Stigma

color: 12A. Style length: Approximately 7.0 mm. Style color: 155D. Ovary length: Approximately 2.0 mm. Ovary color: 144D.

Seed and fruit production: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Osteospermum* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Osteospermum* plant named 'Balvoyelo', substantially as herein shown and described.

* * * * *



FIG. 1




FIG. 2