

(12) **United States Plant Patent**
Schröder

(10) **Patent No.:** **US PP21,196 P3**
(45) **Date of Patent:** **Aug. 10, 2010**

(54) **VARIETY OF *CUPHEA* PLANT NAMED
‘VIEBURG’**

(50) Latin Name: *Cuphea llavea*
Varietal Denomination: **Vieburg**

(75) Inventor: **Ralf Schröder**, Lüdinghausen (DE)

(73) Assignee: **Grünewald Veredelings B.V.**,
’s Gravenzande (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 171 days.

(21) Appl. No.: **12/283,767**

(22) Filed: **Sep. 16, 2008**

(65) **Prior Publication Data**

US 2010/0071105 P1 Mar. 18, 2010

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./420**

(58) **Field of Classification Search** **Plt./420**
See application file for complete search history.

Primary Examiner—Annette H Para

(74) *Attorney, Agent, or Firm*—The Webb Law Firm

(57) **ABSTRACT**

A new and distinct *Cuphea llavea* plant named ‘Vieburg’ with
large, reddish-purple bicolored flowers and an upright growth
habit.

2 Drawing Sheets

1

Botanical classification: *Cuphea llavea*.
Varietal denomination: ‘Vieburg’.

BACKGROUND OF THE INVENTION

The present invention comprises a new and distinct cultivar
of *Cuphea llavea* known by the varietal name ‘Vieburg’. The
new variety was discovered in ’s Gravenzande, The Nether-
lands on Jul. 7, 2006. The new variety has been asexually
reproduced via tip cuttings and the first date of asexual repro-
duction was Aug. 15, 2006 in ’s Gravenzande, The Nether-
lands. The new variety is the result of a planned breeding
program cross between unpatented *Cuphea llavea* plant
Seedling C8 (female parent) and unpatented *Cuphea llavea*
plant Seedling C5-2 (male parent). The purpose of the breed-
ing program was to develop new varieties with bigger flowers
and a broad range of color. The new variety exhibits strong
growth and an upright habit that are similar to both parents,
but does have a bigger flowers and an improved flower color
when compared to both parents. ‘Vieburg’ exhibits similar
vigor to *Cuphea llavea* ‘Tiny Winie’ (unpatented and also
known as ‘Torpedo’), but has bigger flowers, different flower
color and more upright growth than ‘Tiny Winie’. The new
variety has been trial and field tested and has been found to
retain its distinctive characteristics and remain true to type
through successive propagations. The following characteris-
tics distinguish ‘Vieburg’ from other *Cuphea llavea* varieties
known to the breeder:

1. Large flowers;
2. Bicolored flowers; and
3. Improved upright growth.

DESCRIPTION OF THE DRAWINGS

The accompanying photographic drawings illustrate the
new variety, with the color being as nearly true as is possible
with color illustrations of this type:

FIG. 1 is a view of an entire plant of the new variety; and

FIG. 2 is an additional photograph of an entire plant of the
new variety.

2

DESCRIPTION OF THE PLANT

The following detailed description sets forth the character-
istics of a 10 week old ‘Vieburg’ plant grown inside a green-
house in a 12 cm container in The Netherlands. The color
readings were taken indoors under natural light. Color refer-
ences are primarily to the R.H.S. Colour Chart of The Royal
Horticultural Society of London.

PLANT

Time to initiate roots: About 12 days at about 18° C.
Time to develop roots: About 16 days at about 18° C.
Time to produce a finished flowering plant from a rooted
cutting: About 10 weeks in a 12 cm container.
Rooting habit: Good, spreading.
Form: Spreading.
Branching habit: Upright.
Height: 35–40 cm.
Plant diameter: 15 cm.
Vigor: Strong.
Branching habit: Spreading upright, not much branching.
Lateral branches:
Length.—18–24 cm.
Diameter.—2.5–3 mm.
Internode length.—4–6 cm.
Texture.—Smooth and hairy.
Color.—145C.
Foliage:
Arrangement.—Alternate.
Number per branch.—6–8.
Size of leaf.—Length: 5.5–6 cm. Width: 2.5–2.7 cm.
Shape of leaf.—Oval.
Shape of apex.—Acute.
Shape of base.—Round.
Texture.—Smooth with some hairs.
Margin description.—Glossy.
Color.—Young leaves: Upper surface: 138A. Lower sur-
face: 138B. Mature leaves: Upper surface: 137A.
Lower surface: 138B to 138C.

Petiole.—Length: 14–16 mm. Diameter: 2.5–3 mm.
Color: 139D.
Veins.—Venation type: Alternate. Color: Young leaves:
Upper surface: 139D with N77A in the middle vein.
Lower surface: 139D. Mature leaves: Upper surface: 5
138D with N77A in the middle vein. Lower surface:
139C to 139D.

FLOWER

Bud:

Time of description.—2 days before opening.
Shape.—Tube.
Diameter.—4–6 mm.
Length.—20–24 mm.
Color.—N170 with 185A hairs.

Natural flowering season: April–October in The Netherlands.

Flower type and habit: Single and lobed.

Number of flowers per lateral branch: 5–7.

Flower longevity on plant: 7–8 days.

Flower:

Diameter.—50×45 mm.
Height.—25–28 mm.
Throat diameter.—12 mm.

Corolla:

Form.—Actinomorphic; fused petals (lobed).
Appearance.—Open flower with a distance between the
petals.
Petals/lobes.—Number: 6. Length: 15–18 mm. Width:
15 mm. Shape: Oval. Apex: Rounded. Base: Acumi-

nate. Margin: Smooth. Texture and appearance: Very
weak. Color: When opening: Upper surface: N60 and
N79B to N79C. Lower surface: N60 and N79B to
N79C. Fully opened: Upper surface: 60B and 79C.
Lower surface: 60B and 79C.

Throat.—Color: 72B to 72C. Texture: Hairy.

Tube color.—N77A and N77B.

Calyx: No calyx or sepals present.

Fragrance: None.

10 Disease resistance: None observed to date.

Temperature tolerance: From 1° C. to 45° C.

REPRODUCTIVE ORGANS

15 Stamens:

Number per flower.—10.

Filament length.—6–10 mm.

Anthers.—Shape: Oval. Length: 1 mm. Color: 195B.

Pollen.—Color: 195B. Amount: Minimal.

20 Pistils:

Length.—20 mm.

Styles.—Length: 11 mm. Color: 196C.

Stigma.—Shape: Round. Color: 196C.

Seed production: Around 20–30.

25

I claim:

1. A new and distinct variety of *Cuphea llavea* plant named
‘Vieburg’ substantially as shown and described herein.

* * * * *



Fig. 1



Fig. 2