



US00PP21353P2

(12) **United States Plant Patent**
Miyazaki(10) **Patent No.:** US PP21,353 P2
(45) **Date of Patent:** Sep. 28, 2010

- (54) **SENECIO PLANT NAMED ‘SUNSENEPIBA’**
- (50) Latin Name: *Senecio hybrida*
Varietal Denomination: Sunsenepiba
- (75) Inventor: **Kiyoshi Miyazaki**, Shiga (JP)
- (73) Assignee: **Suntory Flowers Ltd.**, Tokyo (JP)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **12/387,560**
- (22) Filed: **May 4, 2009**
- (51) **Int. Cl.**
A01H 5/00 (2006.01)

- (52) **U.S. Cl.** **Plt./480**
- (58) **Field of Classification Search** Plt./480
See application file for complete search history.

Primary Examiner—Kent L Bell
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Senecio* plant named ‘Sunsenepiba’, characterized by its upright and mounded plant habit; freely branching growth habit; early flowering response; freely flowering habit; large daisy-type inflorescences with purple-violet and white bi-colored ray florets and darker purple-colored disc florets.

1 Drawing Sheet

1

Botanical designation: *Senecio hybrida*.
Cultivar denomination: ‘Sunsenepiba’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Senecio* plant, botanically known as *Senecio hybrida*, and hereinafter referred to by the name ‘Sunsenepiba’.

The new *Senecio* plant is a naturally-occurring branch mutation of a proprietary selection of *Senecio hybrida* identified as code number RB325, not patented. The new *Senecio* was discovered and selected by the Inventor on a single flowering plant of the parent selection in a controlled greenhouse environment in Higashiomni, Shiga, Japan in February, 2006.

Asexual reproduction of the new *Senecio* plant by terminal cuttings in a controlled greenhouse environment in Higashiomni, Shiga, Japan since March, 2006, has shown that the unique features of this new *Senecio* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Senecio* have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Sunsenepiba’. These characteristics in combination distinguish ‘Sunsenepiba’ as a new and distinct cultivar of *Senecio*:

1. Upright and mounded plant habit.
2. Freely branching growth habit.
3. Early flowering response.
4. Freely flowering habit.
5. Large daisy-type inflorescences with purple-violet and white bi-colored ray florets and darker purple-colored disc florets.

Plants of the new *Senecio* differ from plants of the parent selection in the following characteristics:

1. Plants of the new *Senecio* flower earlier than plants of the parent selection.

2

2. Plants of the new *Senecio* and the parent selection differ in ray floret color as plants of the parent selection have red purple and white bi-colored ray florets.

Plants of the new *Senecio* can be compared to plants of

5 *Senecio cruentus*×*Senecio heritieri* ‘Sunsenereba’, disclosed in U.S. Plant Pat. No. 12,122. In side-by-side comparisons conducted in Higashiomni, Shiga, Japan, plants of the new *Senecio* differed from plants of ‘Sunsenereba’ in the following characteristics:

- 10 1. Plants of the new *Senecio* had darker green-colored leaves than plants of ‘Sunsenereba’.
2. Plants of the new *Senecio* flowered earlier than plants of ‘Sunsenereba’.
3. Plants of the new *Senecio* were more freely flowering than plants of ‘Sunsenereba’.
4. Plants of the new *Senecio* had shorter peduncles than plants of ‘Sunsenereba’.
5. Plants of the new *Senecio* and ‘Sunsenereba’ differed slightly in ray floret color.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

25 The accompanying photographs illustrate the overall appearance of the new *Senecio* plant. These photographs show the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Senecio*.

30 The photograph at the top of the sheet comprises a side perspective view of a typical flowering plant of ‘Sunsenepiba’ grown in a container.

35 The photograph at the bottom of the sheet is a close-up view of typical inflorescences and inflorescence buds of ‘Sunsenepiba’.

DETAILED BOTANICAL DESCRIPTION

40 The aforementioned photographs, following observations and measurements describe plants grown in containers in Higashiomni, Shiga, Japan during the winter in a polyethylene-covered greenhouse and under conditions and practices

which approximate those generally used in commercial *Senecio* production. During the production of the plants, day temperatures averaged 10° C. and night temperatures averaged 5° C. Measurements and numerical values represent averages for typical flowering plants. Plants were six months old when the photographs and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, Fourth Edition, 2001, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Senecio hybrida* ‘Sunsenepiba’.

Parentage: Naturally-occurring branch mutation of a proprietary selection of *Senecio hybrida* identified as code number RB325, not patented.

Propagation:

- Type*.—Terminal vegetative cuttings.
- Time to initiate roots*.—About one week at 20° C.
- Time to produce a rooted young plant*.—About four weeks at 20° C.
- Root description*.—Fibrous; white in color.
- Rooting habit*.—Freely branching; dense.

Plant description:

- Plant form/growth habit*.—Upright and mounded plant habit; daisy-type inflorescences positioned well above the foliar plane; freely branching habit; vigorous growth habit.
- Plant height*.—About 45.3 cm.
- Plant diameter*.—About 50 cm.
- Lateral branches*.—Length: About 28.8 cm. Diameter: About 5.3 mm. Internode length: About 2 cm. Strength: Strong. Texture: Pubescent. Color: Close to 144B tinted with close to 187A.
- Foliage description*.—Arrangement: Alternate, simple. Length: About 6.9 cm. Width: About 8 cm. Shape: Cordate. Apex: Obtuse. Base: Cordate. Margin: Palmettely lobed; crenate; slightly undulate. Texture, upper surface: Sparsely pubescent. Texture, lower surface: Densely pubescent. Venation pattern: Pinnate; reticulate. Color: Developing and fully expanded leaves, upper surface: Close to 137B; venation, close to 193C. Developing and fully expanded leaves, lower surface: Close to 194B; venation, close to 193C. Petioles: Length: About 8.2 cm. Diameter: About 2.7 mm. Texture, upper and lower surfaces: Sparsely pubescent. Color, upper and lower surfaces: Close to 144B tinted with close to 187A.
- Inflorescence description*:
- Appearance*.—Daisy-type inflorescence form with narrowly elliptic-shaped ray florets; inflorescences positioned above the foliar plane, arising from upper leaf axils; disc and ray florets developing acropetally on a capitulum; inflorescences face mostly upright; freely flowering habit with about 77 inflorescences developing per plant.
- Fragrance*.—Faintly fragrant.

Natural flowering season.—In Higashiomii, Shiga, Japan, plants of the new *Senecio* flower continuously from winter to late spring; early flowering response.

Inflorescence longevity.—Inflorescences last about two weeks on the plant; inflorescences persistent.

Inflorescence bud.—Height: About 9.1 mm. Diameter: About 6.5 mm. Shape: Globose. Color: Close to N187A.

Inflorescence size.—Diameter: About 6.6 cm. Depth (height): About 1.1 cm. Disc diameter: About 1.5 cm.

Ray florets.—Shape: Narrowly elliptic. Length: About 2.8 cm. Width: About 7 mm. Apex: Obtuse to rounded, slightly praemorse. Base: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; satiny. Number of ray florets per inflorescence: About 13 in a single whorl. Color: When opening, upper surface: Towards the apex, close to N74A; towards the base, close to 155C. When opening, lower surface: Close to N81B. Fully opened, upper surface: Towards the apex, close to N80A to N80B, color becoming closer to N80C with development; towards the base, close to 155C. Fully opened, lower surface: Close to N81C.

Disc florets.—Shape: Tubular; apex dentate, five-pointed. Length: About 7.9 mm. Diameter: About 1.8 mm. Number of disc floret per inflorescence: About 150. Color, immature and mature: Close to N78A.

Phyllaries.—Quantity per inflorescence: About 14 in a single whorl. Length: About 7.6 mm. Width: About 1.9 mm. Shape: Lanceolate. Apex: Acute. Base: Fused. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper surface: Close to 146A tinted with close to N186C. Color, lower surface: Close to 146B.

Peduncles.—Length: About 5.6 cm. Diameter: About 1.6 mm. Strength: Strong. Aspect: Mostly upright. Texture: Smooth, glabrous. Color: Close to 187A.

Reproductive organs.—Androecium: Present on disc florets only. Anther shape: Ellipsoidal. Anther color: Close to N79C. Pollen amount: Moderate. Pollen color: Close to 9A. Gynoecium: Present on both ray and disc florets. Stigma shape: Bi-parted. Stigma color: Close to 74C.

Seeds/fruits.—Seed and fruit development have not been observed on plants of the new *Senecio*.

Disease/pest resistance: Plants of the new *Senecio* have not been observed to be resistant to pathogens and pests common to *Senecio*.

Temperature tolerance: Plants of the new *Senecio* have been observed to tolerate temperatures ranging from about 0° C. to about 30° C.

It is claimed:

1. A new and distinct *Senecio* plant named ‘Sunsenepiba’ as illustrated and described.

* * * * *

