



US00PP21674P2

(12) **United States Plant Patent**
Dümmen

(10) **Patent No.:** **US PP21,674 P2**
(45) **Date of Patent:** **Jan. 25, 2011**

(54) **PETUNIA PLANT NAMED ‘DUESWECARA’**

(50) Latin Name: *Petunia*×*hybrida*

Varietal Denomination: **Dueswecara**

(75) Inventor: **Tobias Dümmen**, Rheinberg (DE)

(73) Assignee: **Capital Green Investments Ltd.**, Grand Cayman (KY)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/587,681**

(22) Filed: **Oct. 10, 2009**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./356**

(58) **Field of Classification Search** **Plt./356**

See application file for complete search history.

Primary Examiner—June Hwu

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Petunia* plant named ‘Dueswecara’, characterized by its compact, mounding and outwardly spreading to trailing plant habit; freely branching habit; early and freely flowering habit; large light pink-colored flowers; and good garden performance.

1 Drawing Sheet

1

Botanical designation: *Petunia*×*hybrida*.
Cultivar denomination: ‘DUESWECARA’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Petunia* plant, botanically known as *Petunia*×*hybrida* and hereinafter referred to by the name ‘Dueswecara’.

The new *Petunia* plant is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. The objective of the breeding program is to create new *Petunia* cultivars with numerous unique and attractive flowers.

The new *Petunia* plant originated from a cross-pollination made by the Inventor in July, 2006 in Rheinberg, Germany of a proprietary selection of *Petunia*×*hybrida* identified as code number T04-0063-008, not patented, as the female, or seed, parent with a proprietary selection of *Petunia*×*hybrida* identified as code number T05-4553-002, not patented, as the male, or pollen, parent. The new *Petunia* plant was discovered and selected by the Inventor as a single flowering plant within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rheinberg, Germany in May, 2008.

Asexual reproduction of the new *Petunia* plant by terminal cuttings in a controlled greenhouse environment in Rheinberg, Germany since May, 2008, has shown that the unique features of this new *Petunia* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Petunia* have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment and cultural practices such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Dueswecara’. These characteristics in combination distinguish ‘Dueswecara’ as a new and distinct cultivar of *Petunia*:

1. Compact, mounding and outwardly spreading to trailing plant habit.

2

2. Freely branching habit.
3. Early and freely flowering habit.
4. Large light pink-colored flowers.
5. Good garden performance.

Plants of the new *Petunia* can be compared to plants of the female parent selection. Plants of the new *Petunia* differ primarily from plants of the female parent selection in flower color as plants of the female parent selection have darker-colored flowers. In addition, plants of the female parent selection are more upright than plants of the new *Petunia*.

Plants of the new *Petunia* can be compared to plants of the male parent selection. Plants of the new *Petunia* differ primarily from plants of the male parent selection in flower color as plants of the male parent selection have yellow-colored flowers. In addition, plants of the new *Petunia* are more compact than plants of the male parent selection.

Plants of the new *Petunia* can be compared to plants of the *Petunia*×*hybrida* ‘Keiyeyas’, not patented. In side-by-side comparisons conducted in Rheinberg, Germany, plants of the new *Petunia* differed primarily from plants of ‘Keiyeyas’ in the following characteristics:

1. Plants of the new *Petunia* were more compact than plants of ‘Keiyeyas’.
2. Plants of the new *Petunia* had longer leaves than plants of ‘Keiyeyas’.
3. Plants of the new *Petunia* had larger flowers than plants of ‘Keiyeyas’.
4. Plants of the new *Petunia* and ‘Keiyeyas’ differed in flower color as plants of ‘Keiyeyas’ had yellow-colored flowers.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Petunia* plant, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Petunia* plant. The photograph comprises a close-up view of typical leaves and flowers of ‘Dueswecara’.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown in Rheinberg, Germany, under commercial practice during the summer in 10.5-cm containers in a glass-covered greenhouse with day and night temperatures averaging 18° C. and light levels averaging 4,500 lux. Plants were pinched one time three weeks after planting and were 16 weeks old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Petunia* × *hybrida* 'Dueswecara'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Petunia* × *hybrida* identified as code number T04-0063-008, not patented.

Male, or pollen, parent.—Proprietary selection of *Petunia* × *hybrida* identified as code number T05-4553-002, not patented.

Propagation:

Type.—By terminal cuttings.

Time to initiate roots, summer.—About five days at temperatures of 20° C.

Time to initiate roots, winter.—About seven days at temperatures of 20° C.

Time to produce a rooted young plant, summer.—About three weeks at temperatures of 20° C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures of 20° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Compact, mounding and outwardly spreading to trailing growth habit; freely branching habit with about eight to ten lateral branches developing after pinching; moderately vigorous growth habit.

Plant height.—About 15 cm.

Plant diameter.—About 26 cm.

Lateral branch description:

Length.—About 12 cm.

Diameter.—About 4 mm.

Internode length.—About 2 cm.

Strength.—Moderately strong.

Aspect.—Initially upright to outwardly spreading.

Texture.—Pubescent.

Color.—Close to 144A to 144B.

Foliage description:

Arrangement.—Before flowering, alternate, simple; after flowering, opposite, simple.

Length.—About 5.5 cm.

Width.—About 2.7 cm.

Shape.—Spatulate.

Apex.—Obtuse.

Base.—Attenuate.

Margin.—Entire.

Texture, upper and lower surfaces.—Pubescent; leathery.

Venation pattern.—Pinnate; arcuate.

Color.—Developing leaves, upper surface: Close to 137B. Developing leaves, lower surface: Close to 143A to 143B. Fully expanded leaves, upper surface:

Close to 137A; venation, close to 144B. Fully expanded leaves, lower surface: Close to 137C; venation, close to 144B.

Petiole length.—About 3.8 cm.

Petiole diameter.—About 3.5 mm.

Petiole texture, upper and lower surfaces.—Pubescent.

Petiole color, upper and lower surfaces.—Close to 144A.

Flower description:

Flower arrangement and habit.—Large salverform flowers; single flowers arising from leaf axils; freely flowering habit with usually about 25 to 30 open flowers and flower buds per plant; flowers face mostly upright to outwardly.

Fragrance.—None detected.

Natural flowering season.—Plants flower continuously from late spring into autumn in Germany; early flowering habit, plants typically beginning flowering about nine weeks after planting.

Flower longevity.—Individual flowers last about five to six days on the plant; flowers persistent.

Flower diameter.—About 5.5 cm.

Flower length (height).—About 4.5 cm.

Flower throat diameter.—About 1.2 cm.

Flower tube diameter.—About 2.5 mm.

Flower tube length.—About 3 cm.

Flower bud.—Shape: Ovoid. Length: About 3 cm. Diameter: About 5.8 mm. Color: Close to 149A.

Corolla.—Arrangement: Five petals fused at the base and opening into a flared trumpet. Petal lobe length (from throat): About 2.2 cm. Petal lobe width: About 2.8 cm. Petal shape: Roughly spatulate. Petal apex: Rounded. Petal margin: Entire. Petal texture, upper and lower surfaces: Rippled, glabrous. Throat texture: Rippled, glabrous. Tube texture: Rippled, pubescent. Color: Petal lobe, when opening, upper surface: Close to 69D. Petal lobe, when opening, lower surface: Close to 149A tinted with close to 69D. Petal lobe, fully opened, upper surface: Close to 69D; color becoming closer to 69C with development; venation, close to 66A. Petal lobe, fully opened, lower surface: Close to 69D; venation, close to 66C and 144A. Flower throat: Close to 9A; venation, close to 175A. Flower tube: Close to 9B; venation, close to 187B.

Calyx.—Arrangement: One star-shaped calyx tube with five sepals fused at the base per flower. Sepal length: About 1.4 cm. Sepal width: About 6 mm. Sepal shape: Oblong. Sepal apex: Rounded. Sepal margin: Entire. Sepal texture, upper and lower surfaces: Smooth. Color, immature and mature, upper surface: Close to 137A. Color, immature and mature, lower surface: Close to 137A to 137B.

Peduncles.—Length: About 3.6 cm. Diameter: About 1.5 mm. Strength: Moderately strong. Texture: Smooth. Color: Close to 143A.

Reproductive organs.—Stamens: Quantity: Five per flower. Filament length: About 1.2 cm. Filament color: Close to 145C. Anther shape: Ovate. Anther length: About 1.5 mm. Anther color: Close to 6A to 6B. Pollen amount: Abundant. Pollen color: Close to 154C to 154D. Pistils: Quantity: One per flower. Pistil length: About 2.5 cm. Style length: About 2.1 cm. Style color: Close to 144B to 144C. Stigma shape:

Rounded. Stigma color: Close to 144A. Ovary color: Close to 144A. Seed/fruit: Seed and fruit development have not been observed on plants of the new *Petunia*.

Garden performance: Plants of the new *Petunia* have been observed to have good garden performance and tolerate wind, rain and temperatures ranging from about 5° C. to about 40° C.

Pathogen/pest resistance: Plants of the new *Petunia* have not been observed to be resistant to pathogens and pests common to *Petunia*.

It is claimed:

1. A new and distinct *Petunia* plant named 'Dueswecara' as illustrated and described.

* * * * *

