



US00PP21862P3

(12) **United States Plant Patent**  
**Saville**

(10) **Patent No.:** **US PP21,862 P3**  
(45) **Date of Patent:** **Apr. 12, 2011**

(54) **MINIATURE ROSE PLANT NAMED**  
**'SAVAMONT'**

(50) Latin Name: *Rosa chinensis hybrida*  
Varietal Denomination: **Savamont**

(75) Inventor: **John M. Saville**, Newburyport, MA  
(US)

(73) Assignee: **Nor'East Miniature Roses, Inc.**,  
Rowley, MA (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 2386 days.

(21) Appl. No.: **10/015,143**

(22) Filed: **Dec. 10, 2001**

(65) **Prior Publication Data**

US 2003/0110542 P1 Jun. 12, 2003

(51) **Int. Cl.**  
*A01H 5/00* (2006.01)

(52) **U.S. Cl.** ..... **Plt./122**

(58) **Field of Classification Search** ..... **Plt./122**  
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

PP5,770 P \* 7/1986 De Ruitter ..... **Plt./122**  
PP6,517 P \* 1/1989 Benardella ..... **Plt./116**

\* cited by examiner

*Primary Examiner* — Bruce R. Campell

*Assistant Examiner* — W C Haas

(74) *Attorney, Agent, or Firm* — Christie, Parker & Hale,  
LLP

(57) **ABSTRACT**

A new and distinct Miniature Rose plant having attractive  
Red flowers.

**1 Drawing Sheet**

**1**

Botanical designation: *Rosa chinensis hybrida*.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct variety  
of *Rosa chinensis hybrida*. The new variety was created by  
crossing as seed parent the variety known as 'BENALAV'  
(U.S. Plant Pat. No. 6,517) and as pollen parent, the variety  
known as 'RUMIRED' (U.S. Plant Pat. No. 5,770). The vari-  
etal denomination of the new variety is 'SAVAMONT'.

**SUMMARY OF THE INVENTION**

Among the novel characteristics possessed by the new  
variety which distinguishes it from its parents and all other  
varieties of which I am aware, is its attractive Red flowers.

Asexual reproduction by propagation of vegetative cut-  
tings of the new variety as performed in Essex County, Mass.  
show that the foregoing and other distinguishing characteris-  
tics come true to form and are established and transmitted  
through succeeding propagation.

**COMPARISON WITH PARENTS**

The new variety may be distinguished from its parents in  
flower color. The new variety is much taller and more upright  
than its pollen parent 'RUMIRED' and whereas the flower of  
the new variety is more Blue Red in color, flower of  
'RUMIRED' is more of an Orange Red color. The new  
variety differs from its seed parent BENLAV in flower color.  
Flowers of BENLAV are lavender pink and flowers of the new  
variety are red.

**BRIEF DESCRIPTION OF ILLUSTRATION**

The accompanying photograph illustrates a typical speci-  
men of the new variety and shows the flowering thereof from

**2**

bud to full bloom depicted in color as nearly true as it is  
reasonably possible to make in a color illustration of the  
character. Throughout this specification, color references  
and/or values are based upon The Royal Horticultural Society  
Colour Chart and terminology used in the color descriptions  
herein refers to plate numbers in this color chart except where  
common terms of color definition are employed.

**DESCRIPTION OF THE NEW VARIETY**

The following observations were made of 5 year-old plants  
of the new variety grown in 2-gallon plastic containers in  
Essex County, Mass. during the months of July and August.  
Phenotypic expression may vary with environmental, cultural  
and climatic conditions, as well as differences in conditions  
of light and soil.

**FLOWER**

Blooming habit: Continuous; flowers last about a week on the  
plant and then drop off cleanly with the peduncle.

*Bud*.—Medium; length about 1/2 to 5/8 inch just before  
sepals divide; diameter is approximately 1/2 inch in  
width. Form: Obovate. Color: When sepals first divide  
color near 60A; when petals begin to unfurl, color  
near 60A.

*Sepals*.—Lanceolate with occasional flags; extending  
about 5/8 inch beyond tip of bud. Color: Upper Sur-  
face: Near 144A. Under surface: Near 147C.

*Peduncle*.—Length: About 1 to 1 1/2 inches. Aspect:  
Straight. Strength: Erect. Color: Near 144A; texture  
smooth.

**BLOOM**

Size: Average size of flower when fully expanded about 1 7/8  
inches in diameter.

Borne: Clusters of 2 to 8 and also singly.  
 Form: Opens flat becoming and remaining loosely rolled outward.  
 Petalage: Between 24 to 30.  
 Fragrance: Slight.  
 Persistence: Flower drops off cleanly.  
 Lasting quality: Between 7 to 10 days as cut flower.

## PETALS

Texture: Smooth.  
 Appearance: Inside velvety; outside satiny with slightly raised veins.  
 Form: Broad, spatulate.  
 Size: Length about  $\frac{11}{16}$  inch at point of attachment to tip; outer petal width about  $\frac{3}{4}$  inch.  
 Arrangement: Shingle-like; regularly arranged.  
 Color:  
*During first 2 days.*—Petals at center of flower upper side near 46A, under side near 53A; outer petals upper side near 46A, under side near 60A.  
 In full bloom.—Inner petals upper side between 60A and 53A, underside near 53A; outer petals upper side near 53A, under side near 60A.  
*Base of petals (point of attachment).*—White.  
*General tonality from a distance.*—Dark Red.  
*Color change.*—Very slight as bloom ages. General tonality from a distance: Dark Red even after 3 or more days.  
*Petaloids in center.*—About 2 to 6; usually about  $\frac{1}{8}$  to  $\frac{2}{8}$  inches wide and  $\frac{9}{32}$  to  $\frac{11}{32}$  inches long; shape is obovate with acuminate base; apex tends toward obtuse but often the outer margin appears malformed being emarginate to varying degrees.  
*Color.*—Same as inner petals.  
*Texture.*—Smooth.

## REPRODUCTIVE ORGANS

Arrangement of stamens:  
 Filaments and anthers: Regularly arranged around styles.  
 Stamens: About 100 or more.  
 Filaments: About  $\frac{1}{16}$  to  $\frac{1}{8}$  inch; color Deep Yellow near 20A.  
 Anthers: Near 20B.  
 Pollen: Ample quantity; color near 17A.  
 Pistils: About 30.  
 Styles: Thin, uneven length between  $\frac{1}{16}$  to  $\frac{3}{16}$  inch; color near 53C.  
 Stigmas: Near 27B.  
 Receptacle:  
*Shape.*—Round; surface texture pubescent.  
*Size.*—About  $\frac{5}{16}$  inch.  
*Color.*—Off-White near 196D.  
 Hips:  
*Shape.*—Mostly globular, occasionally apple and pear shaped.  
*Size.*—Average diameter at widest point between  $\frac{13}{16}$  and  $\frac{14}{16}$  inch.  
*Color.*—Some areas near 28B; some areas near 167A.  
*Seeds.*—Seeds do not protrude; surface texture glabrous; color near 19C; seeds appear to be viable;

number of seeds per hip is variable and unknown; size and shape of seeds also unknown.

## PLANT

5 Form: Upright, compact.  
 Growth: Vigorous, uniform branching.  
 Size: At maturity, height between 18 to 21 inches; width between 20 to 24 inches.  
 10 Foliage: Generally 5 leaflets, occasionally 3 or 7 leaflets.  
*Size.*—About  $3\frac{1}{2}$  inches from stem to tip measured along the rachis; about  $1\frac{3}{4}$  inches on mature leaflet at terminal leaflet base to tip.  
*Quantity.*—Abundant.  
 15 *Color.*—New foliage: Upper side near 146A flushed with 187A; under side near 146B flushed with 187A. Mature foliage: Upper side near 147A; under side near 147B.  
*Shape.*—Oval pointed; shape of the base of the leaflets is rounded and the apex is acute.  
*Texture.*—Upper side: Semi-glossy and glabrous. Under side: Matte and glabrous.  
*Edge.*—Serrated.  
*Serration.*—Fine on new foliage becoming somewhat coarse and slightly uneven on older foliage.  
 25 Stipules: Very finely bearded; length is about  $\frac{1}{2}$  the length to the first set of leaflets; upper surface color near 147B, under surface near 146C.  
 Rachis: Smooth texture; upper side color near 147A, under side near 144A.  
 30 *Prickles.*—Occasional; between 2 to 3 prickles on mature leaves, occasionally 5; color near 144C.  
 Resistance to disease: Good resistance to insects and Mildew; average resistance to Blackspot, resistance to Rust not  
 35 observed.

## WOOD

New shoots:  
 40 *Bark.*—Smooth.  
*Wood.*—Near 144A.  
 Old wood:  
*Bark.*—Smooth.  
*Wood.*—Near 146A.  
 45 Flowering stem: Length between  $4\frac{1}{2}$  to  $5\frac{1}{2}$  inches.  
 Thorns:  
*Quantity.*—About 4 per inch on main stalks from base; about 6 to 8 per inch on laterals from stalk; fewer closest to bud/flower.  
 50 *Form.*—Flat base, slightly curved downward to a point; irregularly positioned.  
*Length.*—About  $\frac{1}{4}$  inch.  
*Color.*—Immature thorns: Tip near 161C, at the base near 183D. Mature thorns: Near 165B.  
 55 *Prickles.*—None on main stalks.  
 Winter hardiness: Tested hardy zones 4 to 9; winter protection recommended in Zones 5 and colder.  
 I claim:  
 60 1. A new and distinct Miniature Rose plant of the variety substantially as shown and described.

\* \* \* \* \*

