



(12) **United States Plant Patent**  
**Hofmann et al.**

(10) **Patent No.:** **US PP22,124 P2**  
(45) **Date of Patent:** **Sep. 6, 2011**

(54) **PELARGONIUM PLANT NAMED**  
**‘KUEGRAROLI’**

(50) Latin Name: *Pelargonium crispum*×*Pelargonium grandiflorum*  
Varietal Denomination: **Kuegraroli**

(75) Inventors: **Christa Hofmann**, Leipzig (DE); **Klaus Olbricht**, Dresden (DE); **Katrin Meinel**, Dresden (DE)

(73) Assignee: **Kühne Jungflanzen GbR Claus & Tonsten Kühne**, Dresden (DE)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/798,349**

(22) Filed: **Mar. 31, 2010**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(52) **U.S. Cl.** ..... **Plt./324**

(58) **Field of Classification Search** ..... **Plt./324**  
See application file for complete search history.

*Primary Examiner* — Annette Para

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Pelargonium* plant named ‘Kuegraroli’, characterized by its upright, somewhat outwardly spreading and mounded plant habit; compact and moderately vigorous growth habit; freely branching habit; freely flowering habit; dark purple and red purple bi-colored flowers; no cooling treatment required for flower initiation and development; and good garden performance.

**1 Drawing Sheet**

**1**

Botanical designation: *Pelargonium crispum*×*Pelargonium grandiflorum*.

Cultivar denomination: ‘KUEGRAROLI’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of *Pelargonium* plant, botanically known as *Pelargonium crispum*×*Pelargonium grandiflorum* and hereinafter referred to by the name ‘Kuegraroli’.

The new *Pelargonium* plant is a product of a planned breeding program conducted by the Inventors in Dresden, Germany. The objective of the breeding program is to develop new interspecific *Pelargoniums* that can be grown as potted and/or bedding plants.

The new *Pelargonium* plant originated from a cross-pollination made by the Inventors in Dresden, Germany in 2006 of *Pelargonium crispum* ‘Randy’, not patented, as the female, or seed, parent with an unidentified selection of *Pelargonium grandiflorum*, not patented, as the male, or pollen, parent. The new *Pelargonium* plant was discovered and selected by the Inventors as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Dresden, Germany in December, 2006.

Asexual reproduction of the new *Pelargonium* plant by vegetative terminal cuttings in a controlled greenhouse environment in Dresden, Germany since the spring of 2007, has shown that the unique features of this new *Pelargonium* plant are stable and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

Plants of the new *Pelargonium* have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment and cultural practices such as temperature and light intensity without, however, any variance in genotype.

**2**

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Kuegraroli’. These characteristics in combination distinguish ‘Kuegraroli’ as a new and distinct cultivar of *Pelargonium*:

1. Upright, somewhat outwardly spreading and mounded plant habit.
2. Compact and moderately vigorous growth habit.
3. Freely branching habit; pinching is not required.
4. Freely flowering habit.
5. Dark purple and red purple bi-colored flowers.
6. No cooling treatment required for flower initiation and development.
7. Good garden performance.

Plants of the new *Pelargonium* differ primarily from plants of the female parent, ‘Randy’, in the following characteristics:

1. Plants of the new *Pelargonium* are more upright than and not as trailing as plants of ‘Randy’.
2. Plants of the new *Pelargonium* have thicker stems, larger leaves and larger flowers than plants of ‘Randy’.
3. Flowers of plants of the new *Pelargonium* are held closer together than flowers of plants of ‘Randy’.
4. Plants of the new *Pelargonium* and ‘Randy’ differ in flower color as plants of ‘Randy’ have red purple and white bi-colored flowers.

Plants of the new *Pelargonium* differ primarily from plants of the male parent selection in the following characteristics:

1. Plants of the new *Pelargonium* are more compact than and not as vigorous as plants of the male parent selection.
2. Plants of the new *Pelargonium* have thinner stems, smaller leaves and smaller flowers than plants of the male parent selection.
3. Plants of the new *Pelargonium* flower earlier than plants of the male parent selection.
4. Plants of the new *Pelargonium* do not require a cooling treatment for flower initiation and development whereas plants of the male parent selection do require a cooling treatment for flower initiation and development.



Plants of the new *Pelargonium* can be compared to plants of the *Pelargonium grandiflorum* 'Camvio', disclosed in U.S. Plant Pat. No. 20,421. In side-by-side comparisons conducted in Dresden, Germany, plants of the new *Pelargonium* differed from plants of 'Camvio' in the following characteristics:

1. Plants of the new *Pelargonium* were more compact than and not as vigorous as plants of 'Camvio'.
2. Plants of the new *Pelargonium* flowered about five days earlier than plants of 'Camvio'.
3. Plants of the new *Pelargonium* had larger flowers than plants of 'Camvio'.
4. Plants of the new *Pelargonium* and 'Camvio' differed in flower color as plants of 'Camvio' had red purple and dark red bi-colored flowers.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Pelargonium* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Pelargonium* plant. The photograph comprises a side perspective view of a typical flowering plant of 'Kuegraroli' grown in a container.

#### DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown 12-cm containers in Dresden, Germany in a glass-covered greenhouse during the summer and under conditions which closely approximate commercial *Pelargonium* production. During the production of the plants, day and night temperatures averaged 18° C. and light levels averaged 4,500 lux. Plants were 13 weeks old when the photograph and the description were taken. In the detailed description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Pelargonium crispum* × *Pelargonium grandiflorum* 'Kuegraroli'.

Parentage:

*Female, or seed, parent.*—*Pelargonium crispum*, 'Randy' not patented.

*Male or pollen parent.*—Unidentified selection of *Pelargonium grandiflorum*, not patented.

Propagation:

*Type.*—By vegetative terminal cuttings.

*Time to initiate roots, summer.*—About 18 days at temperatures of 20° C.

*Time to initiate roots, winter.*—About 22 days at temperatures of 22° C.

*Time to produce a rooted young plant, summer.*—About 26 days at temperatures of 18° C.

*Time to produce a rooted young plant, winter.*—About 30 days at temperatures of 20° C.

*Root description.*—Fine, fibrous; creamy white in color.

*Rooting habit.*—Freely branching; dense.

Plant description:

*Plant/growth habit.*—Upright, somewhat outwardly spreading and mounded plant habit; inverted triangle; densely foliated; compact and moderately vigorous growth habit; freely basal habit with about four to six lateral branches developing per plant.

*Plant height, to top of umbels.*—About 20 cm.

*Plant height, to top of leaves.*—About 15 cm.

*Plant width.*—About 18 cm.

*Lateral branches.*—Length: About 6 cm. Diameter:

About 3 mm. Internode length: About 2.5 cm.

Strength: Strong. Texture: Slightly pubescent. Color: Close to 138B.

Foliage description:

*Arrangement.*—Alternate; simple.

*Length.*—About 3.6 cm.

*Width.*—About 4.2 cm.

*Shape.*—Palmate, lobed.

*Apex.*—Cuspidate.

*Base.*—Obtuse.

*Margin.*—Serrate.

*Venation pattern.*—Palmate.

*Texture, upper and lower surfaces.*—Pubescent.

*Color.*—Developing leaves, upper surface: Close to 141B. Developing leaves, lower surface: Close to 139C. Fully expanded leaves, upper surface: Close to 137A; venation, close to 137A. Fully expanded leaves, lower surface: Close to 137D; venation, close to 137D. Petiole: Length: About 1.3 cm. Diameter: About 1.5 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 138B.

Flower description:

*Flower arrangement.*—Single rotate flowers arranged in umbels arising from apical leaf axils; umbels displayed above the foliage on moderately strong peduncles; flowers face mostly upright to outwardly.

*Fragrance.*—Slightly fragrant, aromatic.

*Quantity of flowers.*—Freely flowering habit; about six flowers develop per umbel and about 13 umbels per plant.

*Flowering season.*—In Germany, flowering initiates in the spring and continues until the autumn; plants do not require a cooling treatment for flower initiation and development; early flowering habit, plants begin flowering about 60 days after planting.

*Flower longevity.*—Individual flowers last about six days on the plant; flowers not persistent.

*Umbel height.*—About 7 cm.

*Umbel diameter.*—About 6.5 cm.

*Flower diameter.*—About 5.3 cm by 4.8 cm.

*Flower depth (height).*—About 2.8 cm.

*Flower buds.*—Length: About 1.7 cm. Diameter: About 6 mm. Shape: Ovoid. Color: Close to 138A.

*Petals.*—Quantity per flower: Five arranged in a single whorl. Length: About 3.7 cm to 4 cm. Width: About 1.8 cm to 2.5 cm. Shape: Obovate. Apex: Rounded. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening, upper surface: Close to N79B; towards the margins, close to 71B. When opening, lower surface: Close to N79B; towards the margins, close to 71C. Fully opened, upper surface: Close to N79B; towards the margins, close to 71D; venation, similar to surface color; margin color becoming closer to 70B with development. Fully opened, lower surface: Close to N79B; towards the margins, close to 70B; venation, similar to surface color.

*Sepals.*—Quantity per flower: Five arranged in a single whorl. Length: About 1.8 cm. Width: About 4 mm. Shape: Lanceolate. Apex: Acuminate. Base: Obtuse.

Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper surface: Close to 138B. Color, lower surface: Close to 138A.

*Peduncle (umbel stem).*—Length: About 3.5 cm. Diameter: About 2 mm. Strength: Moderately strong. Angle: Mostly erect. Texture: Smooth, glabrous. Color: Close to N138B.

*Pedicel (individual flower stem).*—Length: About 2 cm. Diameter: About 1.5 mm. Strength: Moderately strong. Angle: Mostly erect. Texture: Smooth, glabrous. Color: Close to 138A.

*Reproductive organs.*—Androecium: Stamen quantity per flower: About seven. Filament length: About 1 cm. Filament color: Close to 71B. Anther length: About 2 mm. Anther shape: Ovate. Anther color: Close to 60B. Pollen amount: Moderate. Pollen color: Close to 32A. Gynoecium: Pistil quantity per flower: One. Pistil length: About 1.6 cm. Stigma shape: Five-parted.

Stigma color: Close to 53A. Style length: About 1 cm. Style color: Close to N66A. Ovary color: Close to 130C.

*Seeds/fruits.*—Seed and fruit development has not been observed on plants of the new *Pelargonium*.

Disease/pest resistance: Plants of the new *Pelargonium* have been observed to be resistant to *Xanthomonas campestris* pv. *pelargonii* and *Puccinia pelargonii-zonalis*. Plants of the new *Pelargonium* have not been observed to be resistant to pests and other pathogens common to *Pelargoniums*.

Garden performance: Plants of the new *Pelargonium* have been observed to have good garden performance and to tolerate wind, rain and temperatures ranging from about 5° C. to about 40° C.

It is claimed:

1. A new and distinct *Pelargonium* plant named 'Kuegraroli' as illustrated and described.

\* \* \* \* \*



