

US00PP22320P3

(12) **United States Plant Patent**
Lavee et al.

(10) **Patent No.:** **US PP22,320 P3**
(45) **Date of Patent:** **Dec. 13, 2011**

(54) **OLIVE TREE NAMED ‘SEPOKA’**

(50) Latin Name: *Olea europaea* L.
Varietal Denomination: **SEPOKA**

(75) Inventors: **Shimon Lavee**, Rehovot (IL); **Benjamin Avidan**, Reshon LeZion (IL); **Yair Manni**, Rosh Haain (IL)

(73) Assignee: **State of Isreal, Ministry of Agriculture & Rural Development Agricultural Research Organization**, Bet Dagan (IL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/589,896**

(22) Filed: **Oct. 30, 2009**

(65) **Prior Publication Data**
US 2011/0107474 P1 May 5, 2011

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./158**

(58) **Field of Classification Search** Plt./158
See application file for complete search history.

(56) **References Cited**

OTHER PUBLICATIONS

Print-out of registration number and grant date information for corresponding IL PBR application No. 3515/03 granted as Registration No. 2712 on Aug. 26, 2007, published by the Ministry of Agriculture & Rural Development in Israel PBR Gazette No. 72, Jan. 1, 2008—Dec. 30, 2008 (3 pages). http://www.moag.gov.il/NR/rdonlyres/8D0AE3A4-851E-4E58-A239-C70D49E646D6/0/ILgaz72_jan09.pdf.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — **Foley & Lardner LLP**

(57) **ABSTRACT**

A new and distinct *Olea europaea* L. olive tree variety named ‘SEPOKA’, particularly characterized as a small to medium-sized tree facilitating hand picking, suitable for intensive orchard growing with harvesting from the ground level with only minimal pruning required, good resistance to *Spilocaea oleagina* and elongated shaped fruit, with attractive, shiny appearance, light green-yellow colored skin covered with large greenish mottles which is suitable for both green and black pickling (particularly desirable for green processing).

3 Drawing Sheets

1

Botanical name of the genus and species of the plant claimed: *Olea europaea* L.
Variety denomination: ‘SEPOKA’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct variety of an olive tree, botanically described as *Olea europaea* L. of the Oleaceae family, and hereinafter referred to by the variety denomination ‘SEPOKA’.

The new *Olea europaea* ‘SEPOKA’ is a product of a planned breeding program conducted by the inventors, Shimon Lavee, Benjamin Avidan and Yair Manni, in Bet Dagan, Israel. The objective of the breeding program was to develop a new *Olea europaea* L. variety which is suitable for both green and black pickling, exhibits resistance to *Spilocaea oleagina* (peacock eye leaf disease), requires minimal pruning and facilitates hand-picking.

The new *Olea europaea* ‘SEPOKA’ originated as a result of self pollination of the *Olea europaea* L. ‘KADESH’ (unpatented) made by the inventors in 1990 in Bet Dagan, Israel. The new *Olea europaea* L. ‘SEPOKA’ was observed and selected from the progeny of the stated self-pollination in 1994 by the inventors in a controlled in Bet Dagan, Israel.

Asexual propagation of the new *Olea europaea* L. ‘SEPOKA’ by rooting of vegetative cuttings was first performed in the spring of 1995 in Bet Dagan, Israel, and has demonstrated that the combination of characteristics as herein disclosed for the new variety are firmly fixed and

2

retained through successive generations of asexual reproduction. The new variety propagates true to type.

BRIEF DESCRIPTION OF THE INVENTION

The following traits have been repeatedly observed and are determined to be characteristics of ‘SEPOKA’ which in combination distinguish this olive tree as a new, unique and distinct variety:

1. A small to medium-sized tree facilitating hand picking;
2. Suitable for intensive orchard growing with harvesting from the ground level with only minimal pruning required;
3. Good resistance to *Spilocaea oleagina* L.; and
4. Elongated shaped fruit, with attractive, shiny appearance, light green-yellow colored skin with large greenish mottles which is suitable for both green and black pickling (particularly desirable for green processing).

In comparison to the unpatented, parental variety *Olea europaea* L. ‘KADESH’, the new *Olea europaea* L. ‘SEPOKA’ differs primarily in the traits listed in Table 1.

TABLE 1

Comparison with parent variety.		
Trait	New Variety ‘SEPOKA’	Parent ‘KADESH’ (unpatented)
Trunk circumference (measured at 50 cm	About 40 cm	About 42 cm

TABLE 1-continued

Comparison with parent variety.		
Trait	New Variety 'SEPOKA'	Parent 'KADESH' (unpatented)
above ground)		
Tree Habit	Semi-spreading	Round
Tree Height	Upto about 3.0 m	Upto about 4.0 m

Of the many commercial varieties known to the present inventors, the most similar in comparison to *Olea europaea* L. 'SEPOKA' is *Olea europaea* L. 'BARNEA' (unpatented). 'SEPOKA' differs from 'BARNEA' in the traits described in Table 2:

TABLE 2

Comparison with a well known commercial variety.		
Trait	New Variety 'SEPOKA'	Comparison variety 'BARNEA'
Tree type	Semi-dwarf	Normal
Conspicuousness of marbling	Strong	Weak
Time of ripening	Very late	Medium

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs illustrate the overall appearance of the new *Olea europaea* L. 'SEPOKA' showing the colors as true as is reasonably possible with colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed morphological description, which accurately describe the color of 'SEPOKA'.

FIG. 1 shows a side view perspective of a typical 10-year old specimen of 'SEPOKA' exhibiting its overall appearance.

FIG. 2 shows a close-up view of typical unripe, fully grown fruit of 'SEPOKA'.

FIG. 3 shows a close-up view of typical ripe, fully grown fruit of 'SEPOKA' and it's flesh.

DETAILED MORPHOLOGICAL DESCRIPTION

The new *Olea europaea* L. 'SEPOKA' has not been observed under all possible environmental conditions. The phenotype of the new variety may vary with variations in environment such as temperature, light intensity, and day length without any change in the genotype of the olive tree.

The aforementioned photographs, together with the following observations, measurements and values describe trees of 'SEPOKA' as grown in the olive farm in Bet Dagan, Israel, under conditions which closely approximate those generally used in commercial practice in Israel. Trees of 'SEPOKA' are planted at a distance of 4x7 meters in sandy red loam soil at an elevation of about 30 meters above sea level. Trees of 'SEPOKA' are irrigated by drip system (about 4 liters per hour) during the summer. Average annual rainfall is about 550 mm, with an average 350 mm of rainfall in winter (December to February). NPK fertilization (9:3:9.50 ppm) is administered through the drip system. Mean diurnal minimum temperature in January is 7.2° C., and mean diurnal maximum temperature in July is 30.8° C.

Unless otherwise stated, the detailed morphological description includes observations, measurements and values based on ten-year-old 'SEPOKA' trees grown in the olive farm in Bet Dagan, Israel from 2004 to 2007. Quantitative data are expressed as an average of measurements taken from 10 parts of trees of 'SEPOKA'. The measurements of any individual tree, or any group of trees, of the new variety may vary from the stated average.

Color references are made to The Royal Horticultural Society Colour Chart (R.H.S.), 1986 edition, except where general colors of ordinary significance are used. Color values were taken under conditions of full sunlight in Bet Dagan, Israel.

All of the trees of 'SEPOKA', insofar as they have been observed, have been consistent in all the characteristics described below.

Classification:

Botanical.—*Olea europaea* L.

Parentage: Self pollination of *Olea europaea* L. 'KADESH' (unpatented)

Propagation:

Method.—Rooting of cuttings.

Growing conditions:

Light intensities.—Full sunlight.

Temperature.—Mean diurnal minimum temperature in January is 7.2° C., and mean diurnal maximum temperature in July is 30.8° C.

Fertilization.—NPK fertilization (9:3:9.50 ppm) is administered through the drip system.

Growth regulators.—No growth regulators are used.

Pruning or trimming requirements.—The olive tree of 'SEPOKA' as described is grown without training. When the trees are freely grown, the size and the shape assumed by the plants are not typical of *Olea europaea* L. species.

Plant.—Type: Semi-dwarf. Growth habit: semi-spreading. Vigor: strong. Height: up to about 3 m. Diameter (Spread of Canopy: about 2.8 m. Circumference of trunk (at 50 cm height): about 40 cm. Attitude of branches: erect. Density of canopy: dense. Aspect of bark: semi-rough. Bark color: gray RHS 198 A. Lenticles: not visible. Abnormal leaves: present. Shape of abnormal leaves: falsiform.

Main branch.—Number of main branches per tree: 3 or 4 resulting from pruning. Length: the length of the branch is up to 3 meters. Diameter: 10-15 cm. Color: gray RHS 198 B. Angle of main brunches with axis of the tree: about 30°. Surface: slightly rough. Lenticles: not visible.

One year old Shoot.—Shape in cross section: round. Diameter: 5-9 mm. Color: gray RHS 156 A. Surface: smooth. Lenticles: present. Density of lenticles: sparse. Shape of lenticel: round. Size of lenticle: minuscule. Color of lenticel: gray between RHS 196A and RHS 196 B.

Fruiting shoot.—Color: light grey. Length of node: medium 22-30 mm. Feathers (side branching of shoot): medium.

Leaves.—Arrangement: the arrangement of the leaves is typical of *Olea europaea* L. species (two opposite leaves per each node).

Leaf blade.—Size: very small to small. Ratio length/width: medium. Length: 48-58 mm. Width: 8-11 mm. Shape: elliptic. Glossiness: present. Color of upper side: medium green RHS 137 C. Color of lower side:

grey-green RHS 148 D. Curvature of longitudinal axis of blade: flat. Twisting: present. Margin: entire. Undulation of margin: very weak. Shape of apex: acute. Shape of base: acute. Upper surface: smooth. Lower surface: smooth. Pubescence of upper side: 5 glabrous. Pubescence of lower side: glabrous. Color of main vein of upper side: dark green RHS 146 B. Color of main vein of lower side: dark green RHS 146 B.

Petiole.—Length: 5-7 mm. Diameter: about 1 mm. 10 Color: grey-green RHS 148 D.

Inflorescence.—Structure: raceme. Shape: elongated. Branching: strong. Axillary flowers: absent. Length: varies up to 15 cm. Diameter: varies up to 5 cm. Number of flowers per inflorescence: numerous. 15 Fruits per inflorescence: 1.

Flower bud.—Shape: obovoid. Size (just before opening): minuscule. Color: greenish between RHS 145 C and RHS 145 D. Pubescence: strong. Bud length: about 4 mm. Bud diameter: about 3 mm. 20

Flower.—Diameter: 5-7 mm. Color: white RHS 155 A. Color of pollen: yellow RHS 3A. Fragrance: mild.

Corolla segment.—Number per flower: 4. Length: about 3 mm. Width: about 2 mm. Shape: elliptic. Shape of apex: rounded. Base: fused. Color of upper side: 25 white RHS 155 A. Color of lower side: white RHS 155 A.

Calyx lobe.—Number per lower: 4. Shape: funnel. Length: about 1 mm. Width: less than 1 mm. Shape of apex: acute. Base: fused. Margin: entire. Color of 30 upper side: typical. Color of lower side: typical.

*Pedice*l.—Length: 1-2 mm. Diameter: less than 1 mm. Surface: smooth. Color: very light green RHS 145 D.

Fruit.—

Fruit (drupe).—In the following description, Position A 35 refers to the position in which the fruit shows its largest asymmetry. Position B can be reached from position A by turning 90 degrees along the longitudinal axes in a way that presents the most developed part of the fruit to the observer (according to UPOV rules). 40 Maturity when described: ripe for eating. Size: medium. Weight: 4-6.2 g. Length: 22-26 mm. Diameter: 13-15 mm. Length/diameter ratio: 2.5. Shape: elongated. Transversal section shape: circular. Green color: RHS 144 A. Color when physiologically ripe: 45 light violet RHS 60 B. Color of flash (freshly cut): beige RHS 159 A. Conspicuousness of marble: strong. Size of mottles: medium. Color of mottles: greenish RHS 145 C. Symmetry in position A: strongly asymmetric. Symmetry in position B: 50 weakly asymmetric. Position of maximum diameter:

central. Shape of apex in position A: rounded. Shape of apex in position B: rounded. Mucron: absent. Position of pistil scar: not central. Shape of base in position A: truncate. Shape of base in position B: rounded. Percentage of stone: 10.5-16.1. Pulp/stone ratio: about 7.04. Pulp/stone detachment: free stone. Conspicuousness of suture: inconspicuous. Fruit suture curvature: not relevant. Oil content for fresh matter%-Percent of oil in mesocarp (Analyzed at 50% black fruit): 17.1. Percentage of dry weight: 27.4. Production per tree: about 28 kg. Oleic acid: not recorded. Polyphenol content: not recorded. Organoleptic characteristics: not recorded.

Stalk.—Length: 13-21 mm. Thickness: about 1 mm. Color: light yellow green RHS 145 C. Width of stalk cavity: narrow, about 4-5 mm. Shape of stalk cavity: circular. Depth of stalk cavity: shallow, about 2 mm. Shape of cross section: circular.

Stone.—Shape in position A: elongated. Shape in position B: elongated. Symmetry in position A: weakly asymmetric. Symmetry in position B: symmetrical. Shape of cross section: circular. Position of largest cross section: central. Grooving: weak. Distribution of grooves: including apex. Number of grooves on basal end: more than 10. Distribution of grooves on basal end: irregular. Shape of distal end in position A: pointed. Shape of distal end in position B: pointed. Mucron: present. Shape of base in position A: pointed. Shape of base in position B: truncate. Conspicuousness of suture: absent. Curvature of suture: not relevant. Size: medium. Length: 17-21 mm. Diameter: 7-8 mm. Color when dry: beige RHS 164 D. Weight: about 0.625 g.

Time of flowering.—Mid April in Bet Dagan, Israel.

Flowering period.—About 10 days in Bet Dagan Israel.

Time of ripening (green maturation).—End of August in Bet Dagan Israel.

Period of ripening.—About 2 Weeks in Bet Dagan Israel.

Resistance to abiotic factors:

Cold.—Observed resistance of down to 0° C. in Bet Dagan, Israel

Resistance to parasites:

Spilocaea oleagina.—‘SEPOKA’ exhibits resistance.

Pseudomonas savastanoi.—‘SEPOKA’ has not been tested.

What is claimed is:

1. A new and distinct *Olea europaea* L. olive tree variety named ‘SEPOKA’, as illustrated and described herein.

* * * * *

FIG. 1




FIG. 2



FIG. 3

