



US00PP22347P2

(12) **United States Plant Patent**  
**Heuger**(10) **Patent No.:** US PP22,347 P2  
(45) **Date of Patent:** Dec. 13, 2011

- (54) **HELLEBORUS PLANT NAMED 'COSEH 820'**
- (50) Latin Name: *Helleborus niger* × *Helleborus argutifolius*  
Varietal Denomination: **COSEH 820**
- (76) Inventor: **Josef Heuger**, Glandorf (DE)
- (\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **13/068,370**
- (22) Filed: **May 9, 2011**
- (51) **Int. Cl.**  
**A01H 5/00** (2006.01)
- (52) **U.S. Cl.** ..... **Plt./439**
- (58) **Field of Classification Search** ..... Plt./439  
See application file for complete search history.

- (56) **References Cited**
- U.S. PATENT DOCUMENTS
- PP21,003 P2 \* 5/2010 Heuger ..... Plt./439  
PP21,504 P2 \* 11/2010 Heuger ..... Plt./439

\* cited by examiner

*Primary Examiner* — Wendy C Haas  
(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**  
A new and distinct cultivar of *Helleborus* plant named 'COSEH 820', characterized by its upright and uniformly mounded plant habit; uniform and freely flowering habit; dark green-colored leaves; long flowering period; single light yellow green and white-colored flowers; and good garden performance.

**2 Drawing Sheets****1**

Botanical designation: *Helleborus niger* × *Helleborus argutifolius*.  
Cultivar denomination: 'COSEH 820'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of *Helleborus* plant, botanically known as *Helleborus niger* × *Helleborus argutifolius*, and hereinafter referred to by the name 'COSEH 820'.

The new *Helleborus* plant is a product of a planned breeding program in Glandorf, Germany. The objective of the breeding program was to create new uniform *Helleborus* plants with unique and attractive plant habit, leaf and flower coloration and resistance to pests.

The new *Helleborus* plant originated from a cross-pollination conducted by the Inventor in Glandorf, Germany in November, 2003 of an unnamed proprietary seedling selection of *Helleborus niger*, not patented, as the female, or seed, parent with an unnamed proprietary seedling selection of *Helleborus argutifolius*, not patented, as the male, or pollen, parent. The new *Helleborus* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination grown in a controlled greenhouse environment in Glandorf, Germany in February, 2006.

Asexual reproduction of the new *Helleborus* plant by divisions in a controlled greenhouse environment in Glandorf, Germany since March, 2006, has shown that the unique features of this new *Helleborus* plant are stable and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

Plants of the new *Helleborus* have not been observed under all possible environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

**2**

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'COSEH 820'. These characteristics in combination distinguish 'COSEH 820' as a new and distinct *Helleborus* plant:

1. Upright and uniformly mounded plant habit.
2. Uniform and freely flowering habit.
3. Dark green-colored leaves.
4. Long flowering period.
5. Single light yellow green and white-colored flowers.
6. Good garden performance.

10 Compared to plants of the parent selections, plants of the new *Helleborus* are more uniformly mounded and are more uniformly and freely flowering.

15 Plants of the new *Helleborus* can be compared to plants of *Helleborus niger* × *Helleborus lividus* 'COSEH 700', disclosed in U.S. Plant Pat. No. 21,003. In side-by-side comparisons conducted in Glandorf, Germany, plants of the new *Helleborus* differed primarily from plants of 'COSEH 700' in leaf color as plants of the new *Helleborus* had lighter green-colored leaves than plants of 'COSEH 700'. In addition, plants of the new *Helleborus* and 'COSEH 700' differed slightly in flower color.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

20 The accompanying colored photographs illustrate the overall appearance of the new *Helleborus* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Helleborus* plant.

25 The photograph on the first sheet comprises a side perspective view of a typical flowering plant of 'COSEH 820' grown in a container.

30 The photograph on the second sheet is a close-up view of a typical flowering plant of 'COSEH 820'.

**DETAILED BOTANICAL DESCRIPTION**

35 The aforementioned photographs and following observations, measurements and values describe plants grown in

15-cm containers during the winter in a glass-covered greenhouse in Glandorf, Germany and under conditions which closely approximate commercial *Helleborus* production. During the production of the plants, day temperatures ranged from 12° C. to 32° C. and night temperatures ranged from 3° C. to 18° C. Plants were 14 months old when the photographs and the description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Helleborus niger* × *Helleborus argutifolius* 'COSEH 820'.

Parentage:

*Female, or seed, parent.*—Unnamed proprietary seedling selection of *Helleborus niger*, not patented.

*Male, or pollen, parent.*—Unnamed proprietary seedling selection of *Helleborus argutifolius*, not patented.

Propagation:

*Type.*—By tissue culture.

*Time to initiate roots.*—About two months at 12° C.

*Time to produce a rooted young plant.*—About six months at 4° C. to 15° C.

*Root description.*—Thick to thin, fleshy; white to brown in color.

*Rooting habit.*—Sparse.

Plant description:

*Plant form and growth habit.*—Herbaceous perennial; upright and uniformly mounding plant habit with flowers held just above the foliar plane; plant shape is globular; and moderately vigorous growth habit.

*Plant height.*—About 28.7 cm.

*Plant diameter (area of spread).*—About 31.3 cm.

Foliage description:

*Arrangement.*—Leaves arranged in a basal rosette; leaves palmately compound with five leaflets per leaf.

*Leaf length.*—About 15.2 cm.

*Leaf width.*—About 17.9 cm.

*Leaflet length.*—About 10.4 cm.

*Leaflet width.*—About 4.8 cm.

*Leaf shape.*—Palmate; orbicular in outline.

*Leaflet shape.*—Elliptic to broadly ovate.

*Leaflet apex.*—Acute.

*Leaflet base.*—Attenuate.

*Leaflet margin.*—Sharply serrate.

*Leaflet texture, upper and lower surfaces.*—Smooth, glabrous; leathery.

*Leaflet venation pattern.*—Pinnate.

*Leaflet color.*—Developing leaves, upper surface: Close to 137A. Developing leaves, lower surface: Close to 146B. Fully developed leaves, upper surface: Between N137A and 147A; venation, close to 144A to 144B. Fully developed leaves, lower surface: Close to 147B; venation, close to 144B to 144C.

*Petiole.*—Length: About 6.7 cm. Diameter: About 5 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 144C.

Flower description:

*Flower shape and habit.*—Single rotate flowers arranged in terminal and axillary cymes; freely and uniform flowering habit with about 75 flowers and flower buds per plant; flowers facing outwardly to slightly nodding and slightly upright.

*Fragrance.*—None detected.

*Natural flowering season.*—Plants begin flowering about ten months after planting; long flowering period; plants flower from late winter to early spring in Germany.

*Flower longevity on the plant.*—About ten days; flowers not persistent.

*Flower buds.*—Length: About 1.9 cm. Diameter: About 1.1 cm. Shape: Ovate. Color: Close to 145C.

*Inflorescence height.*—About 24.2 cm.

*Inflorescence diameter.*—About 22.9 cm.

*Flower diameter.*—About 6.6 cm.

*Flower depth (height).*—About 2.4 cm.

*Petals.*—Transformed into nectaries.

*Sepals.*—Quantity and arrangement: Five arranged in a single whorl. Length: About 3.5 cm. Width: About 2.6 cm. Shape: Broadly ovate to broadly elliptic. Apex: Rounded. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening, upper surface: Close to 145B; towards the margins, close to 145C. When opening, lower surface: Close to 145C. Fully opened, upper surface: Close to 145C to 145D; towards the margins, close to NN155A; towards the base, close to 145B; with development, color becoming closer to 144C. Fully opened, lower surface: Close to 145C; towards the base, close to 145A to 145B.

*Peduncles.*—Strength: Moderately strong. Length: About 13.5 cm. Diameter: About 7 mm. Aspect: About 10° from vertical. Texture: Smooth, glabrous. Color: Close to 145B to 145C heavily dotted with close to 182B.

*Pedicels.*—Strength: Moderately strong. Length: About 2.7 cm. Diameter: About 2 mm. Aspect: Erect to about 30° from the peduncle axis. Texture: Smooth, glabrous. Color: Close to 145A.

*Reproductive organs.*—Stamens: Quantity per flower: About 60. Filament length: About 1.5 cm. Filament color: Close to 157D. Anther shape: Reniform. Anther length: About 2 mm. Anther color: Close to 150B. Pollen amount: Scarce. Pollen color: Close to 158D. Pistils: Quantity per flower: About five. Pistil length: About 1.1 cm. Stigma shape: Club-shaped. Stigma color: Close to 150D. Style length: About 9 mm. Style color: Close to 150D. Ovary color: Close to 145B to 145C. Nectaries (transformed petals): Quantity per flower: About 14. Length: About 8 mm. Diameter, apex: About 2 mm. Diameter, base: About 1 mm. Shape: Flattened triangular. Color: Close to 144B to 144C; towards the apex, close to N144A.

*Seeds/fruits.*—Seed and fruit development have not been observed.

Garden performance: Plants of the new *Helleborus* have been observed to have good garden performance and to tolerate rain and wind. Plants of the new *Helleborus* have been observed to tolerate high temperatures of about 35° C. and are hardy to USDA Hardiness Zone 5.

Pathogen/pest resistance: Plants of the new *Helleborus* have not been shown to be resistant to pathogens and pests common to *Helleborus*.

It is claimed:

1. A new and distinct *Helleborus* plant named 'COSEH 820' as illustrated and described.



