



US00PP22415P2

(12) **United States Plant Patent**
Hurkman

(10) **Patent No.:** **US PP22,415 P2**
(45) **Date of Patent:** **Dec. 27, 2011**

(54) **PETUNIA PLANT NAMED ‘BALSUNMIBU’**

(50) Latin Name: *Petunia*×*hybrida*
Varietal Denomination: **Balsunmibu**

(75) Inventor: **Margaret M. Hurkman**, Santa Maria,
CA (US)

(73) Assignee: **Ball Horticultural Company**, West
Chicago, IL (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/924,344**

(22) Filed: **Sep. 24, 2010**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./356.16**

(58) **Field of Classification Search** Plt./356.16,
Plt./356

See application file for complete search history.

Primary Examiner — Annette Para

(74) *Attorney, Agent, or Firm* — Audrey Charles

(57) **ABSTRACT**

A new and distinct cultivar of *Petunia* plant named ‘Balsun-
mibu’, characterized by its deep violet-blue colored flowers,
dark green-colored foliage, and moderately vigorous, semi-
upright growth habit, is disclosed.

1 Drawing Sheet

1

Latin name of genus and species of plant claimed: *Petunia*×
hybrida.

Variety denomination: ‘Balsunmibu’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Petunia* plant botanically known as *Petunia*×*hybrida* and
hereinafter referred to by the cultivar name ‘Balsunmibu’.

The new cultivar originated in a controlled breeding pro-
gram in Arroyo Grande, Calif. during June 2006. The objec-
tive of the breeding program was the development of *Petunia*
cultivars with single type flowers, unique flower coloration,
and a moderately vigorous, semi-upright growth habit.

The new *Petunia* cultivar is the result of cross-pollination.
The female (seed) parent of the new cultivar is the proprietary
Petunia×*hybrida* breeding selection designated 3141-1-1, not
patented, characterized by its dark violet-blue colored flow-
ers, dark green-colored foliage, and moderately vigorous,
semi-upright growth habit. The male (pollen) parent of the
new cultivar is the proprietary *Petunia*×*hybrida* breeding
selection designated 2730-2-1, not patented, characterized by
its dark purple-colored flowers, medium green-colored foli-
age, and moderately vigorous, semi-upright growth habit.
The new cultivar was discovered and selected as a single
flowering plant within the progeny of the above stated cross-
pollination during May 2007 in a controlled environment at
Arroyo Grande, Calif.

Asexual reproduction of the new cultivar by terminal stem
cuttings since May 2007 at Arroyo Grande, Calif. and West
Chicago, Ill. has demonstrated that the new cultivar repro-
duces true to type with all of the characteristics, as herein
described, firmly fixed and retained through successive gen-
erations of such asexual propagation.

SUMMARY OF THE INVENTION

The following characteristics of the new cultivar have been
repeatedly observed and can be used to distinguish ‘Balsun-
mibu’ as a new and distinct cultivar of *Petunia* plant:

1. Deep violet-blue colored flowers;
2. Dark green-colored foliage; and
3. Moderately vigorous, semi-upright growth habit.

2

Plants of the new cultivar differ from plants of the female
parent primarily in flower size and leaf size. Plants of the new
cultivar have larger-sized flowers and smaller-sized leaves
than plants of the female parent. Plants of the new cultivar
differ from plants of the male parent primarily in flower color.

Of the many commercially available *Petunia* cultivars, the
most similar in comparison to the new cultivar is Cascadias
Noble Blue, not patented. However, in side by side compari-
sons, plants of the new cultivar differ from plants of Cascadias
Noble Blue in at least the following characteristics:

1. Plants of the new cultivar have a more semi-upright
growth habit, shorter plant height and larger plant width,
than plants of Cascadias Noble Blue; and
2. Plants of the new cultivar have a deeper violet-blue
flower color than plants of Cascadias Noble Blue.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographs show, as nearly true as it is
reasonably possible to make the same in color illustrations of
this type, typical flower and foliage characteristics of the new
cultivar. Colors in the photographs differ slightly from the
color values cited in the detailed description, which accu-
rately describes the colors of ‘Balsunmibu’. The plants were
grown in 4-inch pots for 5 weeks in a greenhouse at West
Chicago, Ill. Plants were given one pinch at transplant.

FIG. 1 illustrates a side view of the overall growth and
flowering habit of ‘Balsunmibu’.

FIG. 2 illustrates a close-up view of an individual flower of
‘Balsunmibu’.

DETAILED BOTANICAL DESCRIPTION

The new cultivar has not been observed under all possible
environmental conditions to date. Accordingly, it is possible
that the phenotype may vary somewhat with variations in the
environment, such as temperature, light intensity, and day
length, without, however, any variance in genotype.

The chart used in the identification of colors described
herein is The R.H.S. Colour Chart of The Royal Horticultural
Society, London, England, 2001 edition, except where gen-
eral color terms of ordinary significance are used. The color

values were determined in June 2010 under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe plants produced from cuttings from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown in West Chicago, Ill. in 4-inch pots for 5 weeks utilizing a soilless growth medium. Plants were given one pinch at transplant. Greenhouse temperatures were maintained at approximately 70° F. to 77° F. (21° C. to 25° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Greenhouse light levels of 2,500 footcandles to 6,000 footcandles were maintained during the day. Measurements and numerical values represent averages of typical plants.

Botanical classification: *Petunia*×*hybrida* cultivar Balsunmibu.

Parentage:

Female parent.—Proprietary *Petunia*×*hybrida* breeding selection designated 3141-1-1, not patented.

Male parent.—Proprietary *Petunia*×*hybrida* breeding selection designated 2730-2-1, not patented.

Propagation:

Type cutting.—Terminal stem.

Time to initiate roots.—Approximately 6 to 9 days.

Time to produce a rooted cutting.—Approximately 21 to 28 days.

Root description.—Fibrous.

Rooting habit.—Freely branching.

Plant description:

Commercial crop time.—Approximately 6 to 8 weeks from a rooted cutting to finish in a 10 cm pot.

Growth habit and general appearance.—Moderately vigorous, semi-upright.

Size.—Height from soil level to top of plant plane: Approximately 12.4 cm. Width: Approximately 32.8 cm.

Branching habit.—Freely branching, pinching improves basal branching. Quantity of main branches per plant: Approximately 5.

Branch.—Strength: Moderate. Length: Approximately 14.6 cm. Diameter: Approximately 4.0 mm. Length of central internode: Approximately 1.2 cm. Texture: Densely glandular pubescent with a mixture of long and short length hairs. Gland color: Colorless. Color of young stem: 144A with a strong overlay of 187A. Color of mature stem: 144A with an overlay of 187A.

Foliage description:

General description.—Quantity of leaves per main branch: Approximately 13. Fragrance: Slight. Form: Simple. Arrangement on flowering stem: Opposite.

Leaves.—Aspect: Perpendicular to acute angle to stem. Shape: Ovate to elliptic. Margin: Entire. Apex: Acute. Base: Attenuate. Venation pattern: Pinnate. Length of mature leaf: Approximately 4.5 cm. Width of mature leaf: Approximately 1.9 cm. Texture of upper surface: Sparsely glandular pubescent. Gland color: Colorless. Texture of lower surface: Moderately glandular pubescent. Gland color: Colorless. Color of upper surface of young foliage: 137C with venation of 144B. Color of lower surface of young and mature foliage: 138B with venation of 144B. Color of upper surface of mature foliage: Darker than 137A with venation of 144B.

Petiole.—Length: Approximately 4.0 mm. Width: Approximately 3.0 mm. Texture: Densely glandular

pubescent with a mixture of long and short length hairs. Gland color: Colorless. Color: 144B.

Flowering description:

Flowering habit.—‘Balsunmibu’ is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

Lastingness of individual flower on the plant.—Approximately 10 to 12 days.

Flower description:

General description.—Type: Simple, salverform. Quantity per plant: Approximately 19. Fragrance: Slight.

Bud.—Rate of opening: Generally takes 2 to 3 days for bud to progress from first color to fully open flower. Quantity per plant: Approximately 8.

Bud just before opening.—Shape: Oblong. Length: Approximately 4.5 cm. Diameter at apex: Approximately 8.0 mm. Diameter at base: Approximately 3.0 mm. Texture: Densely glandular pubescent. Gland color: Colorless. Color of petals: N92D. Color of tube: N92D with venation of N92A.

Corolla.—Diameter: Approximately 6.8 cm.

Petals.—Quantity: 5, fused to form a tube. Shape: Obovate. Appearance: Velvety. Margin: Entire, slightly wavy. Apex: Broadly acute to rounded. Length from tube: Approximately 3.2 cm. Length of free portion: Approximately 1.5 cm. Width: Approximately 3.2 cm. Texture of upper surface: Glabrous. Texture of lower surface: Sparsely glandular pubescent. Color of upper surface when first and fully open: Darker with more purple than N89A and midveins of N186A. Color of lower surface when first and fully open: Closest to 90A with midveins of N92A.

Corolla tube.—Length: Approximately 3.5 cm. Diameter at distal end: Approximately 1.0 cm. Diameter at proximal end: Approximately 3.0 mm. Texture of inner surface: Glabrous. Texture of outer surface: Densely glandular pubescent. Gland color: Colorless. Color of inner surface: Darker than N92A. Color of outer surface: Darker than N92D with venation of N92A.

Sepals.—Quantity per flower: 5, fused at base. Shape: Linear. Apex: Acute. Length: Approximately 1.6 cm. Width: Approximately 2.0 mm. Texture of upper surface: Densely glandular pubescent. Texture of lower surface: Densely glandular pubescent. Color of upper surface: 137A transitioning to 144A at base. Color of lower surface: 137C with an overlay of 187A on upward facing surface.

Peduncle.—Strength: Strong. Aspect: Acute angle to stem. Length: Approximately 2.7 cm. Diameter: Approximately 2.0 mm. Texture: Densely glandular pubescent with a mixture of long and short length hairs. Gland color: Colorless. Color: 144A with a strong overlay of 187A on upward facing surface.

Reproductive organs.—Androecium: Stamen quantity: 5, basifixed. Stamen length: Approximately 2.5 cm. Filament length of fixed portion: Approximately 9.0 mm. Filament color: 155D with an overlay of 86A. Anther shape: Bilobed. Anther length: Approximately 1.0 mm. Anther color: 158A with an overlay of 86A. Pollen amount: Abundant. Pollen color: 158D. Gynoecium: Pistil quantity: 1 per flower. Pistil length: Approximately 2.7 cm. Stigma shape: Funnel. Stigma length: Approximately 2.0 mm. Stigma color: 146A

with 86A at base. Style length: Approximately 2.2 cm.
Style color: 145B with an overlay of 86A at apex.
Ovary length: Approximately 3.0 mm. Ovary color:
144C.
Seed and fruit production: Neither seed nor fruit production
has been observed.

Disease and pest resistance: Resistance to pathogens and
pests common to *Petunia* has not been observed.
What is claimed is:
1. A new and distinct cultivar of *Petunia* plant named
5 ‘Balsunmibu’, substantially as herein shown and described.

* * * * *



FIG. 1



FIG. 2