



US00PP22425P2

(12) **United States Plant Patent**
Heuger(10) **Patent No.:** US PP22,425 P2
(45) **Date of Patent:** Dec. 27, 2011

- (54) **HELLEBORUS PLANT NAMED 'COSEH 800'**
- (50) Latin Name: *Helleborus niger*×*Helleborus lividus*
Varietal Denomination: **COSEH 800**
- (76) Inventor: **Josef Heuger**, Glandorf (DE)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **13/068,372**
- (22) Filed: **May 9, 2011**
- (51) **Int. Cl.**
A01H 5/00 (2006.01)
- (52) **U.S. Cl.** **Plt./439**
- (58) **Field of Classification Search** Plt./439
See application file for complete search history.

- (56) **References Cited**
- U.S. PATENT DOCUMENTS
- PP19,413 P2 * 11/2008 van Paemel Plt./439
PP21,063 P2 * 6/2010 Heuger Plt./439
- * cited by examiner

Primary Examiner — Wendy C Haas
(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**
A new and distinct cultivar of *Helleborus* plant named 'COSEH 800', characterized by its upright and uniformly mounded plant habit; uniform and freely flowering habit; dark green-colored leaves; long flowering period; single yellow green to green white-colored flowers; and good garden performance.

2 Drawing Sheets**1**

Botanical designation: *Helleborus niger*×*Helleborus lividus*.

Cultivar denomination: 'COSEH 800'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Helleborus* plant, botanically known as *Helleborus niger*×*Helleborus lividus*, and hereinafter referred to by the name 'COSEH 800'.

The new *Helleborus* plant is a product of a planned breeding program in Glandorf, Germany. The objective of the breeding program was to create new uniform *Helleborus* plants with unique and attractive plant habit, leaf and flower coloration and resistance to pests.

The new *Helleborus* plant originated from a cross-pollination conducted by the Inventor in Glandorf, Germany in November, 2003 of an unnamed proprietary seedling selection of *Helleborus niger*, not patented, as the female, or seed, parent with an unnamed proprietary seedling selection of *Helleborus lividus*, not patented, as the male, or pollen, parent. The new *Helleborus* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination grown in a controlled greenhouse environment in Glandorf, Germany in February, 2006.

Asexual reproduction of the new *Helleborus* plant by divisions in a controlled greenhouse environment in Glandorf, Germany since March, 2006, has shown that the unique features of this new *Helleborus* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Helleborus* have not been observed under all possible environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

5 The following traits have been repeatedly observed and are determined to be the unique characteristics of 'COSEH 800'. These characteristics in combination distinguish 'COSEH 800' as a new and distinct *Helleborus* plant:

1. Upright and uniformly mounded plant habit.
2. Uniform and freely flowering habit.
3. Dark green-colored leaves.
4. Long flowering period.
5. Single yellow green to green white-colored flowers.
6. Good garden performance.

10 Compared to plants of the parent selections, plants of the new *Helleborus* are more uniformly mounded and are more uniformly and freely flowering.

15 Plants of the new *Helleborus* can be compared to plants of *Helleborus niger*×*Helleborus lividus* 'COSEH 710', disclosed in U.S. Plant Pat. No. 21,063. In side-by-side comparisons conducted in Glandorf, Germany, plants of the new *Helleborus* differed primarily from plants of 'COSEH 710' in flower color as plants of 'COSEH 710' had light green-colored flowers with reddish pink overtones. In addition, plants 20 of the new *Helleborus* 'COSEH 710' differed slightly in leaf color.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

25 The accompanying colored photographs illustrate the overall appearance of the new *Helleborus* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of 30 the new *Helleborus* plant.

The photograph on the first sheet comprises a side perspective view of a typical flowering plant of 'COSEH 800' grown in a container.

35 The photograph on the second sheet is a close-up view of a typical flowering plant of 'COSEH 800'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photographs and following observations, measurements and values describe plants grown in

15-cm containers during the winter in a glass-covered greenhouse in Glandorf, Germany and under conditions which closely approximate commercial *Helleborus* production. During the production of the plants, day temperatures ranged from 12° C. to 32° C. and night temperatures ranged from 3° C. to 18° C. Plants were 14 months old when the photographs and the description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Helleborus niger* × *Helleborus lividus* 'COSEH 800'.

Parentage:

Female, or seed, parent.—Unnamed proprietary seedling selection of *Helleborus niger*, not patented.

Male, or pollen, parent.—Unnamed proprietary seedling selection of *Helleborus lividus*, not patented.

Propagation:

Type.—By tissue culture.

Time to initiate roots.—About two months at 12° C.

Time to produce a rooted young plant.—About six months at 4° C. to 15° C.

Root description.—Thick to thin, fleshy; white to brown in color.

Rooting habit.—Sparse.

Plant description:

Plant form and growth habit.—Herbaceous perennial; upright and uniformly mounding plant habit with flowers held just above the foliar plane; plant shape is globular; and moderately vigorous growth habit.

Plant height.—About 34.7 cm.

Plant diameter (area of spread).—About 37.9 cm.

Foliage description:

Arrangement.—Leaves arranged in a basal rosette; leaves palmately compound with five leaflets per leaf.

Leaf length.—About 13.7 cm.

Leaf width.—About 15 cm.

Leaflet length.—About 9.4 cm.

Leaflet width.—About 4.2 cm.

Leaf shape.—Palmate; orbicular in outline.

Leaflet shape.—Broadly elliptic to obovate.

Leaflet apex.—Acute.

Leaflet base.—Attenuate.

Leaflet margin.—Sharply serrate; slightly undulate.

Leaflet texture, upper and lower surfaces.—Smooth, glabrous; leathery.

Leaflet venation pattern.—Pinnate.

Leaflet color.—Developing leaves, upper surface: Close to 146A. Developing leaves, lower surface: Close to 148B. Fully developed leaves, upper surface: Close to 147A; venation, close to 144B. Fully developed leaves, lower surface: Close to 148B; venation, close to N146C.

Petiole.—Length: About 11.7 cm. Diameter: About 6 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 146C to 146D; heavily dotted with close to 187A and 187D.

Flower description:

Flower shape and habit.—Single rotate flowers arranged in terminal and axillary cymes; freely and uniform flowering habit with about 60 flowers and flower buds per plant; flowers facing outwardly to slightly nodding and slightly upright.

Fragrance.—None detected.

Natural flowering season.—Plants begin flowering about ten months after planting; long flowering period; plants flower from late winter to early spring in Germany.

Flower longevity on the plant.—About ten days; flowers not persistent.

Flower buds.—Length: About 1.8 cm. Diameter: About 1.1 cm. Shape: Ovate. Color: Close to 145C to 145D.

Inflorescence height.—About 31.6 cm.

Inflorescence diameter.—About 12.1 cm.

Flower diameter.—About 6.3 cm.

Flower depth (height).—About 2.5 cm.

Petals.—Transformed into nectaries.

Sepals.—Quantity and arrangement: Five arranged in a single whorl. Length: About 3 cm. Width: About 2.5 cm. Shape: Broadly ovate to broadly elliptic. Apex: Rounded. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening, upper surface: Close to 145C to 145D or 150D. When opening, lower surface: Close to 150D. Fully opened, upper surface: Close to 150D or 157D; center and towards the base, close to 148D; with development, color becoming closer to 144B to 144C. Fully opened, lower surface: Close to 145D moderately to strongly tinged with close to 182D.

Peduncles.—Strength: Moderately strong. Length: About 22.1 cm. Diameter: About 6 mm. Aspect: About 10° from vertical. Texture: Smooth, glabrous. Color: Close to 145B to 145C moderately dotted with close to 182A to 182B.

Pedicels.—Strength: Moderately strong. Length: About 5.1 cm. Diameter: About 2 mm. Aspect: Erect to about 30° from the peduncle axis. Texture: Smooth, glabrous. Color: Close to 150D.

Reproductive organs.—Stamens: Quantity per flower: About 45. Filament length: About 1.7 cm. Filament color: Close to 157D. Anther shape: Reniform. Anther length: About 2 mm. Anther color: Close to 150B. Pollen amount: Scarce. Pollen color: Close to 158D. Pistils: Quantity per flower: About five. Pistil length: About 1.1 cm. Stigma shape: Club-shaped. Stigma color: Close to 157D. Style length: About 9 mm. Style color: Close to 157D. Ovary color: Close to 145D. Nectaries (transformed petals): Quantity per flower: About 14. Length: About 8 mm. Diameter, apex: About 2 mm. Diameter, base: About 1 mm. Shape: Flattened triangular. Color: Close to 144B; towards the apex, close to N144B.

Seeds/fruits.—Seed and fruit development have not been observed.

Garden performance: Plants of the new *Helleborus* have been observed to have good garden performance and to tolerate rain and wind. Plants of the new *Helleborus* have been observed to tolerate high temperatures of about 35° C. and are hardy to USDA Hardiness Zone 5.

Pathogen/pest resistance: Plants of the new *Helleborus* have not been shown to be resistant to pathogens and pests common to *Helleborus*.

It is claimed:

1. A new and distinct *Helleborus* plant named 'COSEH 800' as illustrated and described.



