



US00PP22477P2

**(12) United States Plant Patent
Stemkens****(10) Patent No.: US PP22,477 P2
(45) Date of Patent: Jan. 31, 2012**(54) *VERBENA* PLANT NAMED 'APRICENA'(50) Latin Name: *Verbena*×*hybrida*
Varietal Denomination: **Apricena**(75) Inventor: **Henricus Godefriedus Wilhelmus
Stemkens**, Enkhuizen (NL)(73) Assignee: **Syngenta Crop Protection AG**, Basel
(CH)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.(21) Appl. No.: **12/924,306**(22) Filed: **Sep. 24, 2010**(51) Int. Cl. *A01H 5/00* (2006.01)(52) U.S. Cl. **Plt./308**(58) **Field of Classification Search** Plt./308
See application file for complete search history.*Primary Examiner* — Susan McCormick Ewoldt(74) *Attorney, Agent, or Firm* — S. Matthew Edwards(57) **ABSTRACT**A new *Verbena* plant named 'Apricena' particularly distin-
guished by the light red (apricot) flower color; medium to
dark green foliage, small sized creeping and trailing plant
habit with good branching and is early to flower.**1 Drawing Sheet****1**Latin name of the genus and species of the plant claimed:
Verbena×*hybrida*.

Varietal denomination: 'Apricena'.

BACKGROUND OF THE NEW PLANTThe present invention comprises a new *Verbena*, botani-
cally known as *Verbena*×*hybrida*, and hereinafter referred to
by the variety name 'Apricena'.'Apricena' is a product of a planned breeding program. The
new cultivar has light red (apricot) flower color; medium to
dark green foliage, small sized creeping and trailing plant
habit with good branching and is early to flower.'Apricena' originated from an open pollinated hybridiza-
tion made in the summer of 2003 in a controlled breeding
environment in Enkhuizen, Netherlands. The female parent
was the unpatented, proprietary plant designated 'E0735-4'
with red flowers, less incised foliage and is more vigorous.The male parent of 'Apricena' was an unknown plant. The
resultant seed was sown in February 2004.'Apricena' was selected as one flowering plant within the
progeny of the stated cross in the August 2004 in a controlled
environment in Enkhuizen, Netherlands.The first act of asexual reproduction of 'Apricena' was
accomplished when vegetative cuttings were propagated
from the initial selection in August 2004 in a controlled envi-
ronment in Enkhuizen, Netherlands.**BRIEF SUMMARY OF INVENTION**Horticultural examination of plants grown from cuttings of
the plant initiated in August 2004, and continuing thereafter,
has demonstrated that the combination of characteristics as
herein disclosed for 'Apricena' are firmly fixed and are
retained through successive generations of asexual reproduc-
tion.'Apricena' has not been observed under all possible envi-
ronmental conditions. The phenotype may vary significantly
with variations in environment such as temperature, light
intensity and day length.A Plant Breeder's Right for this cultivar was applied for in
Canada on Feb. 9, 2010 (10-6830). 'Apricena' has not been
made publicly available more than one year prior to the filing
of this application.**2**The following traits have been repeatedly observed and are
determined to be basic characteristics of the new variety. The
combination of these characteristics distinguishes this *Ver-
bena* as a new and distinct variety.**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**The accompanying photographic drawing shows typical
flower and foliage characteristics of 'Apricena' with colors
being as true as possible with an illustration of this type. The
photographic drawing shows a flowering potted plant of the
new variety and a close-up of the flowers.**DETAILED BOTANICAL DESCRIPTION**The aforementioned photographs were taken in April 2010
from plants growing in a greenhouse trial in Gilroy, Calif.
USA. These plants were growing in 4 inch pots and were
approximately 15-16 weeks of age.The plant descriptions and measurements were taken in
late May 2010 in Enkhuizen, Netherlands on about 12 week
old plants that had been planted into 12 cm pots and were
grown on benches in a greenhouse at a minimum temperature
(heating temperature) of 12° C. From April on, a minimum
night temperature of 8° C. was held, while day temperatures
were much higher, depending on outdoor temperature.Color references are made to The Royal Horticultural Soci-
ety Colour Chart (R.H.S.) 2001.**TABLE 1**

DIFFERENCES BETWEEN THE NEW VARIETY 'APRICENA' AND A SIMILAR VARIETY		
	'Apricena'	'Comivpec' (U.S. Plant Pat. No. 15,430)
Plant type:	More trailing	Upright
Flower type:	Single	Double
Flower color:	RHS 49C	RHS 27A

Plant:

Form, growth and habit.—Herbaceous, initially spread-
ing, later becoming decumbent to trailing.*Plant height.*—About 7 cm.

- Plant height (inflorescence included).*—About 9-11 cm.
Plant width.—About 35 cm.
- Roots:
Number of days to initiate and produce roots.—About 10-14 days at about 22 degrees C.
Type.—Fine, fibrous, free branching.
Color.—RHS N155B.
- Foliage:
Arrangement.—Alternate, simple.
Immature, leaf color, upper surface.—RHS 137C.
Lower surface.—Between RHS 138B and RHS 138C.
Mature, leaf color, upper surface.—RHS 137A.
Lower surface.—RHS 138A to RHS 138B.
Length.—5.0-6.2 cm.
Width.—2.4-3.3 cm.
Shape.—Ovate.
Base shape.—Acute to attenuate.
Apex shape.—Obtuse.
Margin.—Pinnatifid; entire, with incisions of about 0.7-1.4 cm.
Texture, upper surface.—Slightly glossy, with sparse, short hair.
Lower surface.—Short, stiff hair mainly along the veins; some glandular hairs.
Color of veins, upper surface.—RHS 144B becoming indistinct.
Color of veins, lower surface.—RHS 145B.
Petiole color.—RHS 145B; with RHS 145C on the under side.
Length.—0.1-0.4 cm.
Diameter.—0.1-0.3 cm.
Texture.—Hirtellous.
- Stem:
Characteristics.—Side branches develop potentially at every node.
Color of stem.—RHS 137C.
Length of stem.—Approximately 15-22 cm.
Diameter.—0.4 cm.
Length of internodes.—Approximately 4-5 cm.
Texture.—Rough; densely covered with hair; hirsute.
Color of peduncle.—RHS 138B.
Length of peduncle.—Most often 4-7 cm.
Peduncle diameter.—1.5-2.1 cm.
Texture.—Rough, densely covered with short glandular hair; hirtellous.
- Inflorescence:
Type.—Umbel-like, but actually a spike, umbrella-shaped to roughly semi-spherical.
Blooming habit.—Flowers continuously.
Quantity of inflorescences per plant.—17-21.
Lastingness of individual blooms on the plant.—About 4 days in the greenhouse.
Fragrance.—None.
Inflorescence horizontal diameter.—3.9-4.8 cm.
Inflorescence depth (height).—2.2-3.2 cm.
Quantity of flowers and buds per inflorescence.—24-36.
- Bud (just when starting to show color):
Color.—RHS N155C.
Length.—2.1-2.4 cm.

- Width.*—0.3-0.5 cm.
Shape.—Mainly tube-shaped with a bulbous end.
- Floret:
Form and type.—Sessile; salverform, composed of 5 partly fused petals with a base of a narrow tube.
Immature color, upper surface.—RHS 34C.
Lower surface.—RHS 34D.
Mature color, upper surface.—RHS 49C.
Lower surface.—RHS 49D.
Floret diameter.—1.6-2.2 cm.
Floret depth.—1.6-2.3 cm.
Length of petals.—0.7-0.9 cm.
Width of petals.—0.6-0.8 cm.
Petal shape.—Obovate.
Apex shape.—Emarginate.
Margin.—Entire.
Petal texture, upper surface.—Smooth and glabrous.
Lower surface.—Smooth and glabrous.
Corolla tube length.—1.7-2.1 cm.
Corolla tube color inside.—RHS 150C.
Tube color outside.—RHS 150D or sometimes RHS 155D.
Corolla texture, inside.—Glabrous.
Outside.—Glabrous basally; glandular hairs at the flare.
- Calyx:
Type.—Five sepals whose margins are fused to each other along their length, with a transparent membrane of less than 0.1 cm in width and with one smaller sepal attached to the base of the calyx.
Color of sepals.—RHS 143B to RHS 143C.
Length of sepals.—1.0-1.2 cm.
Width of sepals.—0.2-0.3 cm.
Sepal shape.—Linear.
Apex shape.—Acute.
Margins.—Entire.
Texture.—Densely covered with short hair.
- Reproductive organs:
Pistil.—1.
Length.—1.6-2.2 cm.
Style color.—RHS 144C.
Style length.—1.3-1.6 cm.
Stigma color.—RHS 143B.
Stamens.—Anthers and filaments fused to upper half of corolla tube; four anthers with two pollen sacs per anther.
Color of filaments.—RHS 150D to RHS 155B.
Length filaments.—0.1 cm.
Anther color.—RHS N144B.
Length of anthers.—0.1-0.15 cm.
Color of pollen.—RHS 1B.
Pollen amount.—Moderate.
Fertility/seed set.—Has not been observed on this hybrid.
- Disease/pest resistance: Disease/pest resistance has not been observed on this hybrid.
 What is claimed is:
 1. A new and distinct variety of *Verbena* plant named 'Apricena' substantially as illustrated and described herein.

