



(12) **United States Plant Patent**
Kennedy

(10) **Patent No.:** **US PP22,674 P2**
(45) **Date of Patent:** **Apr. 17, 2012**

(54) **PRIMULA PLANT NAMED ‘K 74’**

(50) Latin Name: *Primula acaulis*
Varietal Denomination: **K 74**

(76) Inventor: **Joseph Olivier Kennedy**, Ballycastle
(IE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/932,944**

(22) Filed: **Mar. 10, 2011**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./472**

(58) **Field of Classification Search** **Plt./472**
See application file for complete search history.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — Penny J. Aguirre

(57) **ABSTRACT**

A new cultivar of *Primula* plant, ‘K 74’, characterized by its
foliage that is a blend of dark green and bronze, its large
flowers that are white in color, its relatively vigorous growth
habit, and its healthy foliage that has been observed to be
disease resistant.

2 Drawing Sheets

1

Botanical classification: *Primula acaulis*.
Variety denomination: ‘K 74’.

**CROSS REFERENCE TO A RELATED
APPLICATION**

This application is co-pending with a U.S. Plant Patent
Application filed for a plant derived from the same cross in the
Inventor’s breeding program that is entitled *Primula* Plant
Named ‘K 72’ (U.S. Plant patent application Ser. No. 12,932,
961).

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Primula* plant, botanically known as *Primula* ‘K 74’ and
will be referred to hereinafter by its cultivar name, ‘K 74’. The
new cultivar of *Primula* is a hardy herbaceous perennial
grown for landscape use.

The new cultivar was derived from a controlled breeding
program conducted by the Inventor at his nursery in Bally-
castle, Northern Ireland. The overall purpose of the breeding
program that was established in the 1980s is to make selec-
tions of *Primula* plants with unique and improved character-
istics to plants from the ‘Garryard’ family of primroses, a
group of *Primula* cultivars that emerged in the 1950s and had
the common characteristic of having bronze colored foliage.

‘K 74’ was selected in the Inventor’s greenhouse in 2006 as
a single unique plant from amongst the seedlings derived
from a cross made in 2003 between two unnamed plants of
Primula acaulis from the Inventor’s breeding program as the
parents.

Asexual reproduction of the new cultivar was first accom-
plished by in vitro propagation in 2009 under the direction of
the Inventor in Ballycastle, Northern Ireland. Asexual propa-
gation has determined the characteristics to be stable and
reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are
determined to be the characteristics of the new cultivar. These
attributes in combination distinguish ‘K 74’ as a unique cul-
tivar of *Primula*.

2

1. ‘K 74’ exhibits foliage that is a blend of dark green and
bronze.
2. ‘K 74’ exhibits large flowers that are white in color.
3. ‘K 74’ exhibits a vigorous growth habit for a dark foli-
aged cultivar of *Primula*.
4. ‘K 74’ exhibits healthy foliage that has been observed to
be disease resistant.

In comparison to its parent plants, ‘K 74’ is more vigorous
and has better bronzing to the foliage and has flowers that are
larger and clearer white in color. ‘K 74’ can be most closely
compared to the cultivar ‘K 72’ from the same breeding
program. ‘K 72’ is similar to ‘K 74’ in having good vigor and
large flowers, however ‘K 72’ differs in having slightly more
bronzing to the foliage and in having red flowers. ‘K 74’ can
also be compared to the cultivar ‘Garryard’ (syns. ‘Garryard
Guinever’, ‘Guinever’), not patented. ‘Garryard’ differs from
‘K 74’ in having foliage that is less bronze in color and in
having flowers that are smaller in size and pink in color.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying colored photographs illustrate the over-
all appearance and distinct characteristics of the new
Primula. The plant in the photograph is 6 months in age as
grown in a 10.5-cm container in a cold-storey greenhouse in
The Netherlands.

The photograph in FIG. 1 provides a side view of ‘K 74’ in
bloom.

The photograph in FIG. 2 provides a close-up view of a
flower of ‘K 74’.

The photograph in FIG. 3 provides a close-up view of the
foliage of ‘K 74’.

The colors in the photograph may differ slightly from the
color values cited in the detailed botanical description, which
accurately describe the colors of the new *Primula*.

DETAILED BOTANICAL DESCRIPTION

The following is a detailed description of six month-old
plants of the new cultivar as grown in 10.5-cm containers in a
cold-storey greenhouse in The Netherlands. The phenotype
of the new cultivar may vary with variations in environmental,
climatic, and cultural conditions, as it has not been tested

under all possible environmental conditions. The color determination is in accordance with The 2007 R.H.S. Colour Chart of The Royal Horticultural Society, London, England, except where general color terms of ordinary dictionary significance are used.

General description:

Blooming period.—Typically March to May in Ballycastle, The Netherlands.

Plant habit.—Herbaceous perennial, clump-forming, tight rosette.

Height and spread.—Reaches about 20 cm in height and about 35 cm in width.

Cold hardiness.—Observed to be hardy to -10° C. in western Europe.

Diseases.—Observed to be disease resistant under the conditions tested.

Root description.—Fibrous roots.

Branching habit.—Basal rosettes.

Propagation.—Tissue culture preferred.

Growth habit.—Vigorous for a bronzed foliated cultivar of *Primula*.

Foliage description:

Leaf shape.—Oblanceolate.

Leaf division.—Simple.

Leaf base.—Broad cuneate.

Leaf apex.—Rounded.

Leaf venation.—Pinnate, on upper surface; 148B to 148C in color fading towards the base into 182B to 182C, lower surface; 148D in color fading towards the base into 182D, sparsely pubescent on main veins, secondary veins and tertiary veins with thin and soft hairs, average length 0.75 mm, 156D to 157D in color.

Leaf margins.—Crenulate, finely sinuate.

Leaf arrangement.—Rosette.

Leaf surface.—Rugose to bullate and moderately glossy, finely pubescent on lower surface.

Leaf color.—Newly emerged foliage, upper surface; 146A to 147A, newly emerged foliage, lower surface; 148A, mature foliage, upper surface; 147A, mature foliage, lower surface; 148A.

Leaf size.—About 11.4 cm in length and 4.4 cm in width.

Leaf quantity.—Average of 24 per rosette.

Leaf attachment.—Sessile.

Inflorescence description:

Inflorescence type.—Funnel form, lower 50% fused into a tube, rosette formed flowers.

Flower number.—Average of 5 per rosette.

Flower fragrance.—None.

Flower aspect.—Upright to slightly outward.

Flower longevity.—About 10 days, depending on temperature and sunlight exposure.

Flower type.—Single.

Flower size.—Average of 3.5 cm in diameter and 2.8 cm in depth.

Flower buds.—Elliptic in shape, average of 8 flower buds per rosette, average of 1.4 cm in length and 5 mm in diameter, color; 145A to 145B then fading to top and turning into 178A, flower buds ribbed (5 ribs); 184D.

Sepals.—5, Narrow oblong, lower 55% fused, margin entire, average of 2 cm in length and 3 mm in width, acuminate apex, fused base, fused arrangement into a campanulate calyx with upper 45% free, surface is dull, main vein moderately pubescent with short soft hairs, average length of hairs 1 mm, color; N155B, color: young and mature upper surface; 146C, tinged at the tip with 176B to 176C, young lower surface; 145A to 145B, top 178A, main vein 184D, mature lower surface; 148A, top 178A, main vein 184D.

Petals.—An average of 5 per flower, rotate arrangement, spatulate, free upper 50% of the petals orbicular, surface is slightly glossy and slightly velvety, smooth texture, upper part of the tube on the lower surface is sparsely pubescent with very short hairs with an average length of 0.3 mm, 149D to 150D in color, margin sinuate, base cuneate, apex is rounded, average of 3.3 cm in length and 1.8 cm in width, color: opening flowers upper surface; NN155D, veined 73D, base (“eye”) 9A to 12A, tube 150A to 150B, fully opened flowers upper surface; N155C to N155D, veined 73D, base (“eye”) 9A to 12A, tube 150A to 150B, opening and fully opened flowers lower surface; NN155D, veined 73D, tube 150B to 150C.

Peduncle.—Average of 2 mm in diameter and 6 cm in length, moderate strength, 186A to 186B in color, glabrous surface.

Pedicel.—None.

Reproductive organs:

Pistils.—About 1 per flower, average of 1.6 cm in length, club shaped stigma N144A in color, style is an average of 1.5 cm in length and 145C and strongly tinged 170C in color, and ovary is 145A to 145B in color.

Stamens.—Average 5, no filaments present, anthers are basifixed and oblong in shape, average of 2 mm in diameter, and N200A in color, pollen is moderate in quantity and 11A in color.

Fruit.—Fruit and seed production was not observed under the conditions tested.

It is claimed:

1. A new and distinct variety of *Primula* plant named ‘K 74’ as described and illustrated herein.

* * * * *




FIG. 1




FIG. 2



FIG. 3