



US00PP22678P2

(12) **United States Plant Patent**  
**Kivistik**(10) **Patent No.:** US PP22,678 P2  
(45) **Date of Patent:** Apr. 24, 2012(54) **CLEMATIS PLANT NAMED ‘UNAI 007’**(50) Latin Name: *Clematis jackmanii*Varietal Denomination: **UNAI 007**(76) Inventor: **Aili Kivistik**, Kose Parish (EE)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/927,359**(22) Filed: **Nov. 12, 2010**(51) **Int. Cl.**  
**A01H 5/00** (2006.01)(52) **U.S. Cl.** ..... **Plt./228**(58) **Field of Classification Search** ..... Plt./228  
See application file for complete search history.*Primary Examiner* — Kent L Bell(57) **ABSTRACT**

A new and distinct *Clematis* cultivar named ‘UNAI 007’ is disclosed, characterized by good resistance to normal diseases and pests of *Clematis*, continuous flowering from May through September, with flower buds continuously produced from the leaf axils and tepals colored Red-Purple with a Purple-Violet central bar. Plants are exceptionally floriferous and compact, making them especially well suited for container production. The new cultivar is a *Clematis*, suitable for ornamental garden purposes.

**1 Drawing Sheet****1**

Latin name of the genus and species: *Clematis jackmanii*.  
Variety denomination: ‘UNAI 007’.

**BACKGROUND OF THE INVENTION**

The new cultivar is a product of a planned breeding program conducted by the inventor that began in 1978. The objective of the breeding program was to develop *Clematis jackmanii* varieties with abundant, continuous, long season flowering, and plants with a good resistance to diseases and pests typical of *Clematis*. In 1982 the inventor, Aili Kivistik a citizen of Estonia, organized a group of unnamed, unpatented *Clematis jackmanii* varieties for an open pollination breeding project. Seeds originating from this open pollination were planted out, and the new variety was selected in 1991. Specific parents of the new variety are unknown, as the seed was bulk collected, without noting the variety collected from. All work was conducted at a non-commercial nursery in Roogja, Estonia.

In 1993 the inventor first propagated ‘UNAI 007’ by vegetative cuttings at the same nursery in Roogja, Estonia. Subsequent generations have shown that the unique features of this cultivar are stable and reproduced true to type.

**SUMMARY OF THE INVENTION**

The cultivar ‘UNAI 007’ has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature, day length, and light intensity, without, however, any variance in genotype. The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘UNAI 007’. These characteristics in combination distinguish ‘UNAI 007’ as a new and distinct *Clematis* cultivar:

1. Good resistance to diseases and pests common to *Clematis*.
2. Continuous flowering from May through September.
3. Very floriferous plants.
4. Flower buds produced continuously from the leaf axils.
5. Very compact plant habit especially well suited for container production.

**2**

6. Tepals colored Red-Purple with a Purple-Violet central bar.

**COMPARISON TO KNOWN VARIETIES**

Plants of the new cultivar ‘UNAI 007’ are similar to known forms of *Clematis jackmanii* in most horticultural characteristics, however, plants of the new cultivar ‘UNAI 007’ are shorter and more compact than common forms of *Clematis jackmanii* known to the inventor. Additionally, plants of ‘UNAI 007’ flower continuously from May through September, whereas many known forms of *Clematis jackmanii* flower reliably 2 months only.

The new cultivar ‘UNAI 007’ can also be compared to the known variety *Clematis jackmanii* ‘Superba’. Plants are similar in most horticultural characteristics, however, plants of the new cultivar produces flowers with 6 tepals, compared to 4 tepals on *Clematis jackmanii* ‘Superba’. Additionally the new cultivar begins blooming one month earlier, and have a different tepal color.

The new cultivar ‘UNAI 007’ can also be compared to the known variety *Clematis jackmanii* ‘Tie Dye’ U.S. Plant Pat. No. 18,913. Plants are similar in most horticultural characteristics, however, plants of the new cultivar are shorter and more compact than ‘Tie Dye’. Additionally the new cultivar begins blooming one month earlier, and flowers one month later than ‘Tie Dye’.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying photograph in FIG. 1 illustrates in full color a typical flowering plant of ‘UNAI 007’ grown outdoors in Estonia. The plant is less than one year old. The photograph was taken using conventional techniques and although colors may appear different from actual colors due to light reflectance it is as accurate as possible by conventional photographic techniques.

**DETAILED BOTANICAL DESCRIPTION**

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used. The following observations and measurements describe

'UNAI 007' plants grown outdoors in Roogaja, Estonia. Plants are approximately 1 year old, in a 1 liter nursery container. The growing conditions are outdoors and unshaded. Temperatures ranged from approximately -2° C. to 20° C. at night to 5° C. to 35° C. during the day. Measurements and numerical values represent averages of typical plant types.

Botanical classification: *Clematis jackmanii* 'UNAI 007'

#### PROPAGATION

10

##### Time to rooting:

*Initiation*.—Approximately 30 days in the Summer at approximately 25° C.

*Rooted cuttings production*.—Approximately 50 days in the Summer at approximately 25° C.

15

Root description: Fine, fibrous.

#### PLANT

Growth habit: Climbing vine.

20

Height: Approximately 50 to 100 cm, measured from the soil line of the pot.

Plant spread: Approximately 50 cm when trained as a vine. Left untrained, approximately 70 cm.

25

Growth rate: Rapid. During active growing season in Spring, approximately 50 cm per month.

Branching characteristics: Very well branched, one main branch with at least 10 lateral branches.

Length of main stem: Approximately 50 cm.

Diameter of main stem: Approximately 0.2 to 0.5 cm.

30

Main stem color: Near RHS Brown 200B.

Main stem texture: Glabrous.

Internode length: Approximately 10 to 15 cm.

Strength: Strong, flexible, semi-woody.

#### FOLIAGE

35

##### Leaf:

*Type*.—Ternate leaves arranged opposite.

*Description ternate leaves*.—Leaves per lateral Branch: Approximately 36 leaflets. Average Length: Approximately 14 cm. Average Width: Approximately 9 cm. Leaflet Petiole: Average Length: Approximately 1.0 cm. Color: Near RHS Green 136A. Diameter: Approximately 0.2 cm. Texture: Glabrous.

40

Tendrils: Not present.

45

#### FLOWER

##### Bloom period:

*Natural season*.—Continuously from approximately May through September.

50

Flower type and habit: Single, axillary; flowers facing outward.

Flower longevity on plant: Approximately 5 days, after fully opened. Self-cleaning.

55

Quantity of flowers: Very free, continuous flowering. Mature plants have at least 50 open flowers during the flowering season.

##### Flower size:

*Diameter*.—Approximately 7 cm.

*Depth*.—Approximately 1 cm.

60

##### Corolla:

*Tepal arrangement*.—Single overlapping whorl.

*Quantity*.—Approximately 6 per flower.

*Size*.—Length: Approximately 3 cm. Width: Approximately 2 cm.

65

*Shape*.—Nearly spatulate.

*Margin*.—Entire.

*Apex*.—Acute.

*Base*.—Truncate.

*Aspect*.—Slightly concave, undulation throughout petal blade, more pronounced on margin.

*Texture*.—Smooth, velvety.

##### Color:

*Tepals*.—When opening: Upper surface: Near RHS Red-Purple 72A with central bar near Purple-Violet 80B. Lower surface: Near RHS Red-Purple 72A with central bar near Red-Purple 72D. Fully opened: Upper surface: Near RHS Red-Purple 72A with central bar near Purple-Violet 80B. Lower surface: Near RHS Red-Purple 72A with central bar near Red-Purple 72D. Ageing: Upper surface: Near RHS Red-Purple 72C with central bar near Purple-Violet 80C. Lower surface: Near RHS Red-Purple 72A with central bar near Red-Purple 72D.

##### Bud:

*Shape*.—Ovate.

*Length*.—Approximately 2 cm.

*Diameter*.—Approximately 1 cm.

*Color*.—Mainly Near RHS Green 138C.

##### Peduncles:

*Length*.—Approximately 0.6 cm.

*Width*.—Approximately 0.3 cm.

*Angle*.—Average 45°.

*Strength*.—Moderate.

*Color*.—Near RHS Green 138A.

*Fragrance*.—None.

#### REPRODUCTIVE ORGANS

##### Stamens:

*Number*.—Approximately 60 per flower.

*Anther length*.—0.2 cm.

*Anther color*.—Near RHS Purple 79A.

*Anther shape*.—Oblong.

*Pollen quantity*.—Moderate.

*Pollen color*.—Near RHS Yellow-Green 1D.

*Filament length*.—Approximately 0.2 cm.

*Filament aspect*.—Straight to slightly outwardly curved.

*Filament color*.—Near RHS Yellow-Green 1D.

##### Pistil:

*Number*.—Approximately 30 per flower.

*Length*.—Approximately 0.4 cm.

*Style color*.—Near RHS Yellow-Green 149D.

*Style length*.—Approximately 1.5 cm.

*Stigma*.—Shape: Linear. Color: Near RHS Yellow-Green 149C.

#### OTHER CHARACTERISTICS

Seeds: Per flower, typically 30 round seeds, approximately 0.15 diameter. Colored near RHS Green 137B.

Disease/pest resistance: Very good resistance to pathogens and pests common to *Clematis* have been observed.

Temperature tolerance: Observed to tolerate temperatures from approximately -5° C. to at least 35° C.

What is claimed is:

1. A new and distinct cultivar of *Clematis* plant named 'UNAI 007' as herein illustrated and described.



Fig. 1