

(12) **United States Plant Patent**
Kennedy

(10) **Patent No.:** **US PP22,711 P2**
(45) **Date of Patent:** **May 1, 2012**

(54) **PRIMULA PLANT NAMED ‘K 72’**

(50) Latin Name: *Primula acaulis*
Varietal Denomination: **K 72**

(76) Inventor: **Joseph Olivier Kennedy**, Ballycastle
(IE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/932,961**

(22) Filed: **Mar. 10, 2011**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./472**

(58) **Field of Classification Search** **Plt./472**
See application file for complete search history.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — Penny J. Aguirre

(57) **ABSTRACT**

A new cultivar of *Primula* plant, ‘K 72’, characterized by its
foliage that is a blend of dark green and bronze, its large
flowers that are deep red in color, its relatively vigorous
growth habit, and its healthy foliage that has been observed to
be disease resistant.

2 Drawing Sheets

1

Botanical classification: *Primula acaulis*.
Variety denomination: ‘K 72’.

**CROSS REFERENCE TO A RELATED
APPLICATION**

This application is co-pending with a U.S. Plant Patent
Application filed for a plant derived from the same cross in the
Inventor’s breeding program that is entitled *Primula* Plant
Named ‘K 74’ (U.S. Plant patent application Ser. No. 12/932,
944).

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of *Primula* plant, botanically known as *Primula* ‘K 72’ and
will be referred to hereinafter by its cultivar name, ‘K 72’. The
new cultivar of *Primula* is a hardy herbaceous perennial
grown for landscape use.

The new cultivar was derived from a controlled breeding
program conducted by the Inventor at his nursery in Bally-
castle, Northern Ireland. The overall purpose of the breeding
program that was established in the 1980s is to make selec-
tions of *Primula* plants with unique and improved character-
istics to plants from the ‘Garryard’ family of primroses, a
group of *Primula* cultivars that emerged in the 1950s and had
the common characteristic of having bronze colored foliage.

‘K 72’ was selected in the Inventor’s greenhouse in 2006 as
a single unique plant from amongst the seedlings derived
from a cross made in 2003 between two unnamed plants of
Primula acaulis from the Inventor’s breeding program as the
parents.

Asexual reproduction of the new cultivar was first accom-
plished by in vitro propagation in 2009 under the direction of
the Inventor in Ballycastle, Northern Ireland. Asexual propa-
gation has determined the characteristics to be stable and
reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

The following traits have been repeatedly observed and are
determined to be the characteristics of the new cultivar. These
attributes in combination distinguish ‘K 72’ as a unique cul-
tivar of *Primula*.

2

1. ‘K 72’ exhibits foliage that is a blend of dark green and
bronze.
2. ‘K 72’ exhibits large flowers that are deep red in color.
3. ‘K 72’ exhibits a vigorous growth habit for a dark foli-
aged cultivar of *Primula*.
4. ‘K 72’ exhibits healthy foliage that has been observed to
be disease resistant.

In comparison to its parent plants, ‘K 72’ is more vigorous and
has better bronzing to the foliage and has flowers that are
larger and deeper red in color. ‘K 72’ can be most closely
compared to the cultivar ‘K 74’ from the same breeding
program. ‘K 74’ is similar to ‘K 72’ in having good vigor and
large flowers, however ‘K 74’ differs in having slightly less
bronzing to the foliage and in having white flowers. ‘K 72’ can
also be compared to the cultivar ‘Garryard’ (syn. ‘Garryard
Guinever’, ‘Guinever’), not patented. ‘Garryard’ differs from
‘K 72’ in having foliage that is less bronze in color and in
having flowers that are smaller in size and pink in color.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying colored photographs illustrate the over-
all appearance and distinct characteristics of the new
Primula. The plant in the photograph is 6 months in age as
grown in a 9-cm container in a cold-storey greenhouse in The
Netherlands.

The photograph in FIG. 1 provides a side view of ‘K 72’ in
bloom.

The photograph in FIG. 2 provides a close-up view of a
flower of ‘K 72’.

The photograph in FIG. 3 provides a close-up view of the
foliage of ‘K 72’. The colors in the photograph may differ
slightly from the color values cited in the detailed botanical
description, which accurately describe the colors of the new
Primula.

DETAILED BOTANICAL DESCRIPTION

The following is a detailed description of six month-old
plants of the new cultivar as grown in 9-cm containers in a
cold-storey greenhouse in The Netherlands. The phenotype
of the new cultivar may vary with variations in environmental,
climatic, and cultural conditions, as it has not been tested

under all possible environmental conditions. The color determination is in accordance with The 2007 R.H.S. Colour Chart of The Royal Horticultural Society, London, England, except where general color terms of ordinary dictionary significance are used.

General description:

Blooming period.—Typically March to May in Ballycastle, The Netherlands.

Plant habit.—Herbaceous perennial, clump-forming, tight rosette.

Height and spread.—Reaches about 20 cm in height and about 35 cm in width.

Cold hardiness.—Observed to be hardy to -10° C. in western Europe.

Diseases.—Observed to be disease resistant under the conditions tested.

Root description.—Fibrous roots.

Branching habit.—Basal rosettes.

Propagation.—Tissue culture preferred.

Growth habit.—Vigorous for a bronzed foliaged cultivar of *Primula*.

Foliage description:

Leaf shape.—Oblanceolate.

Leaf division.—Simple.

Leaf base.—Broad cuneate.

Leaf apex.—Rounded.

Leaf venation.—Pinnate, on upper surface; 183C to 183D stronger towards base in color, lower surface; 177A to 177B in color, sparsely pubescent on main veins, secondary veins and tertiary veins with thin and soft hairs, average length 0.75 mm, 156D in color.

Leaf margins.—Crenulate, finely sinuate.

Leaf arrangement.—Rosette.

Leaf surface.—Rugose to bullate and moderately glossy.

Leaf color.—Newly emerged foliage and mature foliage, upper surface; 147A to N200A, newly emerged foliage, lower surface; 147B to 148A, mature foliage, lower surface; 146A to 147B.

Leaf size.—About 11.7 cm in length and 4.4 cm in width.

Leaf quantity.—Average of 28 per rosette.

Leaf attachment.—Sessile.

Inflorescence description:

Inflorescence type.—Funnel form, lower 50% fused into a tube, rosette formed flowers.

Flower number.—Average of 2 per rosette.

Flower fragrance.—Fresh acid-sweet, faint.

Flower aspect.—Upright to slightly outward.

Flower longevity.—About 10 days, depending on temperature and sunlight exposure.

Flower type.—Single, polyanthus.

Flower size.—Average of 3.2 cm in diameter and 2.8 cm in depth.

Flower buds.—Narrow elliptic in shape, average of 11 flower buds per rosette, average of 2.1 cm in length and 6 mm in diameter, color; N186C then fading to a lighter color turning into 187A to 187B.

Sepals.—5, Narrow oblong, lower 60% fused, margin entire, average of 2.2 cm in length and 3 mm in width, acuminate apex, fused base, fused into a campanulate calyx with upper 40% free, surface is dull, main vein moderately pubescent with short soft hairs, average length of hairs 0.5 mm, color; N155B, color: young and mature upper surface; 183A to 183B, young and mature lower surface; 183A to 183, striped lighter in color; 184B.

Petals.—An average of 7 per flower, rotate arrangement, spatulate, free upper 55% of the petals orbicular, lower 45% of the petals fused into a tube, surface is very slightly glossy and velvety, smooth texture, margin sinuate, apex is emarginate, average of 2.7 cm in length and 1.8 cm in width, color: opening flowers upper surface; 59A, to 187A, base (“eye”) 17B, tube 178B, fully opened flowers upper surface; 59A to 187A, base (“eye”) 17B, tube 178B, opening and fully opened flowers lower surface; 59A to 187A, tube 177A, base 183A.

Peduncle.—Average of 2 mm in diameter and 5.9 cm in length, moderate strength, 184A to 184B in color, glabrous surface.

Pedicel.—None.

Reproductive organs:

Pistils.—About 1 per flower, average of 8 mm in length, club shaped stigma 187A in color, style is an average of 7 mm in length and 187C in color, and ovary is 153B to 153C in color.

Stamens.—Average 5, no filaments present, anthers are basifixed and oblong in shape, average of 2 mm in diameter, and N200A in color, pollen is moderate in quantity and 11A in color.

Fruit.—Fruit and seed production was not observed under the conditions tested.

It is claimed:

1. A new and distinct variety of *Primula* plant named ‘K 72’ as described and illustrated herein.

* * * * *




FIG. 1



FIG. 2




FIG. 3