



US00PP22714P2

(12) **United States Plant Patent**
Olesen

(10) **Patent No.:** **US PP22,714 P2**
(45) **Date of Patent:** **May 8, 2012**

(54) **HYBRID TEA ROSE PLANT NAMED**
‘POULPMT007’

(50) Latin Name: **Rosa hybrid**
Varietal Denomination: **Poulpmt007**

(75) Inventor: **Mogens Nyegaard Olesen**, Fredensborg
(DK)

(73) Assignee: **Poulsen Roser A/S**, Fredensborg (DK)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/930,439**

(22) Filed: **Jan. 7, 2011**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./133**

(58) **Field of Classification Search** Plt./133
See application file for complete search history.

Primary Examiner — Annette Para

(57) **ABSTRACT**

A new garden rose plant of the hybrid tea class which has abundant, orange blend flowers and attractive foliage. This new and distinct variety has shown to be uniform and stable in the resulting generations from asexual propagation.

2 Drawing Sheets

1

Botanical designation: *Rosa hybrid*.
Variety denomination: ‘Poulpmt007’.

SUMMARY OF THE INVENTION

The present invention constitutes a new and distinct variety of garden rose plant which originated from a controlled crossing between the female seed parent, an unnamed, non-commercial seedling, and the male pollen parent, also an unnamed, non-commercial seedling.

The two parents were crossed during the summer of 2000 and the resulting seeds were planted in a controlled environment in Fredensborg, Denmark. The new variety, named ‘Poulpmt007’, originated as a single seedling from the stated cross.

The new variety may be distinguished from its female seed parent primarily by flower color. The seed parent has light pink flowers, while ‘Poulpmt007’ has orange blend flowers.

The new variety may be distinguished from its male pollen parent by flower color. The pollen parent has light pink flowers, while ‘Poulpmt007’ has orange blend flowers.

The objective of the hybridization of this rose variety was to create a new and distinct variety for garden use with unique qualities, such as:

1. Uniform and abundant orange blend hybrid tea shaped flowers;
2. Vigorous, but compact growth when propagated both as a budded rose and on its own roots;
3. Exceptional disease resistance.

This combination of qualities is not present in previously available commercial cultivars of this type, known to the inventor, and distinguish ‘Poulpmt007’ from all other varieties of which we are aware.

As part of the rose development program, Mogens N. Olesen germinated the seeds from the aforementioned hybridization during winter of 2000 and conducted evaluations on the resulting seedlings in a controlled environment in Fredensborg, Denmark. ‘Poulpmt007’ was selected in the spring of 2001 by the inventor as a single plant from the progeny of the aforementioned hybridization.

Asexual reproduction of ‘Poulpmt007’ by traditional budding and rooted cuttings was first done by Mogens N. Olesen

2

in the nursery in Fredensborg, Denmark in July, 2001. This initial and other subsequent asexual propagations conducted in controlled environments have demonstrated that the characteristics of ‘Poulpmt007’ are true to type and are transmitted from one generation to the next.

BRIEF DESCRIPTION OF THE DRAWING

The accompanying color illustration shows as true as is reasonably possible to obtain in color photographs of this type, the typical characteristics of the buds, flowers, leaves, and stems, of ‘Poulpmt007’. Specifically illustrated in the drawing are:

FIG. 1; Open flowers, flower buds, flower petals detached, reproductive flower parts.

FIG. 2; Mature and juvenile leaves, mature and juvenile bare stems.

DETAILED DESCRIPTION OF THE VARIETY

The following is a description of ‘Poulpmt007’, as observed in its growth in in a field nursery in Jackson County, Oreg. Observed plants are 3 years of age, and were grown on *Rosa multiflora* understock. Color references are made using The Royal Horticultural Society (London, England) Colour Chart, 2001, except where common terms of color are used.

For a comparison, several physical characteristics of the rose variety ‘Poulrim’, U.S. Plant Pat. No. 12,465 are compared to ‘Poulpmt007’ in Chart 1.

CHART 1

	‘Poulpmt007’	‘Poulrim’
Petal Count	20 petals	25 to 30 petals
Flower Diameter	50 to 60 mm	100 to 120 mm
General Tonality of Flower Color	On open flower Red Group 52B with intonations of Red-Purple Group 58B.	On open flower blend of Greyed-Orange 170C, Orange 29A with intonations of Red Group 39A.

FLOWER AND FLOWER BUD

Blooming habit: Repeat.

Flower bud:

Size.—Upon opening, 30 mm in length from base of 5
receptacle to end of bud. Bud diameter is 15 mm.

Bud form.—Ovoid.

Bud color.—As sepals unfold, petals are Red-Purple
Group 59B.

Sepal inner surface.—Color: Yellow-Green Group 10
147C with strong intonations of Greyed-Purple
Group 183C. Surface: Moderate pubescence
observed.

Sepal outer surface.—Color: Yellow-Green Group 15
146A. Anthocyanic pigments the color of Greyed-
Purple Group 183A observed. Texture: Smooth.

Sepal Shape.—Apex: Cirrhose. Base: Flat at union with
receptacle.

Sepal Margin.—Margins have medium foliaceous 20
appendages on three of the five sepals.

Sepal size.—30 mm long by 10 mm wide.

Receptacle.—Texture: Smooth. Shape: Broad funnel.
Size: 5 mm in height by 11 mm wide. Color: Yellow-
Green Group 144B.

Pedicel.—Surface: Somewhat rough with small prick- 25
les. Length: 50 mm on average. Diameter: 3 mm on
average. Color: Yellow-Green Group 144A. Weak
anthocyanic pigments the color of Greyed-Red Group
180A observed. Strength: Strong.

Flower bud development: Singularly.

Flower bloom:

Fragrance.—None.

Duration.—The blooms have a duration on the plant of
approximately 10 days. Petals fall cleanly away from 35
plant after flowers have fully matured.

Size.—Flower diameter is 50 to 60 mm when open.
Flower depth is 55 mm on average.

Flower shape.—High centered, semi double, with a
high pointed center which is tightly closed.

Shape of flower, side view.—Open flowers are flat at the
upper portion and concave at the lower portion.

Petalage: Under normal conditions, flowers have 20 petals
total (average), 1 to 3 of which are petaloids.

Petal color:

Upon opening, outer petals.—Upper surface: Orange-
Red Group 33B at the base, blending with Red Group
52B at the middle zone. Red Group 52B at margins.
Lower surface: Red-Purple Group 58B at the mar- 50
ginal and middle zones. Blending with Red Group
51A at the middle and basal zone. Red Group 51A at
the basal zone.

Upon opening, inner petals.—Upper surface: Red
Group 52B at the marginal zone, blending with
Orange-Red Group 33B at the middle and basal zone. 55
Primarily Orange-Red Group 33B at the basal zone.
Lower surface: The base color is Red-Purple Group
58B. An overlay of Red Group 53C at the margins.
Basal zone has blended shades of Red Group 51A.

Basal petal spots, upon opening.—Upper surface: Yel- 60
low Group 5A. Lower surface: Yellow Group 4B.

After opening, inner and outer petals.—Upper surface:
Red-Purple Group 52C blended with Orange-Red
Group 33C at the mid and basal zone. Lower surface:
Red-Purple Group 58B blending with Red Group 50B 65
at the basal zone.

Basal petal spots, after opening.—Upper surface: Yel-
low Group 5C. Lower surface: Yellow Group 5C.

General tonality: On open flower Red Group 52B with into-
nations of Red-Purple Group 58B. No change in the gen-
eral tonality at the end of the 10th day.

Petals:

Petal reflex.—Flat.

Margin.—Entire and uniform.

Shape.—Broad elliptic. Apex shape: Rounded. Base
shape: Rounded.

Size.—40 to 45 mm (l)×40 to 50 mm (w).

Texture.—Smooth.

Thickness.—Above average.

Petaloids:

Quantity.—1 to 3.

Shape.—Asymmetric with an acute base and apex.

Color.—Upper surface is Red Group 52B at the mar-
ginal zone, blending with Orange-Red Group 33B at
the middle and basal zone. Primarily Orange-Red
Group 33B at the basal zone. The lower surface is
Red-Purple Group 58B. An overlay of Red Group
53C at the margins. Basal zone has blended shades of
Red Group 51A. Basal petal spots on the petaloids
are, Yellow Group 5A at the upper portion, and Yellow
Group 4B underneath.

Size.—35 mm long by 12 mm wide on average.

Reproductive organs:

Pollen.—None observed.

Anthers.—Size: 2 mm in length. Color: Yellow Group
13A. Quantity: 200 on average.

Filaments.—Color: Yellow Group 11A. Length: 12 mm.

Pistils.—Length: 12 mm. Quantity: 70 on average.

Stigmas.—Level in location relative to the length of the
filaments and the height of the anthers. Color: Yellow
Group 1C.

Styles.—Color: Red Group 39B.

Hips: None Observed in the field nursery in Jackson County
Oreg.

PLANT

Plant growth: Upright to bushy. When grown as a budded field
grown plant on *Rosa multiflora* understock, the average
height of the plant is 80 cm and the average width is 70 cm.

Stems:

Color.—Juvenile growth: Greyed-Purple Group 183C.
Mature growth: Yellow-Green Group 144B.

Length.—On average, canes are 35 cm from the base of
the plant to the flowering portion.

Diameter.—5 mm.

Internodes.—On mature canes, there is an average dis-
tance of 35 mm between nodes.

Surface texture.—Young wood: Smooth. Older wood:
Smooth.

Prickles:

Incidence.—5 prickles per 10 cm of stem.

Size.—Average length of prickles on mature stems is 9
mm.

Shape.—Linear at upper surface, concave underneath.

Color.—Juvenile prickles: Yellow-Green Group 144B
with intonations of Greyed-Red Group 182A. Mature
prickles: Greyed-Yellow Group 161B.

Plant foliage: Normal number of leaflets leaves in middle of
the stem: 5, leaflets.

Compound leaf.—130 mm (l)×100 (w) on average.

Quantity.—3 leaves per 10 cm of stem on average.
Leaf bearing angle to the stem.—60 degrees.
Color of mature foliage.—Upper side: Yellow-Green Group 147A. Lower side: Yellow-Green Group 147B.
Color of juvenile foliage.—Upper side: Greyed-Purple Group 187A. Lower side: Greyed-Purple Group 187B.
Plant leaves and leaflets:
Stipules.—Size: 23 mm in length. Quantity: 2 per compound leaf. Shape: Linear, slightly broad based with outward extending apices. Margins: Entire with few stipitate glands. Color: Yellow-Green Group 146A.
Petiole.—Length: 25 mm on average. Diameter: 2 to 3 mm.
Upper surface.—Color: Yellow-Green Group 146A.
Lower surface.—Color: Yellow-Green Group 146C.
Rachis.—Length: 30 mm on average.
Upper surface.—Color: Yellow-Green Group 146A.
Lower surface.—Color: Yellow-Green Group 146C.

Leaflet.—Edge: Serrated. Size: Average size of the terminal leaflet on normal leaves is 75 mm in length by 42 mm wide. Shape: Elliptical. Base: Acute. Apex: Acute. Texture: Smooth. Thickness: Thick. Arrangement: Odd pinnate. Venation: Reticulate. Glossiness: Moderately glossy.
Disease resistance: Above average resistance to powdery and downy mildew, rust, black spot, and Botrytis under normal growing conditions in Jackson County, Oreg.
Cold hardiness: The variety is tolerant to USDA Cold Hardiness Zone 6.
Heat tolerance: The variety has been found to be suitable for climate conditions found in the American Horticulture Society heat zone 7.
The invention claimed is:
1. A new and distinct variety of rose plant of the hybrid tea rose class named ‘Poulpmt007’, substantially as illustrated and described herein, due to its abundant orange blend flowers, disease resistance, and extended period of bloom.

* * * * *



