

US00PP23125P2

(12) United States Plant Patent Sheehan

(10) Patent No.:

US PP23,125 P2

(45) Date of Patent:

Oct. 23, 2012

(54) GRAPEVINE 'SHEEGENE-20'

(50) Latin Name: Vitis vinifera

Varietal Denomination: Sheegene-20

(75) Inventor: **Timothy P. Sheehan**, Vancouver, WA

(US)

(73) Assignee: Sheehan Genetics, LLC, Portervile, CA

(US)

(*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: 13/135,332

(22) Filed: Jul. 1, 2011

(51) Int. Cl. A01H 5/00

(2006.01)

Primary Examiner — Annette Para

(74) Attorney, Agent, or Firm — Baker Manock & Jensen, PC; Eric C. Cole, Esq.

(57) ABSTRACT

A new and distinct variety of grapevine characterized by the production of medium to large, red-colored seedless grapes that mature in late September, approximately nine weeks later than 'Flame Seedless' (unpatented) when grown in the San Joaquin Valley of Central California. The grapes of this new variety are produced on strong woody stems and branches and are well adapted to commercial handling.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed: The claimed plant relates to a new and distinct variety of *Vitis vinifera* to be known as 'Sheegene-20'.

Variety denomination: The new variety of *Vitis vinifera* is the result of hybridization of 'Princess' (unpatented), the pollen parent, and 'Red Globe' (U.S. Plant Pat. No. 4,787), the seed parent. The new variety was first hybridized by Timothy P. Sheehan in the spring of 2000. The new variety was asexually propagated in the dormant season of 2003, grafted on one hundred seventy-two vines of 'Freedom' virus-free grape rootstock near Fresno, Calif. The new variety was planted in a *Vitis vinifera* variety block located near Fresno, Calif., on the north side of Muscat Avenue approximately five miles east of State Route 145. The new variety produces a medium-large, red-colored seedless grape with good flavor. The new variety has been shown to maintain its distinguishing characteristics through asexual propagation.

CROSS REFERENCE TO RELATED APPLICATIONS

The claimed plant is not subject of a related application.

STATEMENT OF ANY FEDERALLY-SPONSORED RESEARCH AND DEVELOPMENT

The claimed plant is not subject of Federally-sponsored research or development.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct variety of grapevine that produces a medium-large, red-colored seedless grape that matures during late September when grown in the San Joaquin Valley of Central California. The new invention resembles 'Flame Seedless' (unpatented).

SUMMARY OF THE INVENTION

The 'Sheegene-20' grapevine is characterized by producing a medium-large, red-colored seedless grape that good

2

flavor and it is productive and mature for harvesting and shipment in late September when grown in the San Joaquin Valley of Central California. The new variety can be compared to 'Flame Seedless' (unpatented) but the grapes produced by the new variety mature approximately nine weeks later, among other distinguishing characteristics.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawing is a color photograph that displays leaves along the left side, leaves with tendrils along the right side, clusters of the immature and mature grapes in the center, and on the bottom two grapes cut in half displaying the color of the flesh and the shape of the berries.

DETAILED BOTANICAL DESCRIPTION

Referring more to the horticultural description of the new and distinct variety of grapevine, the following has been observed under the ecological conditions prevailing at the origin vineyard located near Fresno, Calif., in the San Joaquin Valley of Central California. All major color descriptions are by reference to the Dictionary of Color by Maerz & Paul, First Edition, published in 1930. Common colors names are also used in several instances.

Vine:

Size.—Medium to large for two-year old vine.

Vigor.—Very good.

Chilling requirements.—Normal for grapevines in the San Joaquin Valley of Central California.

Figure.—Wide cordons forming a "T" shape figure.

Production capacity.—Very good.

Regularity of bearing.—Regular.

Trunk:

Size.—Medium to large; 4½ inches [11.43 cm] in circumference, 15 inches [38.10 cm] above graft.

Surface texture.—Rough, shaggy.

Color of bark.—Pl.7 H10 Piccadilly, Mauvewood +. Mature cane color.—Pl.7 H9 Liver Brown and Autumn Oak.

3

Nodes.—Three nodes on canes. *Size.*—0.16 inches [4 mm]. Petals (color).—Pl.18 J7 Spring gr. Length between nodes.—3 to 4 inches [7.62 to 10.16] Petals (size).—Small; 0.08 inches [2 mm]. cm. Lenticel numbers.—None. Pistil color.—Pl.17 L8 Neva gr. Date of visible berry set.—May 12, 2010. Shoot length.—17 to 32 inches [43.18 to 81.28 cm]. Shoot shape.—Contour drooping. Size of berries.—Small; 0.08 inches [2 mm]. *Tendrils numbers.*—Single at nodes. Fruit: *Tendrils size.*—Large; 5½ to 8¼ inches [13.97 to 20.96] Solids.—19.2 Brix. Acids.—0.45. cm]. *Tendrils form.*—Bifurcate. Sugar/acid ratio.—42.6. *Juice pH.*—3.32. Tendrils texture.—Firm. Bud shape.—Triangular. Seeds.—None. Bud size.—0.02 inches [6 mm] in width, 0.03 inches [8 Capstem pedicel.—0.24 to 0.35 inches [6 to 9 mm]. Berry weight.—11.36 grams. mm] in length. Branches: Juice color.—Pl.13 L4 Near Bistre gr. Size.—Medium to large; 6 inches [15.24 cm] in circum-Cluster size.—Medium to large. Cluster average length (not including stem).—7.31 ference. Cordons.—32 to 35 inches from trunk (81.28 to 88.90) inches [18.57 cm]. Cluster average diameter.—4.82 inches [12.24 cm]. cm). Surface texture.—Slightly rough with groove. Cluster average weight.—1 lbs. 2 oz. [511.2 grams]. Color (one year or older wood).—Pl.7 H9 Liver Compactness.—Compact. Brown+Autumn Oak. Cluster form.—Conical. Color (immature branches).—Pl.19 K6 Sea gr. on lower *Stem* (*generally*).—2.17 inches [5.51 cm]. Stem (caliper).—0.16 to 0.24 inches [4 to 6 mm]. side of branch; Pl.52 L8 *Cineraria* on upper side of Berry size.—Medium to large. branch. Berry form.—Ovate. Leaves: Size.—Medium to large. Berry numbers.—45 to 75 berries per bunch. Density.—Dense. Berry size (average dimension along longitudinal axis) .—0.98 to 1.26 inches [2.49 to 3.20 cm]. Average length.—4.72 inches [11.99 cm]. Berry size (average dimension along transverse axis).— Average width.—6.89 to 7.09 inches [17.50 to 18.01 30] 0.79 to 0.87 inches [20 to 22 mm]. cm. Skin: *Form.*—Pentagonal. Skin thickness.—Medium. Texture (upper surface).—Smooth. Texture (lower surface).—Slightly glabrous. Texture.—Tough. Color (upward disposed surface).—Pl.23 J9 Mt. Vernon 35 *Tendency to crack.*—None. Blush color.—P1.55 L7 Near Rubient. gr. Color (downward disposed surface).—Pl.21 L7 Med. Ground color.—Pl.12 L1; Oil Y. *Pulp.*—Near Pl.13 K3 Olive gr. Dark gr. Lenticels.—None. Leaf vein color.—Pl.17 L7 Viridine Y. *Marginal form.*—Slightly undulated. 40 Flesh: Leaf vein thickness.—0.008 inches [2 mm]. Flesh color.—Pl.12 K3 Olive gr. Leaf margin.—Toothed. Juice production.—Good. Glandular characteristics.—None. Flavor.—Very good. Aroma.—Mild. Petiole size.—Large. *Petiole length.*—4.53 to 5.12 inches [11.51 to 13.00 cm]. 45 *Texture*.—Firm. Petiole color (lower surface).—Pl.20 L8 Near Eden gr. Ripening.—Even. Petiole color (upper surface).—Pl.16 L8 Dark Cardinal. Eating quality.—Very good. Petiole sinus form.—Upside down 'U' shape. *Use.*—Fresh Market. Resistance to disease.—Uknown. Lobe (average).—Four. Harvesting.—Late August in the San Joaquin Valley of Tooth number.—Four to six. 50 Central California. Tooth shape.—Triangular. Inflorescence: Shipping and handling qualities.—Good. *Size.*—6.70 to 7.50 inches [17.02 to 19.05 cm]. Having thus described and illustrated our new variety of grapevine, I claim: *Number per spur.*—Two to three. 1. A new variety of grapevine to be known as 'Sheegene-Number per vine.—Many; up to sixty-four. Quantity.—Proficient. 20' and parts thereof, substantially as illustrated and described, characterized principally by its production of Flowers: Flower buds.—Small; 0.04 to 0.08 inches [1 to 2 mm]. medium to large, red-colored seedless grapes that mature later in the season of ripening than does 'Flame Seedless' Flower buds surface.—Glabrous.

Flower buds.—Generally hoary.

Date of full bloom.—May 12, 2010.

Date of bloom.—May 9, 2010.

* * * *

60 (unpatented) that it resembles.

