

US00PP23183P2

# (12) United States Plant Patent

## van Haaster

## (10) Patent No.:

## US PP23,183 P2

## (45) **Date of Patent:**

## Nov. 13, 2012

### DAHLIA PLANT NAMED 'FIDAHHYPROBICO'

Latin Name: **Dahlia variabilis** Varietal Denomination: **Fidahhyprobico** 

Johanna G. H. van Haaster, Hillegom Inventor:

(NL)

Assignee: **Fides B.V.**, De Lier (NL)

Subject to any disclaimer, the term of this Notice:

patent is extended or adjusted under 35

U.S.C. 154(b) by 81 days.

Appl. No.: 12/931,175

(22)Jan. 26, 2011 Filed:

Int. Cl. A01H 5/00 (2006.01)

U.S. Cl. Plt./321

(58)See application file for complete search history.

Primary Examiner — June Hwu

(74) Attorney, Agent, or Firm — C. A. Whealy

#### (57)**ABSTRACT**

A new and distinct cultivar of Dahlia plant named 'Fidahhyprobico', characterized by its upright, somewhat outwardly spreading and mounding plant habit; early and freely flowering habit; large inflorescences with light red and yellow bi-colored ray florets; and good postproduction longevity.

## 1 Drawing Sheet

Botanical designation: Dahlia variabilis. Cultivar denomination: 'FIDAHHYPROBICO'.

#### BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of Dahlia plant, botanically known as Dahlia variabilis and hereinafter referred to by the name 'Fidahhyprobico'.

The new *Dahlia* plant is a product of a planned breeding program conducted by the Inventor in Hillegom, The Netherlands. The objective of the breeding program is to create new pot-type Dahlia plants that have a freely branching and flowering habit, attractive ray floret coloration and good postproduction longevity.

The new Dahlia plant originated from an open-pollination in Hillegom, The Netherlands of an unnamed selection of Dahlia variabilis, not patented, as the female, or seed, parent with an unknown selection of Dahlia variabilis as the male, or pollen, parent. The new *Dahlia* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated open-pollination in a controlled environment in Hillegom, The Netherlands.

Asexual reproduction of the new *Dahlia* plant by cuttings in a controlled environment in Hillegom, The Netherlands 25 of this type. Colors in the photographs may differ slightly since 2006 has shown that the unique features of this new Dahlia plant are stable and reproduced true to type in successive generations.

#### SUMMARY OF THE INVENTION

Plants of the new *Dahlia* have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Fidahhyprobico'. These characteristics in combination distinguish 'Fidahhyprobico' as a new and distinct cultivar of Dahlia plant:

1. Upright, somewhat outwardly spreading and mounding plant habit.

- 2. Early and freely flowering habit.
- 3. Large inflorescences with light red and yellow bi-colored ray florets.
- 4. Good postproduction longevity.

Compared to plants of the female parent selection, plants of the new Dahlia differ primarily in ray floret coloration and plant shape.

Plants of the new Dahlia can be compared to plants of Dahlia 'Fidahhyppi', disclosed in U.S. Plant Pat. No. 21,254. In side-by-side comparisons conducted in De Lier, The Netherlands, plants of the new Dahlia differed primarily from plants of 'Fidahhyppi' in the following characteristics:

- 1. Plants of the new *Dahlia* had smaller inflorescences than plants of 'Fidahhyppi'.
- 2. Plants of the new *Dahlia* and 'Fidahhyppi' differed in ray floret color.

#### BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying photograph illustrates the overall appearance of the new *Dahlia* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions from the color values cited in the detailed botanical description which accurately describe the colors of the new *Dahlia* plant. The photograph comprises a side perspective view of a typical flowering plant of 'Fidahhyprobico' grown in a con-30 tainer.

### DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and the following obser-35 vations and measurements describe plants grown during the autumn in 15-cm containers in a glass-covered greenhouse in De Lier, The Netherlands and under conditions and practices which approximate those generally used in commercial potted Dahlia production. During the production of the plants, day and night temperatures averaged 18° C. Plants were nine weeks old when the photograph and description were taken. In the following description, color references are made to The

10

45

50

Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Dahlia variabilis* 'Fidahhyprobico'. Parentage:

Female, or seed, parent.—Unnamed selection of Dahlia variabilis, not patented.

Male, or pollen, parent.—Unknown selection of Dahlia variabilis, not patented.

#### Propagation:

*Type.*—By cuttings.

Time to initiate roots, summer.—About six days at temperatures of about 22° C.

Time to initiate roots, winter.—About eight days at temperatures of about 20° C.

Time to produce a rooted young plant, summer.—About 12 days at temperatures of about 22° C.

Time to produce a rooted young plant, winter.—About 16 days at temperatures of about 20° C.

Root description.—Fine, fibrous; tuber development has not been observed.

Rooting habit.—Freely branching; dense.

#### Plant description:

Plant form and growth habit.—Upright, somewhat outwardly spreading and mounding plant form; broad inverted triangle; three primary lateral branches develop, each primary lateral branch with numerous secondary branches; inflorescences held above the foliar plane on strong peduncles; bushy and dense habit; moderately vigorous growth habit.

Plant height.—About 27 cm.

Plant diameter or spread.—About 28 cm.

Lateral branches.—Length: About 18 cm. Diameter: About 9 mm. Internode length: About 1.5 cm to 2 cm. Aspect: Erect to somewhat outwardly spreading. <sup>35</sup> Strength: Moderately strong. Texture: Smooth, glabrous. Color: Close to 144A.

#### Foliage description:

Arrangement.—Leaves opposite; leaves may be simple or compound with three or five leaflets; measure- 40 ments are for simple leaves.

Length.—About 11.5 cm.

Width.—About 6 cm.

Shape.—Ovate.

Apex.—Acute.

Base.—Attenuate.

Margin.—Serrate to dentate.

Venation pattern.—Pinnate.

Texture, upper and lower surfaces.—Slightly pubescent; slightly rough.

Color.—Developing leaves, upper surface: Close to 137B. Developing leaves, lower surface: Close to 138B. Fully expanded leaves, upper surface: Close to N137A; venation, close to N137A. Fully expanded leaves, lower surface: Close to 138B; venation, close to 138B.

Petioles.—Length: About 4.5 cm. Diameter: About 2 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 144A.

## Inflorescence description:

Appearance and arrangement.—Rotate double inflorescence form with ray florets forming acropetally on a receptacle; inflorescences positioned above the foliar plane on strong peduncles; inflorescences face

upright to outwardly; freely flowering habit, about 20 inflorescences develop per plant.

Fragrance.—None detected.

Time to flower.—Plants flower continuously from spring through the autumn in The Netherlands; early flowering habit, plants begin flowering about seven to eight weeks after planting.

Post-production longevity.—Inflorescences maintain good substance for about two weeks on the plant; inflorescences persistent.

*Inflorescence bud.*—Height: About 4 mm. Diameter: About 7 mm. Shape: Globular, flattened. Color: Close to 144A.

Inflorescence size.—Diameter: About 8.5 cm. Depth (height): About 3 cm. Disc diameter: About 7 mm. Receptacle height: About 3 mm. Receptacle diameter: About 2 cm.

Ray florets.—Quantity per inflorescence: About 114 arranged in about ten whorls. Length: About 3 cm. Width: About 1.4 cm. Shape: Oblanceolate. Apex: Obtuse. Base: Cuneate. Margin: Entire. Aspect: Initially upright to eventually roughly perpendicular to the peduncle; ray florets cupped. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening, upper surface: Towards the apex, close to 54A; mid-section and towards the base, close to 1A. When opening, lower surface: Close to 1B tinted with close to 54A to 54C. Fully opened, upper surface: Towards the apex, close to 54B to 54C; mid-section and towards the base, close to 1A. When opening, lower surface: Close to 1B tinted with close to 54A to 54C.

Disc florets.—Length: About 8 mm. Diameter: About 1 mm. Shape: Tubular, elongated; apices obtuse. Number of disc florets per inflorescence: About 33. Color, when opening and fully opened: Apex and mid-section: Close to 17A. Base: Close to 145D.

Phyllaries.—Quantity per inflorescence: About eight arranged in a single whorl. Length: About 1.2 cm. Width: About 5 mm. Shape: Ovate. Apex: Acute. Base: Cuneate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper surface: Close to 137A. Color, lower surface: Close to 138A.

Peduncles.—Length: About 8 cm. Diameter: About 3 mm. Aspect: Mostly erect. Strength: Strong. Texture: Smooth, glabrous. Color: Close to 144A tinted with close to 183B.

Reproductive organs.—Androecium: Quantity per floret: About two. Filament length: About 8 mm. Filament color: Close to 17A. Anther shape: Narrowly elliptic. Anther length: About 2 mm. Anther color: Close to 17A. Pollen amount: Moderate. Pollen color: Close to 17A. Gynoecium: Not observed. Seeds/fruits: Seed and fruit development have not been observed.

Disease/pest resistance: Plants of the new *Dahlia* have not been shown to be resistant to pathogens and pests common to *Dahlia*.

Temperature tolerance: Plants of the new *Dahlia* tolerate high temperatures of about 35° C. and are hardy to USDA Hardiness Zone 8.

It is claimed:

1. A new and distinct *Dahlia* plant named 'Fidahhyprobico' as illustrated and described.

\* \* \* \* \*

