



US00PP23291P2

(12) **United States Plant Patent**
Ackerman et al.(10) **Patent No.:** US PP23,291 P2
(45) **Date of Patent:** Jan. 1, 2013(54) **STRAWBERRY PLANT NAMED 'PS-3.108'**(50) Latin Name: *Fragaria ananassa*
Varietal Denomination: PS-3.108(75) Inventors: **Stephen M. Ackerman**, Salinas, CA (US); **Steven D. Nelson**, Watsonville, CA (US); **Michael D. Nelson**, Watsonville, CA (US)(73) Assignees: **Berry R&D, Inc.**, Watsonville, CA (US); **Plant Sciences, Inc.**, Watsonville, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: 13/137,804

(22) Filed: Sep. 14, 2011

(51) **Int. Cl.**

A01H 5/00 (2006.01)

(52) **U.S. Cl.** Plt./209(58) **Field of Classification Search** Plt./209
See application file for complete search history.

Primary Examiner — Annette Para

(74) Attorney, Agent, or Firm — Foley & Lardner LLP

ABSTRACT

This invention relates to a new and distinct variety of strawberry plant named 'PS-3.108'. This new strawberry plant named 'PS-3.108' is primarily adapted to the growing conditions of the central coast of California, and is primarily characterized by its dark red to red fruit color, large to very large fruit size, medium plant vigor, conical fruit shape, firm fruit flesh, and early fruit ripening.

4 Drawing Sheets**1**

Latin name of the genus and species of the plant claimed:
Fragaria ananassa.

Variety denomination: 'PS-3.108'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct strawberry variety named 'PS-3.108'. This new variety is a result of a controlled cross made in 2003 in an ongoing breeding program between strawberry variety designated 'PS-1269' (patented, U.S. Plant Pat. No. 10,686) and strawberry variety designated 'PS-4634' (patented, U.S. Plant Pat. No. 17,487). Due to the combining of the reciprocal seed lots, it is unknown as to which parent variety is the seed parent and which parent variety is the pollen parent. The variety is botanically known as *Fragaria ananassa*.

The seedling resulting from the aforementioned cross was selected from a controlled breeding plot in Monterey County, Calif. in the summer of 2005. After its selection, the new variety was asexually propagated by stolons in both San Joaquin County, Calif. and Siskiyou County, Calif. The new variety was extensively tested over the next several years in fruiting fields in Monterey County, Calif. This propagation has demonstrated that the combination of traits disclosed herein as characterizing the new variety are fixed and remain true to type through successive generations of asexual reproduction.

BRIEF SUMMARY OF THE INVENTION

'PS-3.108' is primarily adapted to the climate and growing conditions of the central coast of California. The nearby Pacific Ocean provides the needed humidity and moderate temperatures to produce a strong vigorous plant and maintain fruit quality during the April through November production months.

The following traits have been repeatedly observed and are determined to be unique characteristics of 'PS-3.108', which in combination distinguish this strawberry plant as a new and distinct variety:

2

1. Dark red to red fruit color;
2. Large to very large fruit size;
3. Medium plant vigor;
4. Conical fruit shape;
5. Firm fruit flesh; and
6. Early fruit ripening

The strawberry variety that is believed to be most closely related to the new variety 'PS-3.108' is 'PS-4634' (patented, U.S. Plant Pat. No. 17,487). In side-by-side comparisons to the similar strawberry variety 'PS-4634', 'PS-3.108' differs by the following combination of characteristics as described in Table 1.

TABLE 1

COMPARISON TO THE STANDARD VARIETY		
Characteristic	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
Fruit color	Ranges from red to dark red	Ranges from orange red to red
Average marketable yield	1,969 grams/plant	1,689 grams/plant
Difference in shape between primary and secondary fruit	Ranges from none to very slight	Moderate
Size of calyx in relationship to fruit diameter	Slightly larger	Same size
Firmness of flesh	Firm	Medium
Expression of hollow center	Weak	Strong
Plant size	Medium	Large
Foliage color	Ranges from light to medium yellow green	Medium green
Foliage shape in cross section	Ranges from lightly to strongly concave	Ranges from lightly concave to flat

For identification, a series of molecular markers have been determined for this new variety.

'PS-3.108' differs from its parents, 'PS-1269' and 'PS-4634' by the following combination of characteristics as described in Table 2.

TABLE 2

COMPARISON TO THE PARENT VARIETIES			
Characteristic	'PS-3.108'	'PS-1269' (U.S. Plant Pat. No. 10,686)	'PS-4634' (U.S. Plant Pat. No. 17,487)
Fruit color	Ranges from red to dark red	Red	Ranges from orange red to red
Average marketable yield	1,969 grams/plant	1,443 grams/plant	1,689 grams/plant
Difference in shape between primary and secondary fruit	Ranges from none to very slight	Ranges from moderate to marked	Moderate
Fruit gloss	Medium	Ranges from weak to medium	Strong

BRIEF DESCRIPTIONS OF THE PHOTOGRAPHS

The accompanying color photographs illustrate the overall appearance of typical specimens of the new strawberry variety 'PS-3.108', at various stages of development as true as it is reasonably possible with color reproductions of this type. Color in the photographs may differ slightly from the color value cited in the botanical descriptions which accurately describe the color of 'PS-3.108'. The depicted plant and plant parts of the new strawberry variety 'PS-3.108' are between eight and nine months old. The photographs were taken in Monterey County, Calif.:

FIG. 1 shows typical fruiting field characteristics of 'PS-3.108', taken in the month of July 2011;

FIG. 2 shows a close-up view of the typical leaf structure of 'PS-3.108', taken in the month of July 2011;

FIG. 3 shows typical mature and immature field fruit of 'PS-3.108', taken in the month of July 2011; and

FIG. 4 shows typical internal and external mature fruit characteristics of 'PS-3.108', taken in the month of August 2011.

DETAILED BOTANICAL DESCRIPTION

The new variety 'PS-3.108' has not been observed under all possible environmental conditions. The characteristics of the new variety 'PS-3.108' may vary in detail, depending upon variations in environmental factors, including weather (temperature, humidity and light intensity), day length, soil type and location. In addition, the characteristics of any parental variety or comparison variety included in Tables 1-9 of the present invention may vary in detail, depending upon variations in environmental factors, including weather (temperature, humidity and light intensity), day length, soil type and location.

The aforementioned photographs, together with the following description of the new variety 'PS-3.108', unless otherwise noted, are based on observations taken during the 2011 growing season in Monterey County, Calif. These measurements and ratings were taken from plants of 'PS-3.108' dug from a high-elevation nursery located in Siskiyou County, Calif. during mid October 2010 and planted between two and three weeks later in Monterey County, Calif. The approximate age of the observed plants is between eight and nine months. Yield observations and fruit quality characteristics are averaged from three years of data collected from the 2008 through 2010 growing seasons. Flower measurements and characteristics are from secondary flowers unless otherwise

noted. Fruit characteristics and measurements are from secondary fruit unless otherwise noted.

Color terminology where noted follows The Royal Horticultural Society Colour Chart, London (2007).

The following tables 3 through 9 describe fruit, plant, stolon, foliage, fruiting truss, flower and pest disease characteristics of the new strawberry 'PS-3.108' in comparison to the similar strawberry variety 'PS-4634'.

TABLE 3

FRUIT CHARACTERISTICS		
Characteristic	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
Color of mature fruit	RHS 46A Ranges from red to dark red	RHS 45B Ranges from orange red to red
Color of internal flesh	RHS 44B Medium red	RHS 44C Ranges from light red to medium red
Fruit length (cm)	4.5	4.3
Fruit width (cm)	3.8	3.9
Fruit length/width ratio	1.17 Slightly longer than broad	1.11 Slightly longer than broad
Calyx diameter (cm)	4.2	4.0
Average fruit weight (gm)	29.6	27.7
30 Achene color, shaded side	RHS 162B Greyed yellow group	RHS 162C Greyed yellow group
Achene color, sun-exposed side	RHS 184B Greyed purple group	RHS 184C Greyed purple group
Achenes per berry	438	437
Achene weight (mg)	0.52	0.40
Marketable fruit yield (gm/plant)	1,969	1,689
40 Fruit size	Ranges from large to very large	Ranges from large to very large
Predominant fruit shape	Conical	Conical
Difference in shapes between primary and secondary fruit	None or very slight	Moderate
Band without achenes	Absent or very narrow	Narrow
Unevenness of fruit surface	Absent or very weak	Weak
45 Evenness of fruit color	Even	Ranges from slightly uneven to even
Fruit glossiness	Medium	Strong
Insertion of achenes	Level with surface	Level with surface
Insertion of calyx	In basin	Ranges from in basin to level
50 Attitude of calyx	Spreading	Ranges from spreading to reflexed
Size of calyx in relation to fruit diameter	Slightly larger	Same size
Adherence of calyx	Strong	Strong
Firmness of fruit skin	Strong	Strong
Firmness of fruit flesh	Firm	Medium
Keeping quality	Very good	Very good
Distribution of red color of the flesh	Marginal and central	Marginal and central
Hollow center expression	Weak	Strong
Fruit flavor	Good	Very good
Soluble solids (% brix)	7.4	8.3
Time of first flowering	Early	Early
Time of first harvesting	Early	Early
Harvest period	Late March to November	Late March to November
Type of bearing	June bearing	June bearing

US PP23,291 P2

5

6

TABLE 4

PLANT CHARACTERISTICS		
Characteristic	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
Plant height (cm)	25.3	28.8
Plant spread (cm)	33.9	39.1
Plant size	Medium	Large
Plant habit	Globose	Globose
Plant density	Medium	Medium
Plant vigor	Medium	Strong

TABLE 5

STOLON CHARACTERISTICS		
Characteristic	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
Stolon color	RHS 146D Yellow green group	RHS 146D Yellow green group
Stolon anthocyanin coloration	RHS 182A Greyed red group	RHS 182D Greyed red group
Stolon anthocyanin intensity	Weak	Ranges from weak to medium
Stolon pubescence	Medium	Ranges from weak to medium
Attitude of hairs	Upward	Upward
Average stolon quantity	Ranges from medium to many	Ranges from medium to many
Stolon diameter at bract (mm)	3.7	3.4

TABLE 6

FOLIAGE CHARACTERISTICS		
Characteristic	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
<u>Foliage:</u>		
Color of upper surface	RHS 146A Ranges from light yellow green to medium yellow green	RHS 137A Medium green
Color of underside	RHS 147C Yellow green group	RHS 147C Yellow green group
Number of leaflets	3	3
Shape in cross section	Strongly to slightly concave	Slightly concave to flat
Interveinal blistering	Ranges from medium to strong	Ranges from medium to strong
Leaf glossiness	Ranges from medium to strong	Ranges from medium to strong
<u>Terminal Leaflet:</u>		
Length (cm)	8.5	9.5
Width (cm)	7.7	6.9
Leaf size	Ranges from large to medium	Ranges from large to medium
Length/width ratio	1.11	1.36
Shape of base	Obtuse	Acute
Shape of teeth	Obtuse	Obtuse
Serrations per leaf	18.7	22.1
<u>Petiole:</u>		
Petiole color	RHS 145A Yellow green group	RHS 145A Yellow green group
Petiole length (cm)	15.6	20.2
Petiole diameter (mm)	4.0	4.4
Petiolule color	RHS 145A Yellow green group	RHS 145A Yellow green group

TABLE 6-continued

FOLIAGE CHARACTERISTICS		
Characteristic	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
Petiolule length (mm)	9.7	7.8
Attitude of hairs	Strongly outward	Strongly outward
Petiole pubescence	Moderate	Ranges from moderate to sparse

15 Stipe:

Color	RHS 146B	RHS 146B
	Yellow green group	Yellow green group
Anthocyanin coloration	RHS 63A	RHS 58A
Anthocyanin intensity	Red purple group	Red purple group
Length (mm)	18.1	18.7
Width (mm)	9.9	8.3

TABLE 7

FRUITING TRUSS CHARACTERISTICS		
Characteristic	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
Anthocyanin coloration	RHS 181C	RHS 182D
Anthocyanin intensity	Greyed red group	Greyed red group
Length at maturity (cm)	Absent or very weak	Absent or very weak
Position relative to foliage	29.5	35.3
Pubescence	Level with	Level with
Attitude at first pick	Medium	Ranges from weak to medium
	Prostrate	Prostrate

TABLE 8

FLOWER CHARACTERISTICS		
Characteristic	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
Petal color	RHS NN 155C	RHS NN 155C
	White group	White group
Sepal color	RHS 137C	RHS 137A
	Green group	Green group
Corolla diameter (mm)	26.3	29.9
Calyx diameter (mm)	36.5	35.7
Petal length (mm)	11.3	12.3
Petal width (mm)	12.0	12.5
Petal length/width ratio	0.94	0.98
Petals/flower	6.8	5.6
Sepal length (mm)	13.9	12.8
Sepal width (mm)	5.8	5.2
Sepal length/width ratio	2.40	2.46
Sepals/flower	13.7	11.3
Size of calyx relative to corolla	Larger	Larger
Size of inner calyx relative to outer calyx	Ranges from smaller to same	Ranges from same to larger
Relative position of petals	Overlapping	Overlapping

65

US PP23,291 P2

7

TABLE 9

PEST AND DISEASE REACTIONS		
	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
Powdery mildew	Moderate	Moderately resistant
Angular leaf spot	Susceptible	Susceptible
Botrytis fruit rot	Moderately susceptible	Moderately susceptible
Two-spotted spider mite	Moderately susceptible	Moderately susceptible

8

TABLE 9-continued

PEST AND DISEASE REACTIONS		
	'PS-3.108'	'PS-4634' (U.S. Plant Pat. No. 17,487)
5 Characteristic	'PS-3.108'	'PS-4634'
Lygus bug	Susceptible	Susceptible
Flower thrips	Moderately resistant	Moderately susceptible

10 We claim:

1. A new and distinct strawberry plant named 'PS-3.108', as herein described and illustrated by the characteristics set forth above.

* * * * *

FIG. 1




FIG. 2




FIG. 3





FIG. 4



UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : PP23,291 P2
APPLICATION NO. : 13/137804
DATED : January 1, 2013
INVENTOR(S) : Ackerman et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page:

Item (73) "Assignees: Berry R&D, Inc., Watsonville, CA
(US); Plant Sciences, Inc., Watsonville,
CA (US)"

should be changed to:

(73) --Assignee: Plant Sciences, Inc., Watsonville, CA (US)--

Signed and Sealed this
Tenth Day of May, 2016



Michelle K. Lee
Director of the United States Patent and Trademark Office