



US00PP23562P2

(12) **United States Plant Patent**
Smith

(10) **Patent No.:** **US PP23,562 P2**
(45) **Date of Patent:** **Apr. 23, 2013**

(54) **CHRYSANTHEMUM PLANT NAMED**
‘CIFZ0002’

(50) Latin Name: *Chrysanthemum*×*morifolium*
Varietal Denomination: **CIFZ0002**

(75) Inventor: **Mark A. Smith**, Alva, FL (US)

(73) Assignee: **Syngenta Crop Science Protection AG**,
Basel (CH)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/385,185**

(22) Filed: **Feb. 6, 2012**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.** **Plt./293**

(58) **Field of Classification Search** Plt./293
See application file for complete search history.

Primary Examiner — Annette Para

(74) *Attorney, Agent, or Firm* — Joshua L. Price

(57) **ABSTRACT**

A new *Chrysanthemum* plant named ‘CIFZ0002’ particularly
distinguished by the small red decorative inflorescences, yel-
low-green foliage, medium ball-shaped plant habit, and a
natural flowering season response of late September.

1 Drawing Sheet

1

Latin name of the genus and species of the plant claimed:
Chrysanthemum×*morifolium*.
Varietal denomination: ‘CIFZ0002’.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new *Chrysanthemum*,
botanically known as *Chrysanthemum*×*morifolium*, and
hereinafter referred to by the variety name ‘CIFZ0002’.

‘CIFZ0002’ is a product of a planned breeding program.
The new cultivar has small red decorative inflorescences,
yellow-green foliage, medium ball-shaped plant habit, and a
natural flowering season response of late September.

‘CIFZ0002’ originates as a natural whole plant mutation of
an unpatented variety designated ‘05-M166’. ‘CIFZ0002’
was discovered and selected by the inventor as a single flow-
ering plant within a large population of the parent cultivar,
grown massed in pots outdoors in a controlled breeding pro-
gram in Alva, Fla. in April 2008. The parent cultivar ‘05-
M166’ has purple inflorescence color, similar plant size and
habit, with a natural season flowering response that is a few
days slower than ‘CIFZ0002’.

The first act of asexual reproduction of ‘CIFZ0002’ was
accomplished when vegetative cuttings were propagated
from the initial selection in May 2008 in a greenhouse in Alva,
Fla.

BRIEF SUMMARY OF INVENTION

Horticultural examination of plants grown from cuttings of
the plant initiated in May 2008, and continuing thereafter, has
demonstrated that the combination of characteristics as
herein disclosed for ‘CIFZ0002’ are firmly fixed and are
retained through successive generations of asexual reproduc-
tion.

‘CIFZ0002’ has not been observed under all possible envi-
ronmental conditions. The phenotype may vary significantly
with variations in environment such as temperature, light
intensity, and day length.

Plant Breeder’s Rights for this cultivar were applied for in
Canada on Feb. 24, 2011, No. 11-7176 and in the Community
Plant Variety Office on Mar. 30, 2011, No. 2011/0787.

2

‘CIFZ0002’ has not been made publicly available more than
one year prior to the filing of this application.

The following traits have been repeatedly observed and are
determined to be basic characteristics of the new variety. The
combination of these characteristics distinguishes this *Chry-*
santhemum as a new and distinct variety.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying photographic drawing shows typical
flower and foliage characteristics of ‘CIFZ0002’ with colors
being as true as possible with an illustration of this type. The
photographic drawing shows a flowering potted plant of the
new variety and a close-up of the flowers.

DETAILED BOTANICAL DESCRIPTION

The plant descriptions, measurements and aforementioned
close-up photograph were collected and/or taken outdoors in
Gilroy, Calif. in early November 2011. These plants were
grown outdoors in gallon pots and were approximately 12
weeks of age.

The aforementioned photograph of the whole plant was
taken in later September 2011 in Monroeville, N.J. and plants
were about 15 weeks old, grown outdoors.

Color references are made to The Royal Horticultural Soci-
ety Colour Chart (R.H.S.) 2001.

TABLE 1

DIFFERENCES BETWEEN THE NEW VARIETY ‘CIFZ0002’
AND A MOST SIMILAR VARIETY

	‘CIFZ0002’	‘Gedi One Sav’ (U.S. Plant Pat. No. 13,928)
Inflorescence diameter:	Smaller	Larger
General plant size:	Smaller	Larger
Black cloth and natural flowering season response:	Faster	Slower

Plant:

Form, growth and habit.—Herbaceous garden-type, stems upright and outwardly spreading, freely branching.

Plant height.—12.0-13.0 cm.

Plant height (inflorescence included).—14.0-15.0 cm.

Plant width.—22.0-24.0 cm.

Roots:

Number of days to initiate roots.—About 4 days at about 22 degrees C.

Number of days to produce a rooted cutting.—4-6 days at 22 degrees C.

Type.—Fine, fibrous, free branching.

Color.—RHS N155B but whiter.

Foliage:

Arrangement.—Alternate.

Immature, leaf color, upper surface.—RHS 147A but a little lighter.

Lower surface.—Closest to RHS 147B.

Mature, leaf color, upper surface.—Closest to RHS 147A.

Lower surface.—Closest to RHS 147B.

Length.—3.1-3.4 cm.

Width.—3.0-3.1 cm.

Shape.—Ovate.

Base shape.—Attenuate.

Apex shape.—Mucronulate.

Margin.—Palmately lobed, irregularly incised, serrate.

Texture, upper surface.—Bifid T-shaped hairs.

Lower surface.—Bifid T-shaped hairs.

Color of veins, upper surface.—RHS 138B.

Color of veins, lower surface.—RHS 138B.

Petiole color.—RHS 138B.

Length.—0.8-1.2 cm.

Diameter.—0.2 cm.

Texture.—Bifid T-shaped hairs.

Stem:

Quantity of main branches per plant.—About 10.

Color of stem.—RHS 138A, but appears lighter due to the hairs.

Length of stem.—6.0-7.0 cm.

Diameter.—0.25-0.3 cm.

Length of internodes.—0.7-1.0 cm.

Texture.—Bifid T-shaped hairs.

Color of peduncle.—RHS 138B but appears lighter due to the hairs.

Length of peduncle.—3.5-5.0 cm.

Peduncle diameter.—0.1 cm.

Texture.—Bifid T-shaped hairs.

Inflorescence:

Type.—Compositae, solitary decorative-type inflorescences; borne terminally above foliage, ray florets arranged acropetally on a capitulum.

Quantity of inflorescences per plant.—About 125.

Lastingness of individual blooms on the plant.—A little less than 6 weeks from first color showing.

Fragrance.—Slightly spicy.

Bud (just when opening/showing color):

Color.—RHS 184A.

Length.—0.9-1.1 cm.

Width.—0.8-0.9 cm.

Shape.—Oblate.

Immature inflorescence:

Diameter.—2.0-2.5 cm.

Color of ray florets, upper surface.—Closest to RHS 53A.

Lower surface.—RHS 182A.

Mature inflorescence:

Diameter.—3.1-3.5 cm.

Depth.—1.5 cm.

Total diameter of disc.—0.1-0.2 cm.

Receptacle height.—0.3 cm.

Receptacle diameter.—0.35.

Ray florets:

Average quantity of florets.—About 90 in several whorls.

Color of florets, upper surface.—RHS 53A, maturing to RHS 185B.

Lower surface.—Closest to RHS 182A maturing to about RHS 182C.

Length.—1.1 cm.

Width.—0.3-0.4 cm.

Shape.—Oblong.

Apex shape.—Praemorse.

Margin.—Entire.

Texture, upper surface.—Papillose.

Lower surface.—Papillose.

Disc florets:

Average quantity of florets.—13-22.

Color of florets.—RHS 155C with RHS 6C apex.

Length.—0.3-0.4 cm.

Width.—0.1 cm.

Shape.—Tubular, elongated.

Apex shape.—Acute, 5 pointed.

Phyllaries:

Quantity.—About 35.

Color, upper surface.—RHS 144A.

Lower surface.—RHS 138A.

Length.—0.5 cm.

Width.—0.15 cm.

Shape.—Lanceolate.

Apex shape.—Acute.

Base.—Fused.

Margins.—Entire; some are papery with RHS 200B margins.

Texture, upper surface.—Glabrous.

Lower surface.—Bifid T-shaped hairs.

Reproductive organs:

Pistil.—1, found on both types of florets.

Length.—0.4-0.5 cm.

Style color.—RHS 155C.

Style length.—0.3-0.4 cm.

Stigma color.—RHS 6C.

Stigma shape.—Bi-parted.

Ovary color.—RHS 155C.

Ovary length.—0.1 cm.

Ovary width.—0.05 cm.

Stamens.—4, found on only on the disc florets.

Color of filaments.—RHS 155C.

Length filaments.—0.2-0.3 cm.

Anther color.—RHS 15C.

Anther length.—0.1 cm.

Anther shape.—Oval.

Pollen amount.—None found.

What is claimed is:

1. A new and distinct variety of *Chrysanthemum* plant named 'CIFZ0002' substantially as illustrated and described herein.

* * * * *



UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : PP23,562 P2
APPLICATION NO. : 13/385185
DATED : April 23, 2013
INVENTOR(S) : Smith

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title Page, Item (73) Assignee:, delete “Syngenta Crop Science Protection AG, Basel (CH)” and insert therefor --Syngenta Crop Protection AG, Basel (CH)--

In the Specification

At column 4, line 6, delete “0.35” and insert therefor --0.35 cm--

Signed and Sealed this
Eighteenth Day of June, 2013



Teresa Stanek Rea
Acting Director of the United States Patent and Trademark Office