

US00PP23810P2

(12) United States Plant Patent Dümmen

(10) Patent No.:

US PP23,810 P2

(45) **Date of Patent:**

Aug. 6, 2013

(54) PETUNIA PLANT NAMED 'DUEPOTPIBLIC'

(50) Latin Name: *Petunia*×*hybrida*Varietal Denomination: **Duepotpiblic**

(75) Inventor: **Tobias Dümmen**, Rheinberg (DE)

(73) Assignee: Capital Green Investments Ltd., Grand

Cayman (KY)

(*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 74 days.

(21) Appl. No.: 13/317,899

(22) Filed: Oct. 31, 2011

(51) Int. Cl. A01H 5/00

(2006.01)

(58) Field of Classification Search

Primary Examiner — Annette Para

(74) Attorney, Agent, or Firm — C. A. Whealy

(57) ABSTRACT

A new and distinct cultivar of *Petunia* plant named 'Duepot-piblic', characterized by its compact, mounding and outwardly spreading plant habit; freely branching habit; early and freely flowering habit; light violet-colored flowers with dark violet blue-colored venation; and good garden performance.

1 Drawing Sheet

1

Botanical designation: *Petunia*×*hybrida*. Cultivar denomination: 'DUEPOTPIBLIC'.

CROSS-REFERENCED TO CLOSELY RELATED APPLICATIONS

Title: Petunia Plant Named 'Duepotpipuce'

Applicant: Tobias Dümmen

Filed: U.S. Plant patent application Ser. No. 13/317,912.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Petunia* plant, botanically known as *Petunia*×*hybrida* and hereinafter referred to by the name 'Duepotpiblic'.

The new *Petunia* plant is a product of a planned breeding program conducted by the Inventor in Rheinberg, Germany. The objective of the breeding program is to create new compact *Petunia* plants with numerous unique and attractive flowers.

The new *Petunia* plant originated from a cross-pollination made by the Inventor in July, 2008 in Rheinberg, Germany of a proprietary selection of *Petunia*×*hybrida* identified as code number T06-0921-004, not patented, as the female, or seed, parent with a proprietary selection of *Petunia*×*hybrida* identified as code number T06-2749-002, not patented, as the male, or pollen, parent. The new *Petunia* plant was discovered and selected by the Inventor as a single flowering plant within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rheinberg, Germany in May, 30 2010.

Asexual reproduction of the new *Petunia* plant by terminal cuttings in a controlled greenhouse environment in Rheinberg, Germany since June, 2010 has shown that the unique features of this new *Petunia* plant are stable and reproduced 35 true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Petunia* have not been observed under all possible environmental conditions and cultural practices. The

phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Duepotpiblic'. These characteristics in combination distinguish 'Duepotpiblic' as a new and distinct *Petunia* plant:

- 1. Compact, mounding and outwardly spreading plant habit.
- 2. Freely branching habit.
- 3. Early and freely flowering habit.

without, however, any variance in genotype.

- 4. Light violet-colored flowers with dark violet blue-colored venation.
- 5. Good garden performance.

Plants of the new *Petunia* can be compared to plants of the female parent selection. Plants of the new *Petunia* differ primarily from plants of the female parent selection in the following characteristics:

- 1. Plants of the new *Petunia* are more compact than plants of the female parent selection.
- 2. Plants of the new *Petunia* have smaller flowers than plants of the female parent selection.

Plants of the new *Petunia* can be compared to plants of the male parent selection. Plants of the new *Petunia* differ primarily from plants of the male parent selection in flower color as plants of the male parent selection have white-colored flowers.

Plants of the new *Petunia* can be compared to plants of *Petunia*×*hybrida* 'Duepotpipuce', disclosed in a U.S. Plant patent application Ser. No. 13/317,912. Plants of the new *Petunia* differ primarily from plants of 'Duepotpipuce' in flower color as plants of 'Duepotpipuce' have light violet-colored flowers with red purple-colored venation.

Plants of the new *Petunia* can be compared to plants of the *Petunia*×*hybrida* 'Duesurimbuvein', not patented. In sideby-side comparisons conducted in Rheinberg, Germany, plants of the new *Petunia* differed primarily from plants of 'Duesurimbuvein' in the following characteristics:

1. Plants of the new *Petunia* were more compact than plants of 'Duesurimbuvein'.

- 2. Plants of the new *Petunia* had larger leaves than plants of 'Duesurimbuvein'.
- 3. Plants of the new *Petunia* had smaller flowers than plants of 'Duesurimbuvein'.
- 4. Plants of the new *Petunia* and 'Duesurimbuvein' differed 5 in flower color as plants of 'Duesurimbuvein' had whitecolored flowers with violet blue-colored venation.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Petunia* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical descrip- 15 tion which accurately describe the colors of the new *Petunia* plant. The photograph comprises a side perspective view of a typical flowering plant of 'Duepotpiblic' grown in a container.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown during the summer in 12-cm containers in a glass-covered green- 25 house in Rheinberg, Germany and under cultural practices typical of commercial *Petunia* production. During the production of the plants, day and night temperatures averaged 18° C. and light levels averaged 4,500 lux. Plants were pinched one time three weeks after planting and were 16 30 weeks old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Petunia*×*hybrida* 'Duepotpiblic'. Parentage:

Female, or seed, parent.—Proprietary selection of Petunia×hybrida identified as code number T06-0921-004, not patented.

Male, or pollen, parent.—Proprietary selection of Petunia×hybrida identified as code number T06-2749-002, not patented.

Propagation:

Type.—By terminal cuttings.

Time to initiate roots, summer.—About five days at temperatures of about 20° C.

Time to initiate roots, winter.—About seven days at temperatures of about 20° C.

Time to produce a rooted young plant, summer.—About 50 three weeks at temperatures of about 20° C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures of about 20° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Compact, mounding and outwardly spreading plant habit; freely branching habit with about eight to ten lateral branches developing after pinching; moderately vigorous growth habit.

Plant height.—About 12 cm.

Plant diameter.—About 20 cm.

Lateral branch description:

Length.—About 18 cm. Diameter.—About 4 mm. Internode length.—About 2 cm. Strength.—Moderately strong.

Aspect.—Initially upright to outwardly spreading.

Texture.—Pubescent.

Color.—Close to 144A to 144B.

Foliage description:

Arrangement.—Before flowering, alternate, simple; after flowering, opposite, simple.

Length.—About 5.1 cm.

Width.—About 3 cm.

Shape.—Spatulate.

Apex.—Obtuse.

Base.—Attenuate.

Margin.—Entire.

Texture, upper and lower surfaces.—Pubescent; leathery.

Venation pattern.—Pinnate; arcuate.

Color.—Developing leaves, upper surface: Close to 146B. Developing leaves, lower surface: Close to 146C. Fully expanded leaves, upper surface: Close to 146A; venation, close to 146A. Fully expanded leaves, lower surface: Close to 146B; venation, close to 146B.

Petioles.—Length: About 4 mm. Diameter: About 4 mm. Texture, upper and lower surfaces: Pubescent. Color, upper surface: Close to 146A. Color, lower surface: Close to 146B.

Flower description:

Flower arrangement and habit.—Single salverform flowers arising from leaf axils; freely flowering habit with usually about 25 to 30 open flowers and flower buds per plant; flowers face mostly upright to outwardly.

Fragrance.—None detected.

Natural flowering season.—Plants flower continuously during the spring and summer in Germany; early flowering habit, plants typically beginning flowering about nine weeks after planting.

Flower longevity.—Individual flowers last about five to seven days on the plant; flowers persistent.

Flower diameter.—About 5 cm.

Flower length (height).—About 4 cm.

Flower throat diameter.—About 1.2 cm.

Flower tube length.—About 2.7 cm.

Flower tube diameter.—About 3 mm.

Flower bud.—Length: About 3.5 cm. Diameter: About 6 mm. Shape: Ovoid. Color: Close to 187A to 187B.

Corolla.—Arrangement: Five petals fused at the base and opening into a flared trumpet. Petal lobe length (from throat): About 3 cm. Petal lobe width: About 2.2 cm. Petal shape: Roughly spatulate. Petal apex: Rounded. Petal margin: Entire. Petal texture, upper and lower surfaces: Rippled, glabrous. Throat texture: Rippled, glabrous. Tube texture: Rippled, pubescent. Color: Petal lobe, when opening, upper and lower surfaces: Close to 85B. Petal lobe, fully opened, upper surface: Close to 85C; venation, close to 83A. Petal lobe, fully opened, lower surface: Close to 85C; venation, close to 79B. Flower throat: Close to 79A; venation, close to 79A. Flower tube: Close to 79B; venation, close to 79B.

Calyx.—Arrangement: Five sepals fused at the base forming a star-shaped calyx. Sepal length: About 2.2 cm. Sepal width: About 6.5 mm. Sepal shape: Oblong. Sepal apex: Rounded. Sepal margin: Entire. Sepal texture, upper and lower surfaces: Smooth. Color,

immature and mature, upper surface: Close to 146A. Color, immature and mature, lower surface: Close to 146B.

5

Peduncles.—Length: About 3.6 cm. Diameter: About 1.5 mm. Strength: Moderately strong. Texture: 5 Smooth. Color: Close to 144A.

Reproductive organs.—Stamens: Quantity: Five per flower. Filament length: About 1 cm. Filament color: Close to 148D. Anther shape: Ovate. Anther length: About 1 mm. Anther color: Close to 87B. Pollen amount: Abundant. Pollen color: Close to 97A. Pistils: Quantity: One per flower. Pistil length: About 2 cm. Style length: About 1.8 cm. Style color: Close to 145B. Stigma shape: Rounded. Stigma color: Close to

144A. Ovary color: Close to 144B. Seeds and fruits: Seed and fruit development have not been observed on plants of the new *Petunia*.

Garden performance: Plants of the new *Petunia* have been observed to have good garden performance and tolerate wind, rain and temperatures ranging from about 5° C. to about 40° C.

6

Pathogen/pest resistance: Plants of the new *Petunia* have not been observed to be resistant to pathogens and pests common to *Petunia*.

It is claimed:

1. A new and distinct *Petunia* plant named 'Duepotpiblic' as illustrated and described.

* * * * :

