



US00PP23839P2

(12) **United States Plant Patent**  
**Arts**

(10) **Patent No.:** **US PP23,839 P2**  
(45) **Date of Patent:** **Aug. 20, 2013**

(54) **HYDRANGEA PLANT NAMED**  
**'AGRIHYDRADERTIEN'**

(50) Latin Name: *Hydrangea hybrida*  
Varietal Denomination: **Agrihydradertien**

(75) Inventor: **Niels Arts**, Aalsmeer (NL)

(73) Assignee: **Agriom**, De Kwakel (NL)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 98 days.

(21) Appl. No.: **13/200,606**

(22) Filed: **Sep. 24, 2011**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./250**

(58) **Field of Classification Search**  
USPC ..... **Plt./250**  
See application file for complete search history.

*Primary Examiner* — June Hwu

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Hydrangea* plant named 'Agrihydradertien', characterized by its upright and somewhat outwardly spreading plant habit; vigorous growth habit; strong and sturdy stems; freely flowering habit with numerous sterile flowers per inflorescence; large inflorescences with sterile flowers that are initially light green in color and with development becoming red purple in color; and short production time as a cooling treatment is not required for flower development.

**2 Drawing Sheets**

**1**

Botanical designation: *Hydrangea hybrida*.  
Cultivar denomination: 'AGRIHYDRADERTIEN'.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar of *Hydrangea* plant, botanically known as *Hydrangea hybrida* and hereinafter referred to by the name 'Agrihydradertien'.

The new *Hydrangea* plant is a product of a planned breeding program conducted by the Inventor in Aalsmeer, The Netherlands. The objective of the breeding program was to create new freely branching *Hydrangea* plants with strong and sturdy stems, large inflorescences, attractive flower color and not requirement for a cooling treatment for flower development.

The new *Hydrangea* plant originated from a cross-pollination made by the Inventor on Apr. 20, 2007 in De Kwakel, The Netherlands, of a proprietary selection of *Hydrangea hybrida* identified as code number 200105-02, not patented, as the female, or seed, parent with a proprietary selection of *Hydrangea hybrida* identified as code number 201213-01, not patented, as the male, or pollen, parent. The new *Hydrangea* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Glandorf, Germany on Mar. 15, 2009.

Asexual reproduction of the new *Hydrangea* plant by vegetative cuttings in a controlled environment in Glandorf, Germany since Mar. 15, 2010 has shown that the unique features of this new *Hydrangea* plant are stable and reproduced true to type in successive generations.

**SUMMARY OF THE INVENTION**

Plants of the new *Hydrangea* have not been observed under all possible environmental conditions and cultural practices. The phenotype may vary somewhat with variations in envi-

**2**

ronmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Agrihydradertien'. These characteristics in combination distinguish 'Agrihydradertien' as a new and distinct *Hydrangea* plant:

1. Upright and somewhat outwardly spreading plant habit.
2. Vigorous growth habit.
3. Strong and sturdy stems.
4. Freely flowering habit with numerous sterile flowers per inflorescence.
5. Large inflorescences with sterile flowers that are initially light green in color and with development becoming red purple in color.
6. Short production time as a cooling treatment is not required for flower development.

Plants of the new *Hydrangea* can be compared to plants of the female parent selection. Plants of the new *Hydrangea* differ from plants of female parent selection primarily in sterile flower color as plants of the female parent selection have darker-colored sterile flowers than plants of the new *Hydrangea*. In addition, plants of the new *Hydrangea* have larger inflorescences than plants of the female parent selection.

Plants of the new *Hydrangea* can be compared to plants of the male parent selection. Plants of the new *Hydrangea* differ from plants of male parent selection primarily in sterile flower color as plants of the male parent selection have purple-colored sterile flowers.

Plants of the new *Hydrangea* can be compared to plants of the *Hydrangea hybrida* 'Agrihydradrie', disclosed in U.S. Plant Pat. No. 19,726. Plants of the new *Hydrangea* differ from plants of 'Agrihydradrie' in the following characteristics:

1. Plants of the new *Hydrangea* are taller and more vigorous than plants of 'Agrihydradrie'.

2. Plants of the new *Hydrangea* have darker-colored sterile flowers than plants of 'Agrihydradrie'.

## BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the unique appearance of the new *Hydrangea* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ from the color values cited in the detailed botanical description which accurately describe the colors of the new *Hydrangea* plant.

The photograph on the first sheet comprises a side perspective view of a typical flowering plant of 'Agrihydradertien' with developing inflorescences grown in a container.

The photograph on the second sheet is a close-up view of a typical flowering plant of 'Agrihydradertien' with fully developed inflorescences.

## DETAILED BOTANICAL DESCRIPTION

Plants used in the aforementioned photographs and in the following description were grown during the winter in 15-cm containers in a glass-covered greenhouse in De Kwakel, The Netherlands and under environmental conditions and cultural practices which closely approximate commercial *Hydrangea* production conditions. During the production of the plants, day and night temperatures averaged 17° C. Plants of the new *Hydrangea* were one year old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical description: *Hydrangea hybrida* 'Agrihydradertien'.

## Parentage:

*Female, or seed, parent.*—Proprietary selection of *Hydrangea hybrida* identified as code number 200105-02, not patented.

*Male, or pollen, parent.*—Proprietary selection of *Hydrangea hybrida* identified as code number 201213-01, not patented.

## Propagation:

*Type cutting.*—By vegetative cuttings.

*Time to initiate roots, summer.*—About two weeks at temperatures of 23° C.

*Time to initiate roots, winter.*—About 18 days at temperatures of 18° C.

*Time to produce a rooted young plant, summer.*—About four weeks at temperatures of 23° C.

*Time to produce a rooted young plant, winter.*—About five weeks at temperatures of 18° C.

*Root description.*—Thick; whitish brown in color.

*Rooting habit.*—Freely branching; dense.

## Plant description:

*Plant and growth habit.*—Upright to somewhat outwardly spreading plant habit; rounded in shape; strong and sturdy stems; rapid growth rate and vigorous growth habit.

*Plant height.*—About 30 cm to 40 cm.

*Plant diameter or area of spread.*—About 40 cm.

## Lateral branch description:

*Branching habit.*—Freely branching habit with about six lateral branches per plant.

*Length.*—About 30 cm to 35 cm.

*Diameter.*—About 6 mm.

*Internode length.*—About 3.5 cm.

*Stem texture.*—Smooth, glabrous.

*Strength.*—Strong and sturdy.

*Color.*—Close to 144A; at the nodes, close to 187A; lenticels, close to 187A.

## Foliage description:

*Arrangement.*—Opposite, simple.

*Length.*—About 9 cm.

*Width.*—About 6.5 cm.

*Shape.*—Ovate.

*Apex.*—Acute to acuminate.

*Base.*—Obtuse.

*Margin.*—Dentate to serrate.

*Texture, upper surface.*—Smooth to rugose, glabrous.

*Texture, lower surface.*—Rugose, glabrous.

*Venation pattern.*—Pinnate.

*Color.*—Developing and fully expanded leaves, upper surface: Close to 139A; venation, close to 146C. Developing and fully expanded leaves, lower surface: Close to 137C; venation, close to 146D.

*Petiole.*—Length: About 1.5 cm. Diameter: About 3 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 146C.

## Flower description:

*Flower type and habit.*—Single rounded sterile and star-shaped fertile flowers arranged on terminal panicles; panicles globular in shape; flowers face upright to outwardly.

*Fragrance.*—None detected.

*Natural flowering season.*—Short production time as a cooling treatment is not required for flower development; continuous flowering during the summer in Northern Europe.

*Flower longevity.*—Sterile flowers last about four months on the plant, sterile flowers persistent; fertile flowers last about four weeks on the plant, fertile flowers not persistent.

*Quantity of flowers.*—Freely flowering habit; about 40 sterile flowers per panicle and about 15 to 20 fertile flowers per panicle.

*Panicle height.*—About 8 cm.

*Panicle diameter.*—About 15 cm.

*Sterile flower diameter.*—About 3.5 cm.

*Sterile flower depth (height).*—About 1 cm.

*Fertile flower diameter.*—About 5 mm.

*Fertile flower depth (height).*—About 5 mm.

*Petals, sterile flowers only, fertile flowers without petals.*—Quantity and arrangement: About four in a single whorl. Length: About 3.5 mm. Width: About 2 mm. Shape: Ovate. Apex: Acute. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening and fully opened, upper surface: Close to 57C; color becoming closer to 66B. When opening and fully opened, lower surface: Close to 57D.

*Sepals, sterile flowers.*—Quantity and arrangement: About four in a single whorl. Length: About 1.8 cm. Width: About 1.9 cm. Shape: Roughly deltoid. Apex: Retuse. Base: Cuneate. Margin: Entire. Texture, upper surface: Smooth, glabrous. Texture, lower surface: Smooth to rugose, glabrous. Color: When opening, upper surface: Close to 144A to 144B; with development, towards the apex, close to 58A. When opening, lower surface: Close to 144A to 144B. Fully

opened, upper surface: Close to 58A. Fully opened, lower surface: Close to 64D.

*Sepals, fertile flowers*.—Quantity and arrangement: About four in a single whorl. Length: About 3.5 mm. Width: About 2 mm. Shape: Ovate. Apex: Acute. 5  
Base: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper surface: Close to 63C. Color, lower surface: Close to 63B.

*Pedicels, sterile flowers*.—Length: About 2 cm. Diameter: About 1.5 mm. Strength: Strong. Texture: Smooth, glabrous. Color: Close to 144A overlain with close to 64D. 10

*Pedicels, fertile flowers*.—Length: About 4 mm. Diameter: About 1 mm. Strength: Moderately strong. Texture: Smooth, glabrous. Color: Close to 62C. 15

*Reproductive organs, fertile flowers only; sterile flowers without reproductive organs*.—Stamens: Quantity per flower: About eight. Anther shape: Conical. Anther length: About 1 mm. Anther color: Close to 20

145D. Pollen amount: Abundant. Pollen color: Close to 155D. Pistils: Pistil quantity per flower: About three. Pistil length: About 1 mm. Stigma shape: Oval. Stigma color: Close to 155C. Style length: About 1 mm. Style color: Close to 65D. Ovary color: Close to 65D.

*Seeds*.—Length: About 0.5 mm. Diameter: About 0.1 mm. Color: Close to 200C.

Disease/pest resistance: Under commercial production conditions, plants of the new *Hydrangea* have not been observed to be resistant to pathogens or pests common to *Hydrangea*.

Temperature tolerance: Plants of the new *Hydrangea* have been shown to be tolerant to temperatures ranging from about 3° C. to about 38° C.

It is claimed:

1. A new and distinct *Hydrangea* plant named 'Agrihydradertien' as illustrated and described.

\* \* \* \* \*



