



(12) **United States Plant Patent**
Dümmen

(10) **Patent No.:** **US PP23,927 P2**
(45) **Date of Patent:** **Sep. 24, 2013**

(54) **NEW GUINEA *IMPATIENS* PLANT NAMED
'DUEPETBS11'**

(50) Latin Name: *Impatiens hawkeri*
Varietal Denomination: **Duepetbs11**

(75) Inventor: **Tobias Dümmen**, Rheinberg (DE)

(73) Assignee: **Capital Green Investments Ltd.**, Grand
Cayman (KY)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 108 days.

(21) Appl. No.: **13/317,913**

(22) Filed: **Oct. 31, 2011**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./318.1**

(58) **Field of Classification Search**
USPC Plt./318.1, 318.5
See application file for complete search history.

Primary Examiner — June Hwu

(74) Attorney, Agent, or Firm — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Impatiens* plant named
'Duepetbs11' characterized by its upright and outwardly
spreading growth habit; mounded plant habit; freely branch-
ing habit; moderately vigorous growth habit; dark green-
colored leaves; freely and early flowering habit; red purple
and red bi-colored flowers; and good garden performance.

1 Drawing Sheet

1

Botanical designation: *Impatiens hawkeri*.
Cultivar denomination: 'DUEPETBS11'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar
of New Guinea *Impatiens* plant, botanically known as *Impa-
tiens hawkeri* and hereinafter referred to by the name
'Duepetbs11'.

The new *Impatiens* plant is a product of a planned breeding
program conducted by the Inventor in Rheinberg, Germany.
The objective of the breeding program is to create new freely
branching and uniform New Guinea *Impatiens* plants with
large and attractive flowers.

The new *Impatiens* plant originated from a cross-pollina-
tion made by the Inventor in July, 2007 in Rheinberg, Ger-
many of a proprietary selection of *Impatiens hawkeri* identi-
fied as code number N03-0035-006, not patented, as the
female, or seed, parent with a proprietary selection of *Impa-
tiens hawkeri* identified as code number F-16-02, not pat-
ented, as the male, or pollen, parent. The new *Impatiens* plant
was discovered and selected by the Inventor as a single flow-
ering plant from within the progeny of the stated cross-polli-
nation in a controlled greenhouse environment in Rheinberg,
Germany in May, 2010.

Asexual reproduction of the new *Impatiens* plant by termi-
nal cuttings in a controlled greenhouse environment in Rhei-
nberg, Germany since June, 2010 has shown that the unique
features of this new *Impatiens* plant are stable and reproduced
true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Impatiens* have not been observed under
all possible environmental conditions and cultural practices.
The phenotype may vary somewhat with variations in envi-
ronmental conditions such as temperature, daylight and light
intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are
determined to be the unique characteristics of 'Duepetbs11'.

2

These characteristics in combination distinguish
'Duepetbs11' as a new and distinct *Impatiens* plant:

1. Upright and outwardly spreading growth habit;
mounded plant habit.
2. Freely branching habit.
3. Moderately vigorous growth habit.
4. Dark green-colored leaves.
5. Freely and early flowering habit.
6. Red purple and red bi-colored flowers.
7. Good garden performance.

Plants of the new *Impatiens* can be compared to plants of
the female parent selection. Plants of the new *Impatiens* differ
primarily from plants of the female parent selection in the
following characteristics:

1. Plants of the new *Impatiens* are more vigorous than
plants of the female parent selection.
2. Plants of the new *Impatiens* and the female parent selec-
tion differ in flower color.

Plants of the new *Impatiens* can be compared to plants of
the male parent selection. Plants of the new *Impatiens* differ
primarily from plants of the male parent selection in the
following characteristics:

1. Plants of the new *Impatiens* are more vigorous than
plants of the male parent selection.
2. Plants of the new *Impatiens* and the male parent selection
differ in flower color.

Plants of the new *Impatiens* can be compared to plants of
New Guinea *Impatiens* 'Kibahia', disclosed in U.S. Plant Pat.
No. 18,442. In side-by-side comparisons conducted in Rhei-
nberg, Germany, plants of the new *Impatiens* differed prima-
rily from plants of 'Kibahia' in the following characteristics:

1. Plants of the new *Impatiens* were more vigorous than
plants of 'Kibahia'.
2. Plants of the new *Impatiens* and 'Kibahia' differed in
flower color as plants of 'Kibahia' had violet purple and
red purple-colored flowers.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the over-
all appearance of the new *Impatiens* plant showing the colors

as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Impatiens* plant.

The photograph comprises a side perspective view of a typical flowering plant of 'Duepetbs11' grown in a container.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations, measurements and values describe plants grown in 10.5-cm containers during the summer in a glass-covered greenhouse in Rheinberg, Germany and under cultural practices typically used in commercial New Guinea *Impatiens* production. During the production of the plants, night temperatures averaged 18° C. and light levels averaged 4,500 lux. Plants were pinched one time about three weeks after planting and were 16 weeks old when the photograph and description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* 'Duepetbs11'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Impatiens hawkeri* identified as code number N03-0035-006, not patented.

Male, or pollen, parent.—Proprietary selection of *Impatiens hawkeri* identified as code number F-16-02, not patented.

Propagation:

Type.—By terminal cuttings.

Time to initiate roots, summer.—About five days at temperatures of about 20° C.

Time to initiate roots, winter.—About seven days at temperatures of about 20° C.

Time to produce a rooted young plant, summer.—About three weeks at temperatures of about 20° C.

Time to produce a rooted young plant, winter.—About four weeks at temperatures of about 20° C.

Root description.—Fine, fibrous; white in color.

Rooting habit.—Freely branching; dense.

Plant description:

Plant and growth habit.—Upright to outwardly spreading growth habit; mounded plant habit; freely branching habit with about seven lateral branches; moderately vigorous growth habit.

Plant height.—About 16 cm.

Plant diameter.—About 18 cm.

Lateral branch description:

Length.—About 14 cm.

Diameter.—About 6 mm.

Internode length.—About 4 cm.

Strength.—Strong.

Aspect.—Initially upright to outwardly spreading.

Texture.—Smooth, glabrous.

Color.—Close to 147C.

Foliage description:

Arrangement.—Opposite or in whorls; simple.

Length.—About 10 cm.

Width.—About 4 cm.

Shape.—Ovate.

Apex.—Apiculate.

Base.—Obtuse.

Margin.—Serrulate with ciliation.

Texture, upper and lower surfaces.—Smooth, glabrous; leathery.

Venation pattern.—Pinnate; arcuate.

Color.—Developing leaves, upper surface: Close to 139A; venation, close to 59A. Developing leaves, lower surface: Close to 59A; venation, close to 59A.

Fully expanded leaves, upper surface.—Close to 139A tinted with close to 59A; venation, close to 59A.

Fully expanded leaves, lower surface.—Close to 59A; venation, close to 59A.

Petiole length.—About 2.2 cm.

Petiole diameter.—About 4 mm.

Petiole texture, upper and lower surfaces.—Smooth, glabrous.

Petiole color, upper and lower surfaces.—Close to 60C.

Flower description:

Flower type and flowering habit.—Single rounded and flat axillary flowers; freely flowering habit, typically about six open flowers and flower buds per lateral branch; flowers positioned above and beyond the foliar plane, flowers typically face upright or outwardly.

Flower longevity.—Flowers typically last about five to six days under greenhouse conditions; petals self-cleaning, gynoecium persistent.

Fragrance.—None detected.

Natural flowering season.—Year-round under greenhouse conditions; in the garden, flowering from spring until fall in Germany; early flowering habit, plants typically begin flowering about eight weeks after planting.

Flower buds.—Length: About 1.8 cm. Diameter: About 1 cm. Shape: Ovate. Color: Close to 45C and 53A.

Flower size.—Length: About 6 cm. Diameter: About 6.2 cm. Depth: About 3 cm.

Petals.—Quantity and arrangement: Five per flower in a single whorl. Length: About 3 cm. Width: About 3.6 cm. Shape: Obcordate. Apex: Emarginate; rounded. Base: Acute. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous; satiny. Color: When opening, upper surface: Close to 74B; center and star pattern, close to 46B. When opening, lower surface: Close to 47B. Fully opened, upper surface: Close to 74A; center and star pattern, close to 46B; colors becoming closer to 74D and 46D with development. Fully opened, lower surface: Close to 47C.

Sepals.—Quantity and arrangement: Three in a single whorl; one modified into an elongated spur. Length: About 1 cm. Width: About 5 mm. Shape: Oval. Apex: Acute. Base: Obtuse. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 147A. Spur length: About 4.7 cm. Spur diameter: At flower, about 3 mm; at apex, less than 1 mm. Spur texture: Smooth, glabrous. Spur color: Close to 145C and 67D.

Peduncles.—Length: About 5 cm. Diameter: About 2 mm. Angle: Upright to outward. Strength: Moderately strong; flexible. Texture: Smooth, glabrous. Color: Close to 146D.

Reproductive organs.—Stamens: Quantity: Five fused at anthers; filaments free. Anther length: About 7 mm. Anther shape: Oval. Anther color: Close to 10B and 63A. Pollen amount: Abundant. Pollen color: Close to 11C. Pistils: Quantity per flower: One. Pistil length:

About 7 mm. Stigma shape: Crested. Stigma color:
Close to 59A. Style color: Close to 61A. Ovary color:
Close to 59A.
Seeds and fruits.—Seed and fruit production has not
been observed on plants of the new *Impatiens*.
Disease/pest resistance: Plants of the new *Impatiens* have not
been observed to be resistant to pathogens and pests com-
mon to New Guinea *Impatiens*.

Garden performance: Plants of the new *Impatiens* have been
observed to have good garden performance and tolerate
temperatures ranging from about 5° C. to about 40° C.
It is claimed:
1. A new and distinct *Impatiens* plant named ‘Duepetbs11’
as illustrated and described.

* * * * *

