



US00PP23954P2

(12) **United States Plant Patent**
Kordes

(10) **Patent No.:** **US PP23,954 P2**
(45) **Date of Patent:** **Oct. 8, 2013**

(54) **MINATURE ROSE PLANT NAMED**
‘KORPOT001’

(50) Latin Name: *Rosa hybrida*
Varietal Denomination: **KORpot001**

(75) Inventor: **Tim-Hermann Kordes**, Klein
Offenseth-Sparrieshoop (DE)

(73) Assignee: **W. Kordes’ Söhne Rosenschulen**
GmbH & Co KG,
Offenseth-Sparrieshoop (DE)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 89 days.

(21) Appl. No.: **13/385,489**

(22) Filed: **Feb. 21, 2012**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./120**

(58) **Field of Classification Search**
USPC Plt./120
See application file for complete search history.

(56) **References Cited**

PUBLICATIONS

Pluto UPOV PBR QZ citation for ‘KORpot001’ Feb. 22, 2011.*

* cited by examiner

Primary Examiner — Wendy C Haas

(57) **ABSTRACT**

A new and distinct variety of rose with long lasting, novel
cream apricot flowers, and attractive foliage with good dis-
ease resistance. It exhibits above average vigor with abundant
flowers. The new variety propagates well from cuttings and
by grafting. This new and distinct variety has shown to be
uniform and stable in the resulting generations from asexual
propagation.

1 Drawing Sheet

1

Latin name of genus and species: The botanical classifica-
tion of the new rose plant is *Rosa hybrida*.

Variety denomination: The denomination of the new vari-
ety is ‘KORpot001’.

CROSS REFERENCES AND FEDERAL R&D
STATEMENT

There are no cross referenced or related applications. This
variety was developed without the aid of any research grant.

The new variety of rose plant of the present invention
originated from a controlled crossing in a breeding program
of two distinct parents during the summer of 2006. The cross-
ing was between an ‘un-named seedling’, the seed parent, and
another ‘un-named seedling’, the pollen parent, from the
same inventor.

The resulting seeds were planted during the following win-
ter. The resulting seedlings were evaluated and exhibited
distinctive physical and biological characteristics. The new
rose plant was selected as a single plant from the seedling
beds due to its superior characteristics and asexually propa-
gated for further evaluation. This new and distinctive rose
variety is named ‘KORpot001’.

SUMMARY OF THE INVENTION

The new rose plant may be distinguished from its seed
parent, an ‘un-named seedling’, by the following combina-
tion of characteristics:

1. ‘KORpot001’ has cream apricot cup shaped flowers,
whereas the ‘un-named seedling’ has cream white flow-
ers with high centered buds.
2. ‘KORpot001’ has a moderately compact growth habit,
whereas the ‘un-named seedling’ has an upright habit.

2

The new rose plant may be distinguished from its pollen
parent, an ‘un-named seedling’, by the following combina-
tion of characteristics:

1. ‘KORpot001’ has cream apricot flowers, whereas the
‘un-named seedling’ has copper yellow flowers.
2. ‘KORpot001’ has large flowers, whereas the ‘un-named
seedling’ has medium sized flowers.

The objective of the hybridization was to create a new and
distinct rose plant with unique qualities, such as:

1. Uniform growth and flowering;
2. Abundant attractive, recurrent flowers;
3. Attractive and abundant foliage; and
3. Resistance to diseases encountered in landscapes and
gardens.

This combination of qualities is not present in prior rose
cultivars known to the inventor. These objectives have been
substantially achieved and in that distinguish ‘KORpot001’
from all other varieties of which I am aware.

As part of a rose development program, Tim-Hermann
Kordes germinated seeds from the aforementioned hybridiza-
tion and conducted evaluations and observations on the
resulting seedlings in a controlled environment in Offenseth-
Sparrieshoop, Germany. The resulting seedlings exhibited
distinctive physical and biological characteristics. The new
rose plant ‘KORpot001’ was selected in May, 2007 from the
seedling beds to be asexually propagated for further evalua-
tion. The first asexual propagation of ‘KORpot001’ was done
by rooting cuttings in June, 2007 at the inventor’s nursery in
Offenseth-Sparrieshoop, Germany.

This initial and other subsequent propagations conducted
in controlled environments demonstrate that ‘KORpot001’
reproduces true to type in successive generations of asexual
reproduction.

BRIEF DESCRIPTION OF THE DRAWING

The accompanying color drawing shows as true as is rea-
sonably possible to obtain in color photographs of this type,

the typical characteristics of the buds, sepals, reproductive organs, flowers, petals, leaves, prickles, and stems of ‘KORpot001’.

DETAILED BOTANICAL DESCRIPTION

The following is a description of ‘KORpot001’, as observed growing in October, 2011 in a nursery in Jackson County, Oreg. on plants 6 months of age. Color references are made using The Royal Horticultural Society (London, England) Colour Chart, 2001 except where common terms of color are used.

For a comparison, several physical characteristics of the rose variety ‘KORpedia’, a rose variety from the same inventor described and illustrated in U.S. Plant Pat. No. 19,638 and issued on Jan. 20, 2009 are compared to ‘KORpot001’ in Chart 1.

CHART 1

Characteristic	‘KORpot001’	‘KORpedia’
General tonality	Yellow-Orange Group 20C	Yellow Group 2D
Pistil count	25-30	Approximately 90
Petal count	70-85	30-35

Parents: Seed Parent: ‘Un-named seedling’. Pollen Parent: ‘Un-named seedling’.

Classification:

Botanical classification.—*Rosa hybrida* ‘KORpot001’.

Commercial classification.—Miniature rose.

FLOWER AND FLOWER BUD

Blooming habit: Recurrent.

Flower bud:

Size.—Upon opening, 20 mm in length from base of receptacle to distal end of bud and 14 mm diameter at its widest point.

Bud form.—Short. Pointed ovoid.

Bud color.—As sepals first unfold, bud color is Yellow-Green Group 149D. When ¼ open, the upper surface of petals is Yellow-White Group 158D with a basal zone of Yellow Group 9C, and the lower surface is Yellow-Orange Group 18D. Guard petals have intonations of Yellow-Green Group 145C.

Sepals.—Color: Upper surface: Yellow-Green Group 144A. Lower surface: Yellow-Green Group 144A. Size: Average 15-35 mm (l)×7-9 mm (w). Shape: Strong foliaceous appendages on three of the five sepals. Apex: Three sepals are cirrose and two are apiculate. Base: Flat at union with receptacle. Quantity: Five. Surface texture: Upper side: Heavily pubescent. Lower surface: Light pubescence with stipitate glands. Margins: With stipitate glands.

Receptacle:

Surface.—Smooth.

Color.—Yellow-Green Group 144B.

Shape.—Funnel.

Size.—8 mm (h)×7 mm (w).

Peduncle:

Surface.—With fine hairs and stipitate glands.

Length.—60 to 80 mm average length.

Diameter.—2 to 3 mm average diameter.

Color.—Yellow-Green Group 144B.

Strength.—Strong.

Borne.—Singularly.

Flower bloom:

Fragrance.—Very light fruit and citrus scent.

Duration.—On the plant 12 to 14 days. Long lasting. As a cut flower, 6 to 7 days. Senesced petals drop away cleanly.

Size.—Large for a miniature rose. When open, the average flower diameter is 80 mm and the average flower height is 35 mm.

Form.—Shape of flower when viewed from the side: Upon opening, upper part: Flattened convex. Upon opening, lower part: Flattened convex. Open flower, upper part: Flat. Open flower, lower part: Concave.

Color:

Upon opening, petals.—Outermost petals: Outer Side: Yellow-White Group 158D. Inner Side: Yellow-White Group 158D with a basal zone of Yellow Group 10B. Innermost petals: Outer Side: Yellow-Orange Group 23D. Inner Side: Yellow-Orange Group 22D.

Upon opening, basal petal spots.—Basal petal spot, outermost petals: Outer Side: None observed. Inner Side: No distinct basal spot. Basal zone is Yellow Group 12A. Basal petal spot, innermost petals: Outer Side: Yellow Group 12C. Inner Side: Yellow Group 12B.

After opening, petals.—Outermost petals: Outer Side: White Group 155B. Inner Side: Yellow-White Group 158C with basal zone of Yellow Group 8A. Innermost petals: Outer Side: Yellow-Orange Group 20C. Inner Side: Yellow-Orange Group 20C with intonations of Yellow Group 9B in the basal zone.

After opening, basal petal spots.—Basal petal spot, outermost petals: Outer Side: Yellow Group 3D. Inner Side: Yellow Group 3D. Basal petal spot, innermost petals: Outer Side: Yellow Group 9C. Inner Side: Yellow Group 9C.

General tonality: On open flower Yellow-Orange Group 20C. No change in the general tonality at the end of the fourth day. Afterwards, general tonality is Yellow-Orange Group 18C.

Petals:

Petal count.—Very double.

Average range.—Approximately 70 to 85 petals under normal conditions.

Petal reflex.—Petals reflex somewhat.

Petal edge.—Entire.

Petal shape.—Obovate. Apex shape is round. Shape of base is cuneate.

Petal size.—30-40 mm (l)×30 mm (w).

Thickness.—Thick.

Petal arrangement.—Not Formal.

Petaloids:

Petaloid count.—Average of 2-5 per flower.

Petaloid size.—Petaloids are 10-15 mm (l)×3-4 mm (w).

Petaloid color.—Color of inner side is Yellow-Orange Group 18D. Color of outer side is Yellow-Orange Group 16D.

Petaloid texture.—Thick.

Margins.—Reflexed and ruffled.

Petaloid shape.—Most commonly oblanceolate, with some petaloids highly irregular. Apex: Obtuse. Base: Pointed.

Reproductive organs:

Pistils.—Approximately 25-30 present. Stigmas: Location: Slightly superior in position to anthers. Color:

Yellow Group 4C. Styles: Length: About 8 mm long. Color: Green-Yellow Group 1D with intonations of Red Group 46C.

Stamens.—Approximately 20 on average and regularly arranged. Anthers: Size: Average 0.5 mm long. Pollen: Generally present. Color: Greyed-Orange Group 163A. Filaments: Color: Green-Yellow Group 1C. Length: 4 mm.

THE PLANT

Growth: Above average vigor.

Plant habit: Compact upright habit. When grown as a 13 cm pot plant, the average plant height is 25 cm and the average plant width is 20 cm.

Blooming: Floriferous.

Stems:

Stem color.—Young wood: Yellow-Green Group 144A.

Older wood: Yellow-Green Group 144A.

Stem surface.—Young wood: Smooth. Older wood: Smooth.

Prickles: Present.

Incidence.—Average of 3 per each 10 cm of stem.

Size.—Average length: 6 mm.

Color.—Immature prickles: Green-Yellow Group 161. Prickles not observed on mature wood.

Shape.—Linear.

Anthocyanin.—On young prickles, Greyed-Red Group 182C.

Leaves and leaflets: Normally 5 leaflets on normal leaves in middle of the stem.

Venation pattern.—Pyramidal net pattern.

Leaf size.—110 mm (l)×80 mm (w).

Abundance.—Average.

Texture.—Thin. Upper side of leaflet: Semi-glossy. Under side of leaflet: Matte.

Color, mature foliage.—Upper Leaf Surface: Green Group 137A. Lower Leaf Surface: Green Group 138B.

Color, juvenile foliage.—Upper Leaf Surface: Yellow-Green Group 144B. Lower Leaf Surface: Yellow-Green Group 144B.

Anthocyanin intonation.—Present. Intonations of Greyed-Purple Group 183B present on leaflet margins and upper leaf surface.

Stipules:

Size.—8 mm long, 3 mm from distal tip to distal tip.

Stipule color.—Yellow-Green Group 144A.

Margins.—With stipitate glands.

Shape.—Apex: Apiculate. Base: Winged.

Petiole:

Length.—Average 12 mm.

Diameter.—Average 1 mm.

Petiole color.—Yellow-Green Group 144A.

Underneath.—With small, fine prickles and stipitate glands.

Margins.—With stipitate glands.

Petiole rachis:

Length.—Average 18-20 mm.

Diameter.—Average 1 mm.

Color.—Yellow-Green Group 144A.

Margins.—With stipitate glands.

Underneath.—With stipitate glands and a few small prickles.

Leaflets:

Size.—Average size of the terminal leaflet is 50 mm (l)×30 mm (w).

Shape.—Ovate. Base: Obtuse. Apex: Acute.

Margins.—Serrated.

Surface.—Upper surface: Moderately glossy. Lower surface: Matte.

Texture.—Thin.

Arrangement.—Odd pinnate.

Venation.—Reticulate.

Hips/seed formation: Not observed.

Winter hardiness: Unknown.

Disease resistance: Good resistance to powdery mildew (*Sphaerotheca pannosa*) and Botrytis (*Botrytis cinerea*) diseases under normal growing conditions.

I claim:

1. A new and distinct variety of rose plant, as described and illustrated herein.

* * * * *

