



US00PP23999P2

(12) **United States Plant Patent**  
**Bergman**

(10) **Patent No.:** US PP23,999 P2  
(45) **Date of Patent:** Oct. 22, 2013

- (54) **CHRYSANTHEMUM PLANT NAMED  
'CIDZ0021'**
- (50) Latin Name: *Chrysanthemum×morifolium*  
Varietal Denomination: **CIDZ0021**
- (75) Inventor: **Wendy R. Bergman**, Gilroy, CA (US)
- (73) Assignee: **Syngenta Crop Protection AG**, Basel (CH)
- (\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 184 days.
- (21) Appl. No.: **13/317,426**
- (22) Filed: **Oct. 18, 2011**

- (51) **Int. Cl.**  
*A01H 5/00* (2006.01)
- (52) **U.S. Cl.**  
USPC ..... **Plt./296; Plt./286**
- (58) **Field of Classification Search**  
USPC ..... Plt./286, 296  
See application file for complete search history.

*Primary Examiner* — Susan McCormick Ewoldt  
*(74) Attorney, Agent, or Firm* — Joshua L. Price

(57) **ABSTRACT**

A new *Chrysanthemum* plant named 'CIDZ0021' particularly distinguished by the small sized daisy-type inflorescences with greyed orange color, medium green foliage, very compact plant growth habit and is freely branched.

**1 Drawing Sheet**

**1**

Latin name of the genus and species of the plant claimed:  
*Chrysanthemum×morifolium*.  
Varietal denomination: 'CIDZ0021'.

**BACKGROUND OF THE NEW PLANT**

The present invention comprises a new *Chrysanthemum*, botanically known as *Chrysanthemum×morifolium*, and hereinafter referred to by the variety name 'CIDZ0021'.

'CIDZ0021' is a product of a planned breeding program. The new cultivar has small sized daisy-type inflorescences with greyed orange color, medium green foliage, very compact plant growth habit and is freely branched.

'CIDZ0021' originated from a hybridization made in March 2004 in a controlled breeding environment in Salinas, Calif. The female parent was unknown.

The male parent of 'CIDZ0021' was a proprietary, unpatented plant identified as 'F0079' with smaller size inflorescences, smaller plant habit and faster flowering response time.

'CIDZ0021' was selected as one flowering plant within the progeny of the stated cross in the May 2005 in a controlled environment in Fort Myers, Fla.

The first act of asexual reproduction of 'CIDZ0021' was accomplished when vegetative cuttings were propagated from the initial selection in June 2005 in a controlled environment in Fort Myers, Fla.

**BRIEF SUMMARY OF INVENTION**

Horticultural examination of plants grown from cuttings of the plant initiated in June 2005, and continuing thereafter, has demonstrated that the combination of characteristics as herein disclosed for 'CIDZ0021' are firmly fixed and are retained through successive generations of asexual reproduction.

'CIDZ0021' has not been observed under all possible environmental conditions. The phenotype may vary significantly with variations in environment such as temperature, light intensity and day length.

Plant Breeder's Rights for this cultivar were applied for in Canada on Jul. 29, 2011, #11-7346 and in Brazil on Apr. 8,

**2**

2011, #21806.000105/2011. 'CIDZ0021' has not been made publicly available more than one year prior to the filing of this application.

5 The following traits have been repeatedly observed and are determined to be basic characteristics of the new variety. The combination of these characteristics distinguishes this *Chrysanthemum* as a new and distinct variety.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographic drawing shows typical flower and foliage characteristics of 'CIDZ0021' with colors being as true as possible with an illustration of this type.

15 The photographic drawing shows a flowering plant of the new variety, and a close-up of the flowers.

**DETAILED BOTANICAL DESCRIPTION**

20 The plant descriptions, measurements and aforementioned photographs were taken in Gilroy, Calif. in September 2011 under natural light. These plants were propagated and grown in Nipomo, Calif. and shipped to Gilroy, Calif. for the data collection and photographs. These plants were approximately 9 weeks of age, grown as 4 plants together in a 5 inch bulb pan, under greenhouse trial conditions.

25 Color references are made to The Royal Horticultural Society Colour Chart (R.H.S.) 2001.

**TABLE 1**

**DIFFERENCES BETWEEN THE NEW VARIETY  
'CIDZ0021' AND A SIMILAR VARIETY**

	<b>'CIDZ0021'</b>	<b>'Yostockton' (Not patented)</b>
Plant height:	Taller	Shorter
Inflorescence size:	Larger	Smaller
Flowering response time:	7 weeks	6 weeks

40 Plant:

*Form, growth and habit.*—Herbaceous fleurette pot type, stems upright, freely branching, small foliage.

<i>Plant height.</i> —About 10.0 cm.		<i>Immature inflorescence:</i>
<i>Plant height (inflorescence included).</i> —14.0-15.0 cm.		<i>Diameter.</i> —1.0-2.9 cm.
<i>Plant width.</i> —15.0 cm.		<i>Color of ray florets, upper surface.</i> —RHS 160A ground color heavily overlaid and mottled with RHS 175B to RHS 175C with RHS 178A margins.
<i>Roots:</i>		<i>Lower surface.</i> —RHS 160A ground color lightly overlaid and mottled with a muted hue of RHS 175B to RHS 175C with darker margins.
<i>Number of days to initiate roots.</i> —4 days at about 22 degrees C.	5	<i>Mature inflorescence:</i>
<i>Number of days to produce a rooted cutting.</i> —About 10 days at 22 degrees C.	10	<i>Diameter.</i> —3.2-3.5 cm.
<i>Type.</i> —Fine, fibrous, free branching.	10	<i>Depth.</i> —1.0-1.3 cm.
<i>Color.</i> —RHS N155B but whiter.	15	<i>Total diameter of 'disc'.</i> —1.2-1.3 cm.
<i>Foliage:</i>		<i>Receptacle height.</i> —0.25-0.3 cm.
<i>Arrangement.</i> —Alternate.		<i>Receptacle diameter.</i> —0.35 cm.
<i>Immature, leaf color, upper surface.</i> —Closest to RHS 137A but darker.	15	<i>Ray florets:</i>
<i>Lower surface.</i> —Between RHS 137B and RHS 137C.	20	<i>Average quantity of florets.</i> —Average of 36 in 1-2 whorls.
<i>Mature, leaf color, upper surface.</i> —Closest to RHS 137A but darker.	20	<i>Color of florets, upper surface.</i> —RHS 160A ground color heavily overlaid and mottled with RHS 175B to RHS 175C with RHS 178A margins. Color just very faintly fades as it matures.
<i>Lower surface.</i> —Between RHS 137B and RHS 137C.	25	<i>Lower surface.</i> —RHS 160A ground color lightly overlaid and mottled with a muted hue of RHS 175B to RHS 175C with darker margins.
<i>Length.</i> —3.7-4.0 cm.	25	<i>Length.</i> —1.7-1.8 cm.
<i>Width.</i> —2.4-2.8 cm.	25	<i>Width.</i> —0.45 cm.
<i>Shape.</i> —Elliptical.	25	<i>Shape.</i> —Elliptical.
<i>Base shape.</i> —Attenuate.	25	<i>Apex shape.</i> —Slightly emarginated to praemorse, a few are obtuse.
<i>Apex shape.</i> —Mucronulate.	25	<i>Margin.</i> —Entire.
<i>Margin.</i> —Somewhat palmately lobed, slightly serrate.	25	<i>Texture, upper surface.</i> —Papillose.
<i>Texture, upper surface.</i> —Bifid T-shaped hairs.	25	<i>Lower surface.</i> —Papillose.
<i>Lower surface.</i> —Bifid T-shaped hairs.	25	<i>Disc florets:</i>
<i>Color of veins, upper surface.</i> —Closest to RHS 138B.	30	<i>Average quantity of florets.</i> —About 200.
<i>Color of veins, lower surface.</i> —RHS 138B.	30	<i>Color of florets.</i> —RHS 2C with a RHS 9B apex.
<i>Petiole color.</i> —RHS 138B.	30	<i>Length.</i> —0.45-9.5 cm.
<i>Length.</i> —1.3-1.6 cm.	30	<i>Width.</i> —0.1 cm.
<i>Diameter.</i> —0.2 cm.	30	<i>Shape.</i> —Tubular, elongated.
<i>Texture.</i> —Bifid T-shaped hairs.	30	<i>Apex shape.</i> —Acute, 5 pointed.
<i>Stem:</i>		<i>Phyllaries:</i>
<i>Quantity of main branches per plant.</i> —3-4.	35	<i>Quantity.</i> —Approximately 25.
<i>Color of stem.</i> —RHS 138A but appears lighter due to hairs.	35	<i>Color, upper surface.</i> —RHS 137C.
<i>Length of stem.</i> —5.4-6.5 cm.	35	<i>Lower surface.</i> —Closest to RHS 137A but appears lighter due to hairs.
<i>Diameter.</i> —0.25-0.3 cm.	35	<i>Length.</i> —0.35-0.4 cm.
<i>Length of internodes.</i> —0.4-0.6 cm.	40	<i>Width.</i> —0.1-0.15 cm.
<i>Texture.</i> —Bifid T-shaped hairs.	40	<i>Shape.</i> —Ovate to lanceolate.
<i>Color of peduncle.</i> —RHS 138A but appears lighter due to hairs.	45	<i>Apex shape.</i> —Acute.
<i>Length of peduncle.</i> —2.2-2.5 cm.	45	<i>Base.</i> —Fused.
<i>Peduncle diameter.</i> —0.1-0.15 cm.	45	<i>Margins.</i> —Entire; papery with RHS 165A edges.
<i>Texture.</i> —Bifid T-shaped hairs.	45	<i>Texture, upper surface.</i> —Glabrous.
<i>Inflorescence:</i>		<i>Lower surface.</i> —Bifid T-shaped hairs.
<i>Type.</i> —Compositae type, solitary inflorescences daisy-type, borne terminally above foliage, ray florets arranged acropetally on a capitulum.	50	<i>Reproductive organs:</i>
<i>Quantity of short days to flowering (response time).</i> —7 weeks.	50	<i>Found on both florets.</i> —Yes.
<i>Quantity of inflorescences per plant.</i> —26-40.	55	<i>Pistil.</i> —1.
<i>Lastingness of individual blooms on the plant.</i> —About 4 weeks.	55	<i>Length.</i> —0.4-0.45 cm.
<i>Fragrance.</i> —Slightly spicy.	55	<i>Style color.</i> —RHS 2C.
<i>Bud (just when opening/showing color):</i>		<i>Style length.</i> —0.3 cm.
<i>Color.</i> —RHS 178A to RHS 178B on upper side of petal; and RHS 175B to RHS 175C on the lower side with RHS 178B at the apex.	60	<i>Stigma color.</i> —RHS 9A.
<i>Length.</i> —0.4-0.5 cm.	60	<i>Stigma shape.</i> —Bi-parted.
<i>Width.</i> —0.7 cm.	65	<i>Ovary color.</i> —Not observed.
<i>Shape.</i> —Oblate.	65	<i>Found on only disc florets.</i> —Yes.
		<i>Stamens.</i> —4.
		<i>Color of filaments.</i> —RHS 2D but more translucent looking.

US PP23,999 P2

**5**

*Length filaments.*—0.2 cm.

*Anther color.*—Closest to RHS 9C.

*Anther length.*—0.1 cm.

*Anther shape.*—Oblong.

*Color of pollen.*—RHS 14B.

*Pollen amount.*—Some.

*Fertility/seed set.*—Has not been observed on this hybrid.

**6**

*Disease/pest resistance.*—Has not been observed on this hybrid.

What is claimed is:

1. A new and distinct variety of *Chrysanthemum* plant named ‘CIDZ0021’ substantially as illustrated and described herein.

\* \* \* \* \*

