



(12) **United States Plant Patent**
Egger

(10) **Patent No.:** **US PP24,036 P3**
(45) **Date of Patent:** **Nov. 19, 2013**

(54) **KNIPHOFIA PLANT NAMED ‘REDHOT
POPSICLE’**

(50) Latin Name: ***Kniphofia* spp.**
Varietal Denomination: **Redhot Popsicle**

(75) Inventor: **Janet N. Egger**, Wilsonville, OR (US)

(73) Assignee: **Terra Nova Nurseries, Inc.**, Canby, OR
(US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 79 days.

(21) Appl. No.: **13/506,430**

(22) Filed: **Apr. 17, 2012**

(65) **Prior Publication Data**

US 2013/0276198 P1 Oct. 17, 2013

(51) **Int. Cl.**
A01H 5/00 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./443**

(58) **Field of Classification Search**
USPC Plt./443
See application file for complete search history.

Primary Examiner — Annette Para

(74) *Attorney, Agent, or Firm* — Klarquist Sparkman, LLP

(57) **ABSTRACT**

A new and distinct *Kniphofia* plant characterized by its
numerous spikes of coral red flowers the first year, repeat
blooming from June through September in Canby, Oreg., a
compact habit with multiple crowns, narrow, grassy leaves,
and excellent vigor.

1 Drawing Sheet

1

Botanical denomination: *Kniphofia* spp.
Cultivar designation: Redhot Popsicle.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct *Kniphofia*,
and given the cultivar name is ‘Redhot Popsicle’. *Kniphofia*
is in the family Asphodelaceae. ‘Redhot Popsicle’ is a
hybrid resulting from a planned breeding program to produce
a series of compact, reblooming *Kniphofia*. The cross was
made using the following proprietary, unreleased plants:
Kniphofia 26-1, as the seed parent, and *Kniphofia* 20-1, as the
pollen parent. It was selected for best habit, flower color,
reblooming, and crown count from many seedlings of the
cross in Canby, Oreg.

Compared to *Kniphofia* ‘Bressingham Comet’ (an unpatented
plant), the new cultivar is more free flowering with red
flowers rather than yellow to red orange flowers.

Compared to *Kniphofia* ‘Nancy’s Red’ (an unpatented
plant), the new cultivar is shorter and much more free flowering.

SUMMARY OF THE INVENTION

The new cultivar is unique and characterized by:

1. numerous spikes of coral red flowers the first year
2. repeat blooming from June through September in Canby,
Oreg.
3. compact habit with multiple crowns
4. narrow, grassy leaves
5. excellent vigor, fills a one gallon pot the first year from
tissue culture

This new cultivar has been reproduced only by asexual
propagation (division, tissue culture). Each of the progeny
exhibits identical characteristics to the new cultivar. Asexual
propagation by leaf cuttings, tissue culture, and division using
standard techniques as done in Canby, Oreg., shows that the
foregoing characteristics and distinctions come true to form
and are established and transmitted through succeeding

2

propagations. The present invention has not been evaluated
under all possible environmental conditions. The phenotype
may vary with variations in environment without a change in
the genotype of the plant.

BRIEF DESCRIPTION OF THE DRAWING

FIG. 1 shows a plant of a one-year-old *Kniphofia* ‘Redhot
Popsicle’ in the ground in the trial field in July in Canby, Oreg.

DETAILED PLANT DESCRIPTION

The following is a detailed description of the new *Kniphofia*
cultivar based on observations of six-month-old specimens
growing in the ground outside in part sun in Canby,
Oreg. Canby is in Zone 8 on the USDA Hardiness map.
Temperatures range from a high of 95 degrees F. in August to
an average of 32 degrees F. in January. Normal rainfall in
Canby is 42.8 inches per year in the trial fields in Canby, Oreg.
The color descriptions are all based on The Royal Horticultural
Society Colour Chart, 5th edition.

Plant:

Type.—Rhizomatous herbaceous perennial.

Hardiness.—USDA Zones 6 to 9.

Size.—Grows to 41 cm wide and 34 cm tall from the top
of the soil to the top of the foliage.

Form.—Upright and clumping.

Leaf:

Type.—Simple.

Shape.—Strap-like.

Arrangement.—Basal rosette.

Blade size.—Grows to 40 cm long and 10 mm wide.

Margins.—Serrulate.

Apex.—Acuminate.

Base.—Clasping.

Surface texture.—Glabrous on both surfaces.

Venation.—Parallel.

Color.—Top and bottom side Green 137A.

Inflorescence:

Type.—Spike-like scapose raceme.

Number of flowers per raceme.—About 100.

Inflorescence.—Grows to 25 cm long and 5 cm wide.

Peduncle description.—Grows to 35 cm long and 7 mm 5
wide, glabrous to glaucous, Yellow Green 146C.

Pedicle description.—2.5 mm long, glabrous, Greyed
Orange 177D.

Bloom time.—Late June through September in Canby, 10
Oreg.

Lastingness.—An inflorescence lasts for 2 to 4 weeks
depending on the temperatures.

Flower Bud:

Size.—26 mm long and 5 mm wide.

Shape.—Cylindrical.

Surface texture.—Glabrous.

Color.—Orange Red N34C.

Flower:

Type.—Actinomorphic.

Shape.—Cylindrical and tubular.

Size.—Grows to 4 cm deep and 12 mm wide.

Texture.—Waxy.

Surface texture.—Glabrous inside and outside.

Color.—Outside, Orange Red N34C, inside Red 35A
with veins Orange Red N34A.

Corolla description.—6 lobed fused tepals, 27 mm long
and 12 mm wide, each lobe 3 mm to 3.5 mm wide and
5 mm long, ovate to oblong, margin entire, tip
notched; glabrous inside and out.

Pistil description.—One extruding, 33 mm long, ovary 3
mm long 1.8 mm wide, Yellow Green 144A, style 30
mm long, Red 36D, stigma White 155A.

Stamen.—6, 34 mm long, filaments 33 mm long, and
Yellow 4D, anthers 1.5 mm long and Yellow 13A,
pollen Yellow 4B.

Fragrance.—None.

15 *Fruit and seed*: None seen.

Pest and diseases: No known resistances to pests or diseases.
No problems have been observed on this plant grown under
commercial conditions in Canby, Oreg.

I claim:

20 1. A new and distinct *Kniphofia* plant as herein shown and
described.

* * * * *

