



US00PP24120P3

(12) **United States Plant Patent**
Kardos(10) **Patent No.:** US PP24,120 P3
(45) **Date of Patent:** Dec. 31, 2013(54) **LANTANA PLANT NAMED ‘SUNSET ORANGE’**(50) Latin Name: *Lantana* sp.
Varietal Denomination: **Sunset Orange**(75) Inventor: **Joshua H. Kardos**, Watkinsville, GA
(US)(73) Assignee: **Plant Introductions, Inc.**, Watkinsville,
GA (US)(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 119 days.(21) Appl. No.: **13/374,634**(22) Filed: **Jan. 4, 2012**(65) **Prior Publication Data**

US 2013/0174307 P1 Jul. 4, 2013

(51) **Int. Cl.**
A01H 5/00 (2006.01)(52) **U.S. Cl.**
USPC **Plt./227**(58) **Field of Classification Search**
USPC Plt./227
See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

PP19,548 P2 12/2008 Dirr
PP21,452 P2 * 11/2010 Pan Plt./227
PP21,539 P3 11/2010 Griffith
2012/0084892 P1 4/2012 Kardos

OTHER PUBLICATIONS

Lantana Catalog Page (distributed at a trade show Jan. 5, 2011)
“*Lantana* ‘Sunset Orange’”, 1 pg.
Plant Introductions, Inc. webpage, (downloaded Dec. 30, 2011)
“*Lantana* selections”, <http://www.plantintroductions.com/lantanaselections.html>, 3 pgs.

* cited by examiner

Primary Examiner — June Hwu

(74) Attorney, Agent, or Firm — Lathrop & Gage LLP

(57) **ABSTRACT**

A new and distinct cultivar of *Lantana* plant named ‘Sunset Orange’, characterized by its low growing, mounding to spreading growth habit; leathery, scabrous dark green foliage; continuous flowering; and light orange flower buds that open to light orange and age to dark orange.

2 Drawing Sheets**1**

Genus and species of plant claimed: *Lantana* sp.
Variety denomination: ‘Sunset Orange’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Lantana* plant, botanically known as *Lantana* sp., and hereinafter referred to by the cultivar name ‘Sunset Orange’.

The new *Lantana* plant originated from a controlled cross between *Lantana* sp. ‘Chapel Hill Gold’ (U.S. Plant Pat. No. 21,539) and an unnamed seedling of *Lantana camara* known to the inventor as *Lantana camara* DIO (not patented) as part of a planned breeding program to develop low growing, continuously flowering *lantanas* with dark green foliage. The cultivar ‘Sunset Orange’ originated and was discovered in a cultivated environment in a greenhouse in Watkinsville, Ga.

Asexual reproduction of the new cultivar by stem cuttings in Watkinsville, Ga. has shown that all the unique features of this new *Lantana*, as herein described, are stable and reproduced true-to-type through successive generations of such asexual propagation.

SUMMARY OF THE INVENTION

Plants of the new cultivar ‘Sunset Orange’ have not been observed under all possible environmental conditions. The phenotype may vary somewhat with changes in light, temperature, soil and rainfall without, however, any variance in genotype.

2

The following traits have been repeatedly observed and are determined to be unique characteristics of ‘Sunset Orange’. These characteristics in combination distinguish ‘Sunset Orange’ as a new and distinct cultivar: 1. Low growing, 5 mounding to spreading growth habit; 2. Leathery, scabrous dark green foliage; 3. Continuous flowering; and 4. Light orange flower buds that open to light orange and age to dark orange. There are no other cultivars of *Lantana* with this combination of characteristics known to the inventor.

10 Plants of the new *Lantana* ‘Sunset Orange’ differ from plants of the female parent, ‘Chapel Hill Gold’, primarily in flower color and growth habit as plants of ‘Chapel Hill Gold’ have bright golden yellow flower buds and flowers and a spreading growth habit, whereas plants of ‘Sunset Orange’ have light orange flower buds that open to light orange and age to dark orange, and a mounding to spreading growth habit. Plants of the new *Lantana* ‘Sunset Orange’ differ from plants of the male parent, an unnamed seedling known to the inventor as *Lantana* DIO, primarily in flower color and 15 growth habit, as plants of the male parent have flowers that remain light orange and a loose, rounded growth habit, whereas plants of ‘Sunset Orange’ have flowers that open light orange and age to dark orange, and an overall more compact, mounding to spreading growth habit.

20 25 ‘Sunset Orange’ can be compared to the cultivar ‘Bani Redda’ (U.S. Plant Pat. No. 21,452), but differs in the growth habit, foliage, and floral characteristics. ‘Sunset Orange’ has a low growing, mounding to spreading and overall larger growth habit, whereas ‘Bani Redda’ has a compact and dense mounding and overall smaller growth habit. ‘Sunset Orange’

has overall smaller foliage than 'Bani Redda'. 'Sunset Orange' averages 23 flowers per inflorescence, whereas 'Bani Redda' averages 33 flowers per inflorescence. 'Sunset Orange' has light orange flowers that mature to dark orange, whereas 'Bani Redda' has yellow flowers that mature to 5 orange-red.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying color photographs illustrate the flower 10 and foliage characteristics and the overall appearance of the new *Lantana*, showing the colors as true as it is reasonably possible to obtain in color reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately 15 describe the colors of the new *Lantana*.

FIG. 1 illustrates a close-up view of the inflorescences of 'Sunset Orange'.

FIG. 2 illustrates the overall appearance of a group of 20 mature plants of 'Sunset Orange' in containers.

DETAILED DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2001 Edition, 25 except where general terms of ordinary dictionary significance are used. Plants used for the description were approximately six months old and were grown in 8 L containers in full sun under outdoor conditions in a nursery in Watkinsville, Ga. Botanical classification: *Lantana* sp., cultivar 'Sunset 30 Orange'.

Parentage:

Female parent.—*Lantana* sp. cultivar 'Chapel Hill Gold' (U.S. Plant Pat. No. 21,539).

Male parent.—*Lantana camara* unnamed seedling 35 known to the inventor as *Lantana camara* DIO (not patented).

Propagation: Stem cuttings.

Time to initiate roots, summer: About 10 days at 32° C.

Plant description: Herbaceous flowering plant, subshrub, low 40 growing, mounding to spreading growth habit. Freely branching; two lateral branches potentially forming at every node; pinching enhances lateral branch development.

Root description.—Numerous, fine, fibrous and well- 45 branched.

Plant size.—About 45 cm in height from the soil level to the top of the inflorescences, and about 60 cm in diameter. Young stems having a diameter of about 3 mm and a squarish shape. Mature stems having a 50 diameter of about 4 mm or more and a rounded shape.

Stem strength.—Strong, but flexible.

Stem texture.—Coarse, pubescent.

Stem color (young).—144A. Color (mature): 199B.

Internode length.—About 2.2 cm.

Vegetative buds: Opposite in arrangement, valvate, ovoid, 55 pubescent.

Color.—144A.

Size.—About 1 mm in length and about 1 mm in width.

Foliage description:

Arrangement.—Opposite, simple.

Length.—About 3 cm.

Width.—About 2.3 cm.

Shape.—Ovate.

Apex.—Acute.

Base.—Cuneate.

Margin.—Crenate.

Texture (upper surface).—Leathery, moderately waxy, with scabrous pubescence.

Texture (lower surface).—Rough, with hispid pubescence.

Venation pattern.—Pinnate.

Venation color (upper surface).—144B.

Venation color (lower surface).—144A.

Fragrance.—Pungent, mint-like.

Color of developing and mature foliage (upper surface).—147A.

Color of developing and mature foliage (lower surface).—147B.

Petiole length.—About 6 mm.

Petiole diameter.—About 2 mm.

Petiole texture, both surfaces.—Hispid pubescence.

Petiole color (upper and lower surfaces).—144B.

Flower description:

Flower type and habit.—Small salverform flowers arranged in axillary corymbs; flowers face mostly upward or outward. Flowers are self-cleaning. Freely flowering with potentially two inflorescences per node; typically about 20 to 25 flowers per corymb.

Natural flowering season.—Spring until the first frost in fall; flowering is continuous.

Flower longevity on the plant.—About one week.

Fragrance.—None observed.

Inflorescence diameter.—About 3.4 cm.

Inflorescence height.—About 1.5 cm.

Flower bud length.—About 1 cm.

Flower bud diameter.—About 3 mm.

Flower bud shape.—Oblong.

Flower bud color.—21A.

Flower appearance.—Flared trumpet, corolla fused, four-parted; flowers roughly rectangular in shape.

Diameter.—About 9 mm.

Corolla tube length.—About 8 mm.

Pedicels.—None observed, flowers not stalked.

Petals:

Arrangement/appearance.—Single whorl of four petals, fused into flared trumpet.

Petal length from throat.—About 5 mm for the upper petal, about 4 mm for the lower petal, and about 3 mm for the lateral petals.

Petal width.—Upper and lower petals are about 6 mm in width, and lateral petals are about 3 mm in width.

Petal shape.—Spatulate to somewhat orbicular.

Petal apex.—Obtuse.

Petal base.—Fused.

Petal margin.—Entire.

Petal texture, upper and lower surfaces.—Smooth, glabrous.

Petal color (young).—Upper surface: 21A and lower surface: 17C.

Color of throat.—21A.

Color of corolla tube.—17C.

Petal color (mature).—Upper surface: N25B and lower surface: 26B.

Color of throat.—N25B.

Color of corolla tube.—21D.

Calyx:

Arrangement/appearance.—The calyx consists of 5 sepals fused into a single tubular structure. Length is about 4.5 mm. Width is about 1.5 mm.

<i>Sepal base.</i> —Fused.		Pistils:
<i>Sepal apex.</i> —Acute.		<i>Quantity.</i> —One inferior pistil per flower.
<i>Texture, upper and lower surfaces.</i> —Scabrous.		<i>Pistil length.</i> —About 4 mm.
<i>Color, upper and lower surfaces.</i> —146A.	5	<i>Stigma shape.</i> —Rounded.
<i>Bracts:</i>		<i>Stigma size.</i> —About 1 mm in diameter.
<i>Arrangement/appearance.</i> —One bract per flower at the base of the corolla, leaf-like. Length is about 6 mm.		<i>Stigma color.</i> —145C.
Width is about 1.5 mm.		<i>Style length.</i> —About 2 mm.
<i>Shape.</i> —Lanceolate.	10	<i>Style color.</i> —145D.
<i>Apex.</i> —Acute.		<i>Ovary size.</i> —About 1 mm in diameter.
<i>Margin.</i> —Entire.		<i>Ovary color.</i> —145B.
<i>Texture, upper and lower surface.</i> —Scabrous.		Fruit:
<i>Color, upper and lower surfaces.</i> —146A.		<i>Type/appearance.</i> —Drupe.
<i>Peduncles:</i>	15	<i>Shape.</i> —Round.
<i>Length.</i> —About 4.5 cm.		<i>Diameter.</i> —About 5 mm.
<i>Diameter.</i> —About 1.5 mm.		<i>Mature color.</i> —202A.
<i>Angle.</i> —About 45 degrees from the stem.		<i>Number per infructescence.</i> —Ranges from zero to about five.
<i>Strength.</i> —Flexible, but strong.	20	Disease/pest resistance: Plants of the claimed <i>Lantana</i> variety grown in the garden have not been noted to be susceptible or resistant to pathogens and pests common to <i>Lantana</i> .
<i>Color.</i> —144A.		Weather and temperature tolerance: Plants of the new variety have been observed to be cold hardy to U.S. Cold Hardiness Zone 8.
<i>Stamens:</i>	25	I claim:
<i>Quantity/arrangement.</i> —Four per flower, adnate to the inside of the corolla tube.		1. A new and distinct <i>Lantana</i> plant named ‘Sunset Orange’, as illustrated and described herein.
<i>Anther shape.</i> —Oblong.		* * * * *
<i>Anther length.</i> —1 mm.		
<i>Anther width.</i> —Less than 1 mm.		
<i>Anther color.</i> —13A.		
<i>Pollen amount.</i> —Produced in very small quantities, and 13A in color.		

U.S. Patent

Dec. 31, 2013

Sheet 1 of 2

US PP24,120 P3



FIGURE 1



FIGURE 2