



US00PP24203P2

(12) **United States Plant Patent**
Jacobs(10) **Patent No.:** US PP24,203 P2
(45) **Date of Patent:** Jan. 28, 2014

- (54) **ALSTROEMERIA PLANT NAMED 'ZALSARENO'**
- (50) Latin Name: *Alstroemeria hybrida*
Varietal Denomination: Zalsareno
- (75) Inventor: **Henricus Cornelius Maria Jacobs,**
Rijsenhout (NL)
- (73) Assignee: **Van Zanten Plants b.v.**, Hillegom (NL)
- (*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 92 days.
- (21) Appl. No.: **13/385,871**
- (22) Filed: **Mar. 10, 2012**

- (51) **Int. Cl.**
A01H 5/00 (2006.01)
- (52) **U.S. Cl.**
USPC **Plt./309**
- (58) **Field of Classification Search**
USPC Plt./309
See application file for complete search history.

Primary Examiner — Susan McCormick Ewoldt
(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Alstroemeria* plant named 'Zalsareno', characterized by its erect and strong flowering stems; vigorous growth habit; purple and pale yellow-colored flowers; and excellent postproduction longevity.

1 Drawing Sheet

1

Botanical designation: *Alstroemeria hybrida*.
Cultivar denomination: 'ZALSARENO'.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Alstroemeria* plant, botanically known as *Alstroemeria hybrida*, commercially used as a cut flower *Alstroemeria*, and hereinafter referred to by the name 'Zalsareno'.

The new *Alstroemeria* plant is a product of a planned breeding program conducted by the Inventor in Rijsenhout, The Netherlands. The objective of the breeding program is to create new cut flower *Alstroemeria* plants with desirable flower and plant qualities, attractive flower coloration and excellent postproduction longevity.

The new *Alstroemeria* plant originated from a cross-pollination made by the Inventor in Rijsenhout, The Netherlands in June, 2006 of a proprietary *Alstroemeria hybrida* selection identified as code number 2Z171-1, not patented, as the female, or seed, parent with a proprietary *Alstroemeria hybrida* selection identified as code number 871069-2, not patented, as the male, or pollen, parent. The new *Alstroemeria* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Rijsenhout, The Netherlands in July, 2007.

Asexual reproduction of the new *Alstroemeria* plant by rhizome divisions in a controlled greenhouse environment in Rijsenhout, The Netherlands since September, 2007 has shown that the unique features of this new *Alstroemeria* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Alstroemeria* have not been observed under all possible environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environmental conditions such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Zalsareno'.

2

These characteristics in combination distinguish 'Zalsareno' as a new and distinct *Alstroemeria* plant:

1. Erect and strong flowering stems.
2. Vigorous growth habit.
3. Purple and pale yellow-colored flowers.
4. Excellent postproduction longevity.

Plants of the new *Alstroemeria* can be compared to plants of the parent selections. Plants of the new *Alstroemeria* differ from plants of the parent selections primarily in flower color as plants of the female parent selection have dark purple-colored flowers and plants of the male parent selection have yellow-colored flowers.

Plants of the new *Alstroemeria* can be compared to plants of *Alstroemeria hybrida* 'Stabec', disclosed in U.S. Plant Pat. No. 9,041. In side-by-side comparisons conducted in Rijsenhout, The Netherlands, plants of the new *Alstroemeria* differed primarily from plants of 'Stabec' in flower color as plants of 'Stabec' had light red to deep pink-colored flowers.

20 BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Alstroemeria* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Alstroemeria* plant.

30 The photograph comprises a close-up view of a typical flowering stem of 'Zalsareno'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph and following observations and measurements describe plants of the new *Alstroemeria* grown during the late summer and early autumn in ground beds in a glass-covered greenhouse in Rijsenhout, The Netherlands. During the production of the plants, day temperatures ranged from 15° C. to 25° C., night temperatures ranged from 10° C. to 15° C., soil temperatures averaged 15° C. and light levels averaged 5,000 lux. Plants were one year old when the photograph and description were taken. In

the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Alstroemeria hybrida* 'Zalsareno'. 5

Parentage:

Female, or seed, parent.—Proprietary *Alstroemeria hybrida* selection identified as code number 2Z171-1, not patented.

Male, or pollen, parent.—Proprietary *Alstroemeria hybrida* selection identified as code number 871069-2, not patented. 10

Propagation:

Type.—In vitro rhizogenesis.

Time to produce a rooted young plant, summer.—About 15 days at 16° C. to 25° C.

Time to produce a rooted young plant, winter.—About 60 days at 16° C. to 20° C.

Root description.—Fibrous, fleshy, thick; color, close to 155D. 20

Rooting habit.—Freely branching; medium density.

Rhizomes.—Shape: Elongate; rounded. Length: About 10 cm to 30 cm. Diameter: About 3 mm to 10 mm.

Texture: Smooth. Color: Close to 155D.

Plant description:

Plant and growth habit.—Upright; freely branching, bushy appearance; vigorous growth habit.

Plant height.—About 125 cm to 146 cm.

Plant diameter (spread).—About 25 cm.

Flowering stem description:

Aspect.—Erect.

Length.—About 110 cm to 130 cm.

Diameter.—About 6 mm to 8 mm.

Internode length.—About 0.5 cm to 12 cm.

Strength.—Strong.

Texture.—Smooth, glabrous.

Color.—Close to 144B. 35

Foliage description:

Appearance.—Leaves asymmetrical; sessile.

Length.—About 17.4 cm to 21.4 cm.

Width.—About 3.8 cm to 4.8 cm.

Shape.—Lanceolate.

Apex.—Acute.

Base.—Cuneate.

Margin.—Entire; slightly undulate.

Texture, upper and lower surfaces.—Smooth, glabrous. 45

Venation pattern.—Parallel.

Color.—Developing and fully developed leaves, upper surface: Close to 137A; venation, close to 139C. Developing and fully developed leaves, lower surface: Close to 137B; venation, close to 146C. 50

Flower description:

Flower type and habit.—Single cup-shaped flowers arranged in compound umbels; flowers face mostly outwardly; perianth segments separate; freely flowering habit, about seven to eleven flowers developing per flowering stem. 55

Natural flowering season.—Flowering continuous during the spring in The Netherlands; plants begin flowering about 80 to 90 days after planting.

Fragrance.—Not detected.

Flower longevity on the plant.—About four weeks; 60 flowers not persistent.

Flower longevity as a cut flower.—About 12 to 16 days; flowers not persistent.

Flower buds (showing color).—Length: About 3.5 cm to 4.5 cm. Diameter: About 1.2 cm to 1.5 cm. Shape: Roughly ovoid. Color: Close to 61A.

Umbel height.—About 12.5 cm to 16 cm.

Umbel diameter.—About 15 cm to 18 cm.

Flower diameter.—About 6 cm by 6.5 cm.

Flower depth.—About 7 cm to 7.5 cm.

Perianth.—Arrangement: Six arranged in two whorls, each whorl with two lateral and one median segments. Outer perianth, lateral segments: Length: About 5.8 cm to 6.1 cm. Width: About 3.4 cm to 3.9 cm. Shape: Obovate. Apex: Embedded pointed. Base: Attenuate. Margin: Entire; undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close 72A. Color, when opening and fully opened, lower surface: Close to 72A. Outer perianth, median segment: Length: About 5.8 cm to 6.4 cm. Width: About 3.4 cm to 3.8 cm. Shape: Obovate. Apex: Embedded pointed. Base: Attenuate. Margin: Entire; undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 72A. Color, when opening and fully opened, lower surface: Close to 72A. Inner perianth, lateral segments: Length: About 6 cm to 6.3 cm. Width: About 1.8 cm to 2 cm. Shape: Elliptical. Apex: Wishbone-shaped. Base: Attenuate. Margin: Entire; undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 70C; towards the base, close to 4B; at the base, close to 14B; stripes, close to 183A. Color, when opening and fully opened, lower surface: Close to 70C; towards the base, close to 4B; at the base, close to 14B; stripes, close to 183A. Inner perianth, median segment: Length: About 5.3 cm to 5.7 cm. Width: About 1.8 cm to 2 cm. Shape: Elliptical. Apex: Wishbone-shaped. Base: Attenuate. Margin: Entire; undulate. Texture, upper and lower surfaces: Smooth, glabrous. Color, when opening and fully opened, upper surface: Close to 72A; stripes, close to 183A. Color, when opening and fully opened, lower surface: Close to 72A; stripes, close to 183A.

Pedicels.—Length: About 1 cm to 8.5 cm. Diameter: About 2 mm to 3 mm. Strength: Strong. Angle: About 20° from vertical. Texture: Smooth, glabrous. Color, upper and lower surfaces: Close to 144A.

Reproductive organs.—Stamens: Quantity per flower: Six. Anther shape: Elliptical. Anther length: About 9 mm to 10 mm. Anther color: Close to 178A. Pollen amount: Scarce to moderate. Pollen color: Close to 199A. Pistils: Quantity per flower: One. Style length: About 3 cm to 3.5 cm. Style color: Close to 64A. Stigma color: Close to 64A. Ovary color: Close to 144A.

Seeds and fruits.—Seed and fruit development has not been observed on plants of the new *Alstroemeria*.

Disease & pest resistance: Plants of the new *Alstroemeria* have not been observed to be resistant to pathogens and pests common to *Alstroemeria* plants.

Temperature tolerance: Plants of the new *Alstroemeria* have been observed to tolerate temperatures from about 0° C. to about 40° C.

It is claimed:

1. A new and distinct *Alstroemeria* plant named 'Zalsareno' as illustrated and described.

