



US00PP24325P2

(12) **United States Plant Patent**  
**Klemm et al.**

(10) **Patent No.:** **US PP24,325 P2**  
(45) **Date of Patent:** **Mar. 18, 2014**

(54) **OSTEOSPERMUM PLANT NAMED**  
**'KLEOE12201'**

(50) Latin Name: *Osteospermum ecklonis* Norl.  
Varietal Denomination: **KLEOE12201**

(75) Inventors: **Nils Klemm**, Stuttgart (DE); **Silke**  
**Wagner**, Leinfelden-Echterdingen (DE)

(73) Assignee: **Klemm+Sohn GmbH & Co. KG**,  
Stuttgart (DE)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/573,286**

(22) Filed: **Sep. 7, 2012**

(51) **Int. Cl.**  
*A01H 5/00* (2006.01)

(52) **U.S. Cl.**  
USPC ..... **Plt./360**

(58) **Field of Classification Search**  
USPC ..... **Plt./360**  
See application file for complete search history.

*Primary Examiner* — Kent L Bell

(74) *Attorney, Agent, or Firm* — Jondle Plant Sciences  
Division of Swanson & Bratschun, L.L.C.

(57) **ABSTRACT**

A new and distinct *Osteospermum* variety named  
'KLEOE12201' particularly distinguished by light green foli-  
age, good branching, and compact growth habit, is disclosed.

**1 Drawing Sheet**

**1**

Genus and species: *Osteospermum ecklonis* Norl.  
Variety denomination: 'KLEOE12201'.

**BACKGROUND OF THE NEW PLANT**

The present invention comprises a new and distinct variety  
of *Osteospermum*, botanically known as *Osteospermum eck-*  
*lonis* Norl., and hereinafter referred to by the variety name  
'KLEOE12201'. The new variety originated from a con-  
trolled cross conducted in July 2008 in Nairobi, Kenya  
between female *Osteospermum* plant named '07 183' (unpat-  
ented), and male *Osteospermum* plant named 'KLEO04113'  
(EU CPVR No. 21047). A single plant selection was subse-  
quently chosen for further evaluation and asexual propaga-  
tion.

The new variety was first propagated via vegetative cut-  
tings in the spring of 2009 in Stuttgart, Germany, and has been  
asexually reproduced repeatedly by vegetative cuttings for  
three to four generations. The present invention has been  
found to retain its distinctive characteristics through succes-  
sive asexual propagations via vegetative cuttings.

Plant Breeder's Rights for this variety have not been  
applied for. 'KLEOE12201' has not been made publicly  
available or sold more than one year prior to the filing date of  
this application.

**SUMMARY OF THE INVENTION**

The following are the most outstanding and distinguishing  
characteristics of the new variety when grown under normal  
horticultural practices in Stuttgart, Germany.

1. Light green foliage;
2. Good branching; and
3. Compact growth habit.

**DESCRIPTION OF THE PHOTOGRAPH**

This new *Osteospermum* plant is illustrated by the accom-  
panying photograph which shows overall plant habit includ-  
ing blooms, buds, and foliage of the plant in full color. The  
colors shown are as true as can be reasonably obtained by

**2**

conventional photographic procedures. The photograph was  
taken in the spring of 2012 on a 3-month old plant grown in a  
heated glass greenhouse in Stuttgart, Germany, under condi-  
tions which approximate those generally used in normal hor-  
ticultural practice.

**DESCRIPTION OF THE NEW VARIETY**

The following detailed descriptions set forth the distinctive  
characteristics of 'KLEOE12201'. The data which define  
these characteristics were collected from asexual reproduc-  
tions carried out in Stuttgart, Germany. The plant history was  
taken in the spring of 2012 on 5-month old plants grown in 12  
centimeter pots. The plants were grown in a heated glass  
greenhouse and pinched at week 32. Color readings were  
taken under natural and artificial light. Color references are  
primarily to The R.H.S. Colour Chart of The Royal Horticul-  
tural Society of London (R.H.S.), Fifth Edition (2007).

**DETAILED BOTANICAL DESCRIPTION OF THE  
NEW PLANT**

**Classification:**

*Family.*—Asteraceae.  
*Botanical name.*—*Osteospermum ecklonis* Norl.  
*Common name.*—African daisy.  
*Denomination.*—'KLEOE12201'.

**Parentage:**

*Female parent.*—*Osteospermum* plant named '07 183'  
(unpatented).  
*Male parent.*—*Osteospermum* plant named  
'KLEO04113' (EU CPVR No. 21047).

**Plant:**

*Form and habit.*—Herbaceous perennial, usually culti-  
vated as an annual.  
*Growth habit.*—Vigorous.  
*Branching habit.*—Freely branching.  
*Height (from top of soil).*—17.0 cm.  
*Width (including inflorescence).*—28.0 cm.  
*Propagation.*—Vegetative cuttings from tips.

*Time to produce a finished flowering plant.*—13 to 16 weeks.

*Time to initiate and develop roots.*—2 to 4 weeks.

*Root description.*—Freely rooting.

Leaves:

*Arrangement.*—Arranged in a whorl.

*Shape.*—Lanceolate.

*Apex.*—Acute.

*Base.*—Acute.

*Margin.*—Serrate.

*Color (both immature and mature leaves).*—Upper surface: RHS 141A. Lower surface: RHS 138A.

*Length.*—7.0 cm.

*Width.*—3.0 cm.

*Texture.*—Leathery.

*Petiole.*—Absent.

Stems:

*Total number of branches.*—15.

*Length.*—13.0 cm.

*Diameter.*—0.5 cm.

*Internode length.*—1.0 cm.

*Color.*—RHS 141D.

*Texture.*—Smooth.

Inflorescence buds:

*Shape.*—Obovate.

*Length.*—1.2 cm.

*Diameter.*—0.7 cm.

*Color (at tight bud just before the ray florets unfold).*—RHS 137C.

Inflorescence:

*Type.*—Single.

*Blooming habit.*—Spring to autumn.

*Quantity of inflorescences per plant.*—40.

*Lastingness of the inflorescences on the plant.*—12 days.

*Fragrance.*—Absent.

*Inflorescence diameter.*—6.0 cm.

*Disc diameter.*—1.1 cm.

Disc florets:

*Quantity per inflorescence.*—80.

*Shape.*—Oblanceolate.

*Tube color.*—Closed: RHS N79B. Mature: RHS N79C.

*Length.*—0.8 cm.

*Diameter (at apex).*—0.2 cm.

*Apex.*—Obtuse.

*Apex color.*—RHS N79B.

*Base.*—Fused to form a tube.

*Margin.*—Entire.

Ray florets:

*Quantity per inflorescence.*—16.

*Shape.*—Oblanceolate.

*Color.*—Upper surface: RHS N79B. Lower surface: RHS N77B.

*Length.*—3.0 cm.

*Width.*—0.9 cm.

*Apex.*—Obtuse.

*Base.*—Acute.

*Margin.*—Entire.

*Texture.*—Smooth.

Peduncle:

*Length.*—5.0 cm.

*Diameter.*—0.15 cm.

*Texture.*—Rough.

*Color.*—RHS 138A.

Phyllaries:

*Arrangement.*—Single.

*Observed quantity per inflorescence.*—22.

*Shape.*—Lanceolate.

*Color.*—Upper surface: RHS 138A. Lower surface: RHS 138B.

*Length.*—1.1 cm.

*Width.*—0.15 cm.

*Apex.*—Acute.

*Base.*—Obtuse.

*Margin.*—Entire.

*Texture.*—Rough.

Reproductive organs:

*Androecium.*—Location: Base of the disc floret corolla. Stamens: Quantity: 80. Shape: Filamentous. Color: RHS N77A. Filament length: 0.7 cm. Filament diameter: 0.05 cm. Anther: Shape: Elliptical. Color: RHS 200A. Length: 0.3 cm. Diameter: 0.09 cm. Pollen: Color: RHS 13A. Amount: Abundant.

*Gynoecium.*—Location: Base of the ray florets; stigmas in disc florets are degenerated and non-functional. Pistils: Number: 16. Length: 0.8 cm. Diameter: 0.03 cm. Stigma: Color: RHS N79A. Shape: Bipartite. Length: 0.2 cm. Diameter: 0.04 cm. Style: Color: RHS 79C. Length: 0.3 cm. Diameter: 0.03 cm. Shape: Filamentous.

Fruit and seed set: None observed.

Disease and insect/pest resistance: No specific observations were made.

#### COMPARISON WITH PARENTAL AND COMMERCIAL VARIETIES

'KLEOE12201' differs from the female parent *Osteospermum* plant '07 183' (unpatented) in that 'KLEOE12201' has good branching, whereas '07 183' has average branching.

'KLEOE12201' differs from the male parent *Osteospermum* plant 'KLEO04113' (EU CPVR No. 21047) in that 'KLEOE12201' has a compact growth habit and very good basal branching, whereas 'KLEO04113' has an upright growth habit and minimal branching.

'KLEOE12201' differs from the commercial *Osteospermum* plant 'KLEO04110' (U.S. Plant Pat. No. 18,272) in that 'KLEOE12201' has purple flowers, whereas 'KLEO04110' has violet purple flowers.

We claim:

1. A new and distinct variety of *Osteospermum* plant named 'KLEOE12201' as shown and described herein.

\* \* \* \* \*



